

# SINGULAR PERTURBATION OF POLYNOMIAL POTENTIALS AND REAL SPECTRAL LOCI

Alexandre Eremenko and Andrei Gabrielov

[www.math.purdue.edu/~eremenko](http://www.math.purdue.edu/~eremenko)

[www.math.purdue.edu/~agabriel](http://www.math.purdue.edu/~agabriel)

Purdue University

2010

## 0. Brief history

Bender and Wu (1969) studied the even anharmonic oscillator

$$-w'' + (\epsilon x^4 + x^2)w = \lambda w, \quad w(\pm\infty) = 0 \quad (1)$$

as a perturbation of the harmonic oscillator ( $\epsilon = 0$ ). Eigenvalues are analytic functions of  $\epsilon > 0$  but have essential singularity at  $\epsilon = 0$ .

Study of such singularities led Bender and Wu to consideration of *complex potentials* and *boundary conditions in the complex plane*.

Eigenvalues of Problem (1), as functions of complex  $\epsilon$ , have only algebraic singularities for  $\epsilon \neq 0$ , while  $\epsilon = 0$  is a complicated non-isolated singularity. (Simon (1970), Loeffel and Martin (1972), Delabaere, Dillinger and Pham (1997), Eremenko and Gabrielov (2009). Further we refer to this last paper as EG09.)

## 1. Eigenvalue problem:

$$-w'' + P(z)w = \lambda w, \quad (2)$$

$$w(z) \rightarrow 0 \text{ as } z \rightarrow \infty \text{ on } L_1 \text{ and } L_2 \quad (3)$$

where  $P(z) = a_d z^d + \dots + a_1 z$ , and  $L_k = \{r e^{i\theta_k}, r > 0\}$ .

*Separation rays*

$$\operatorname{Re} \left( \int_0^z \sqrt{a_d \zeta^d} d\zeta \right) = 0, \text{ that is } a_d z^{d+2} < 0,$$

divide the plane into  $d+2$  sectors  $S_0, \dots, S_{d+1}$ .  
Solution  $w \neq 0$  of (2) is *subdominant* in  $S_j$  if

$$w(z) \rightarrow 0, \quad z \rightarrow \infty, \quad z \in S_j.$$

For each  $j$ , the space of subdominant in  $S_j$  solutions is 1-dimensional, and no solution can be subdominant in adjacent sectors.

**Definition.** The rays  $L_1, L_2$  are *admissible* for  $P$  if they are not parallel to any separation rays and belong to non-adjacent sectors  $S_j$ .

The spectrum of this problem with admissible  $L_1, L_2$  is discrete and infinite. If  $a_d = 1$  and  $\mathbf{a} = (a_1, \dots, a_{d-1})$  then there exists an entire function  $F$ , called the *spectral determinant*, such that the spectrum is given by the equation

$$F(\mathbf{a}, \lambda) = 0.$$

The set of all solutions of this equation in the  $(\mathbf{a}, \lambda)$  space is called the *spectral locus*.

We study global topology of the spectral locus. For example:

*For every  $d \geq 3$  the spectral locus is a smooth connected hypersurface in  $\mathbb{C}^d$ . (Alexandersson and Gabrielov (2010); case  $d = 3$ : EG09).*

*For  $d = 4$ , the spectral locus of even potentials  $P$  consists of two disjoint smooth connected curves in  $\mathbb{C}^2$  (EG09).*

## 2. Self-adjoint and $PT$ -symmetric problems

If  $P$  is real and  $L_1, L_2 \subset \mathbf{R}$ , the problem is self-adjoint and the spectrum is real.

If  $P(-\bar{z}) = \overline{P(z)}$ , and the rays  $L_1, L_2$  are *interchanged* by the reflection in  $i\mathbf{R}$ , the problem is called  $PT$ -symmetric.\* In this case, the spectral determinant is a *real entire function* but some eigenvalues may be non-real.

For the  $PT$ -symmetric cubic potential

$$P(z) = iz^3 + iaz \quad \text{and} \quad L_1, L_2 \subset \mathbf{R}, \quad (4)$$

the spectrum is real if  $a \geq 0$  (Case  $a = 0$ : Dorey, Dunning, Tateo (2001); general case: Shin (2002)).

\* $P$  in  $PT$  stands for parity and  $T$  for time. For mathematics, it does not matter which reflection to consider. It is important that the potential and the boundary conditions are preserved by the symmetry.

The following computer-generated plot of the real spectral locus of (4) is taken from Trinh's thesis (2002). One of our goals is to *prove* that the spectral locus really looks like this.




Fig 1. Real spectral locus for  $PT$ -symmetric cubic.

Our method is based on “Nevanlinna parametrization” of the spectral locus introduced in EG09, and “degeneration” (singular perturbation) of potentials.

Degeneration results show what happens when  $a \rightarrow \infty$  while  $a_d$  is fixed. By rescaling, this is equivalent to  $a_d \rightarrow 0$ , while  $a$  is bounded. So we consider potentials

$$P_t(z) = tz^d + cz^m + p_t(z) \quad (5)$$

where  $m < d$ ,  $c \in \mathbf{C} \setminus \{0\}$  is fixed,  $\deg p_t < m$ , coefficients of  $p_t$  are bounded, and  $t \searrow 0$ .

We’ll give sufficient conditions for the spectrum of  $P_t$  to converge to the spectrum of  $P_0$  as  $t \searrow 0$ .

First we study the model case  $p_t = 0$ , and then extend the results to the general case.

### 3. Stokes complexes of the binomials

The asymptotic behavior of solutions of the equation  $-w'' + Pw = \lambda w$  depends on

$$\int \sqrt{P(z) - \lambda} \, dz,$$

which leads to the question of the structure of trajectories of the quadratic differential

$$Q(z)dz^2, \quad Q = P - \lambda.$$

The zeros of  $Q$  are called *turning points*. Curves where  $Q(z)dz^2 < 0$  are called *vertical trajectories* and curves where  $Q(z)dz^2 > 0$  *horizontal trajectories*. Vertical (horizontal) trajectories adjacent to the turning points are called the *Stokes lines* (*anti-Stokes lines*).



Stokes lines and turning points form the 1-skeleton of the cell decomposition of the plane which is called the *Stokes complex*. The 2-cells of this decomposition are called *faces*. The multi-valued function

$$\int \sqrt{Q(z)} dz$$

splits into single-valued branches in the faces. Each branch maps its face onto a right half-plane, or a left half-plane, or onto a vertical strip.

## Examples of Stokes complexes




Fig 2. Stokes complex of  $z^4 + iz^3$ .




Fig 3. Stokes complex of  $z^4 + e^{\pi i/4} z^3$ .

Let  $Q(z) = z^d + cz^m$ . Consider the partition  $T$  of the plane into sectors by the Stokes lines of monomials  $z^d dz^2$  and  $cz^m dz^2$  (i.e., by the rays where  $z^{d+2} < 0$  or  $cz^{m+2} < 0$ ).

**Definition.** A sector  $S$  of  $T$  is called *stable* if

(a)  $S$  contains an anti-Stokes line of  $cz^m dz^2$ ,  
and

(b) The closures of the two sectors of  $T$  adjacent to  $S$  do not contain non-zero turning points (roots of  $z^{d-m} = -c$ ).

A sector  $S$  is *marginally stable* if it satisfies (a) and this weaker condition instead of (b):

(b') The interiors of the two sectors adjacent to  $S$  do not contain turning points.




Fig 4. Partition  $T$  for (a)  $Q = z^4 + iz^3$   
and (b)  $Q = z^4 + e^{i\pi/4}z^3$ .

Black solid lines are the Stokes lines of  $z^4 dz^2$ . Red dashed lines are the Stokes lines of the cubic monomial. Dotted lines are anti-Stokes lines of the two monomials. Sectors  $S_1 - S_4$  in (a) and  $S_0 - S_3$  in (b) are stable.

**Theorem 1.** *Marginally stable sectors do not intersect the Stokes complex of  $z^d + cz^m$ .*

*Stable sectors do not intersect the Stokes complex of  $P_t$  in  $\{z : |z| > R\}$  where  $R$  depends on  $c$  and the bounds for the coefficients of  $p_t$ , but does not depend on  $t \geq 0$ .*

**Theorem 2.** *Suppose that the rays  $L_1, L_2$  are admissible for  $z^d$  and  $cz^m$ , and belong to stable sectors of  $T$ . Then the spectrum of the boundary value problem for the potential  $P_t$  with boundary conditions on  $L_1$  and  $L_2$  converges to the spectrum of the boundary value problem for  $P_0$  with the same boundary conditions.*

Convergence of the spectra means that the spectral determinants converge uniformly with respect to  $\lambda$  and coefficients of  $p_t$ , when  $\lambda$  and these coefficients are restricted to a compact set.

In other words, for every eigenvalue  $\lambda_0 \in K$  for  $t = 0$ , there exists a unique eigenvalue  $\lambda_t$  which converges to  $\lambda_0$  as  $t \rightarrow 0$ , and this convergence is uniform with respect to coefficients of  $p_t$  restricted to a compact set.

Our proof of Theorem 2 uses a conformal change of the independent variable

$$\zeta = \int \sqrt{P - \lambda} \, dz$$

which is due to Green and Liouville. Theorem 1 ensures that this change of the variable behaves continuously at  $t = 0$ .

In our applications to spectral loci we need the following Stokes complex:




Fig 5. Stokes complex of  $z^3 + z^2$ . Sectors intersecting the imaginary axis are stable.




Fig 6. Stokes complex of  $z^6 + z$ . There are no stable sectors.



## 4. Real spectral locus for the cubic

$$-w'' + (z^3 - az + \lambda) = 0, \quad w(\pm i\infty) = 0. \quad (6)$$

This problem is equivalent to the  $PT$ -symmetric problem with

$$P(z) = iz^3 + iaz \quad \text{and} \quad L_1, L_2 \subset \mathbf{R}$$

by the change of the variable  $z \mapsto iz$ .

**Theorem 3.** *For every integer  $n \geq 0$ , there exists a simple curve  $\Gamma_n \subset \mathbf{R}^2$ , which is the image of a proper analytic embedding of a line, and which has these properties:*

- (i) *For every  $(a, \lambda) \in \Gamma_n$  problem (6) has an eigenfunction with  $2n$  non-real zeros.*
- (ii) *The curves  $\Gamma_n$  are disjoint and the real spectral locus of (6) is  $\bigcup_{n \geq 0} \Gamma_n$*
- (iii) *The map*

$$\Gamma_n \cap \{(a, \lambda) : a \geq 0\} \rightarrow \mathbf{R}_{\geq 0},$$

*$(a, \lambda) \mapsto a$  is a 2-to-1 covering.*

- (iv) *For  $a \geq 0$ ,  $(a, \lambda) \in \Gamma_n$  and  $(a, \lambda') \in \Gamma_{n+1}$  imply  $\lambda' > \lambda$ .*

## Sketch of the proof.

a) Nevanlinna parametrization of the real spectral locus. Consider the following cell decomposition  $\Phi$  of the Riemann sphere, with labeled faces. Here  $b = e^{i\beta}$ ,  $\beta \in (0, \pi)$ .




Fig 7. Cell decomposition  $\Phi$  of the sphere.

and the following cell decompositions  $\Psi_n$  of the complex plane:




Fig 8. Cell decomposition  $\Psi_2$  of the plane.

Labeled cell decomposition  $\Psi_n$  have the same local structure as  $\Phi$ , so there exists a local homeomorphism  $g : \mathbf{C} \rightarrow \overline{C}$  such that  $\Psi_n = g^{-1}(\Phi)$ . This  $g$  can be chosen so that it commutes with the reflection  $z \mapsto \bar{z}$ . By the Uniformization Theorem, there exists a symmetric homeomorphism  $\phi : \mathbf{C} \rightarrow \mathbf{C}$  such that  $f = g \circ \phi$  is a real meromorphic function. According to Nevanlinna theory, this function satisfies the differential equation

$$\frac{f'''}{f'} - \frac{3}{2} \left( \frac{f''}{f'} \right)^2 = -2(z^3 - az + \lambda)$$

with real  $a, \lambda$ . Now we have  $f = w/w_1$  where  $w, w_1$  are two real linearly independent solutions of the equation

$$-w'' + (z^3 - az + \lambda)w = 0,$$

and from our construction follows that  $w$  is subdominant in the sectors intersecting the imaginary axis. So  $\lambda$  is an eigenvalue. The eigenfunction  $w$  has  $2n$  non-real zeros, by construction. Thus we have a curve  $\Gamma_n$  in the real spectral locus, parametrized by  $b = e^{i\beta}$ ,  $\beta \in (0, \pi)$ .

That the union of these curves exhaust the whole real spectral locus follows by reversing the steps, and applying the classification of cell decompositions from EG09a. This proves (i) and (ii). (iii) follows from a result of Shin that for  $a > 0$  all eigenvalues are real. To show (iv) we apply rescaling and our previous results on the continuous behavior of the spectrum under degeneration  $a \rightarrow +\infty$ . Real affine change of the independent variable gives an equation  $-y'' + (tz^3 + z^2 + \mu)y = 0$ , where  $\mu$  is an explicit increasing function of  $\lambda$ . As  $t \searrow 0$  this tends to a self-adjoint problem (harmonic oscillator). The convergence of spectra is justified using Theorems 1 and 2, see Fig. 5. Then the Sturm–Liouville theory gives the relation between the order of eigenvalues and number of zeros of eigenfunctions.

It remains to verify that the number of non-real zeros of an eigenfunction does not change in this degeneration. For this we consider the degeneration of the cell decomposition  $\Psi_n$ :




Fig 9. Degenerated cell decomposition of the plane: the loops around  $b$  and  $\bar{b}$  are replaced by a single loop.

## 5. Real spectral locus for a $PT$ -symmetric quartic family

This is a 2-parametric family

$$-w'' + (-z^4 + az^2 + cz)w = -\lambda w, \quad w(\pm i\infty) = 0. \quad (7)$$

It is equivalent to the  $PT$ -symmetric family

$$-w'' + (z^4 + az^2 + icz)w = \lambda w, \quad w(\pm\infty) = 0,$$

studied by Bender, *et al* (2001) and Delabaere and Pham (1998).

**Theorem 4.** *The real spectral locus of (7) consists of disjoint smooth analytic properly embedded surfaces  $S_n \subset \mathbf{R}^3$ ,  $n \geq 0$ , homeomorphic to a punctured disk. For  $(a, c, \lambda) \in S_n$ , the eigenfunction has exactly  $2n$  non-real zeros. For large  $a$ , projection of  $S_n$  on the  $(a, c)$  plane approximates the region  $9c^2 - 4a^3 \leq 0$ .*

Numerical computation suggests that the surfaces have the shape of infinite funnels with the sharp end stretching towards  $a = -\infty$ ,  $c = 0$ , and that the section of  $S_n$  by every plane  $a = a_0$  is a closed curve.

Theorem 4 implies that this section is compact for large  $a_0$ .

The following computer-generated plot is taken from Trinh's thesis:




Fig 10. Section of the surfaces  $S_0, \dots, S_3$  by the plane  $a = -9$ .



Proof of Theorem 4 follows the same lines as the proof of Theorem 3. The cell decompositions now look like this:




Fig 11. Cell decomposition  $\Phi$  of the sphere for Theorem 4.

Here the Nevanlinna parameter  $b$  belongs to the upper half-plane punctured at  $i$ , which explains why  $S_n$  is doubly connected.




Fig 12. Cell decomposition  $\Psi_2$  of the plane for Theorem 4.

To study asymptotics of  $S_n$  as  $a$  and  $c$  tend to infinity, after an affine change of the independent variable we obtain

$$-y'' + (-tz^4 + z^3 + \alpha z)y = -\mu y.$$

Theorems 1 and 2 imply that the spectrum changes continuously at  $t = 0$ .




Fig 13. Stokes complex of  $-z^4 + z^3$ . Sectors intersecting the imaginary axis are stable.




Fig 14. Degenerated cell decomposition for Theorem 4.

The rescaled potential converges to the previously studied cubic potential, and we can make conclusions about asymptotic behavior of the spectral locus.

## References

P. Alexandersson and A. Gabrielov, On eigenvalues of the Schrödinger operator with a polynomial potential with complex coefficients, in preparation.

C. Bender, M. Berry, P. Meisinger, Van M. Savage, M. Simsek, Complex WKB analysis of energy-level degeneracies of non-Hermitian Hamiltonians. J. Phys. A 34 (2001), no. 6, L31–L36.

C. Bender, A. Turbiner, Analytic continuation of eigenvalue problems. Phys. Lett. A 173 (1993), no. 6, 442–446.

C. Bender and Tai Tsun Wu, Anharmonic oscillator. Phys. Rev. (2) 184 1969 1231–1260.

E. Delabaere, F. Pham, Eigenvalues of complex Hamiltonians with  $PT$ -symmetry. II. Phys. Lett. A 250 (1998), no. 1-3, 29–32.

E. Delabaere, D. T. Trinh, Spectral analysis of the complex cubic oscillator. J. Phys. A 33 (2000), no. 48, 8771–8796.

P. Dorey, C. Dunning and R. Tateo, Spectral equivalences, Bethe ansatz equations, and reality properties in  $PT$ -symmetric quantum mechanics, J. Phys. A, 34 (2001) 5679–5704.

A. Eremenko, A. Gabrielov, Analytic continuation of eigenvalues of a quartic oscillator, Comm. Math. Phys., v. 287, No. 2 (2009) 431-457.

A. Eremenko and A. Gabrielov, Irreducibility of some spectral determinants, arXiv:0904.1714

V. Grecchi, M. Maioli and A. Matrinez, Padeé summability of the cubic oscillator, J. Math. Phys. A: Math. Theor. 42 (2009) 425208.

R. Nevanlinna, Über Riemannsche Flächen mit endlich vielen Windungspunkten, Acta Math., 58 (1932) 295–373.

K. Shin, On the reality of the eigenvalues for a class of PT-symmetric oscillators, Comm. Math. Phys., 229 (2002) 543–564.

Y. Sibuya, Global theory of a second order linear ordinary differential equation with a polynomial coefficient, North-Holland, Amsterdam–Oxford; American Elsevier, NY, 1975.

B. Simon, Coupling constant analyticity for the anharmonic oscillator, Ann. Physics 58 (1970), 76–136.

Duc Tai Trinh, Asymptotique et analyse spectrale de l'oscillateur cubique, These, 2002.