


Lesson 18: Concavity, Inflection Points & the Second Derivative Test


Lesson 19: Concavity, Inflection Pts & the Second Derivative Test

Concave Up



Like a cup

Concave Down



Like a frown

Concavity of a Function:

Suppose $f''(x)$ exists on an open interval, I .

Then

- Ⓐ If $f''(x) > 0$ for all x in I , then $f(x)$ is **concave up** on I .
- Ⓑ If $f''(x) < 0$ for all x in I , then $f(x)$ is **concave down** on I .

Ex 1: Determine the largest open interval(s) on which $f(x) = x^3 - x$ is concave up or down.

Step 1: Find when $f''(x) = 0$


$$f'(x) = 3x^2 - 1$$

$$f''(x) = 6x = 0$$


$$x = 0$$

Step 2+3: Draw a # line with the pts from ①

Determine test pts using ②



Step 4: Plug those values into f'' to determine whether it's positive or negative




$$f''(x) = 6x$$

$$f''(-1) = 6(-1)$$

$$f''(1) = 6(1)$$

Step 5: Use definition of concavity



Concave Up: $(0, \infty)$

Concave Down: $(-\infty, 0)$

Ex 2: Let $f(x) = \frac{1}{12}x^4 - \frac{1}{3}x^3$.

Ⓐ Determine the largest open interval(s) on which $f(x)$ is increasing or decreasing.

Step 1: Find when $f'(x) = 0$

$$f'(x) = \frac{4}{12}x^3 - \frac{3}{3}x^2 = \frac{1}{3}x^3 - x^2 = 0$$

$$x^2 \left(\frac{1}{3}x - 1 \right) = 0$$


$$x^2 = 0 \quad \left| \quad \frac{1}{3}x - 1 = 0$$

$$x = 0 \quad \left| \quad \frac{1}{3}x = 1$$


$$x = 3$$

Step 2+3: Draw a # line with the pts from ①


Determine test pts using ②



Step 4: Plug those values into f' to determine whether it's positive or negative



Step 5: Use definition of increasing/decreasing



Ex 2: Let $f(x) = \frac{1}{12}x^4 - \frac{1}{3}x^3$.

⑤ Determine the largest open interval(s) on which $f(x)$ is concave up or down.

Step 1: Find when $f''(x) = 0$

$$f'(x) = \frac{4}{12}x^3 - \frac{3}{3}x^2 = \frac{1}{3}x^3 - x^2$$

$$f''(x) = \frac{3}{3}x^2 - 2x = x^2 - 2x = 0$$

$$x(x-2) = 0$$


$$x=0 \quad | \quad x-2=0$$

$$x=2$$


Step 2: Draw a # w/ the pts from ①



Step 3: Determine test pts using ②.



Step 4: Plug those values into f'' to determine whether it's positive or negative



$$f''(x) = x(x-2)$$

$$f''(-1) = -1(-1-2)$$

$$- \cdot - = +$$


$$f''(1) = 1(1-2)$$

$$+ \cdot - = -$$

$$f''(3) = 3(3-2)$$

$$+ \cdot + = +$$

Step 5: Use definition of concavity





Concave Up: $(-\infty, 0) \cup (2, \infty)$
 Concave Down: $(0, 2)$

Ex 2: Let $f(x) = \frac{1}{12}x^4 - \frac{1}{3}x^3$.

Ⓒ Determine the largest open interval(s) on which $f(x)$ is decreasing and concave up.

Use your # lines from Ⓐ and Ⓑ



| | $(-\infty, 0)$ | $(0, 2)$ | $(2, 3)$ | $(3, \infty)$ |
|-------|----------------|----------|----------|---------------|
| f' | - | - | - | + |
| f'' | + | - | + | + |

Answer: $(-\infty, 0) \cup (2, 3)$

Steps in Finding the Inflection Pts

① Find the pts on the curve where the second derivative is 0 or DNE.

i.e. Find where $f''(x) = 0$ and $f''(x)$ DNE

② Test whether the concavity changes @ these pts

i.e. $\leftarrow \begin{matrix} - \\ | \\ c \end{matrix} \begin{matrix} + \\ | \\ c \end{matrix} \rightarrow f''$ or $\leftarrow \begin{matrix} + \\ | \\ c \end{matrix} \begin{matrix} - \\ | \\ c \end{matrix} \rightarrow f''$

where c is a pt found in ①

Ex 3: Find the inflection pt(s) of $f(x) = \frac{3}{5}x^5 - x^4$ if they exist.

Step 1: Find when $f''(x) = 0$

$$f'(x) = \frac{3}{5}(5)x^4 - 4x^3 = 3x^4 - 4x^3$$

$$f''(x) = 12x^3 - 12x^2 = 0$$

$$12x^2(x-1) = 0$$


$$\begin{array}{l|l} 12x^2 = 0 & x-1 = 0 \\ x = 0 & x = 1 \end{array}$$

Step 2+3: Draw a # line with the pts from ①

Determine test pts using ②




Step 4 Plug those values into f' to determine whether it's positive or negative




Step 5: Use definition of inflection pts.

i.e. Look for sign change.



Does the sign change?

No $\Rightarrow x=0$ NOT
Inflection Pt



Does the sign change?

Yes $\Rightarrow x=1$ IS
Inflection Pt

Second Derivative Test

Let $f(x)$ be a function such that $f'(c)=0$ and $f''(x)$ exists on an open interval containing c .

- ① If $f''(c) > 0$ then $f(x)$ has a relative min $f(c)$ at $x=c$.
- ② If $f''(c) < 0$ then $f(x)$ has a relative max $f(c)$ at $x=c$.

If $f''(c)=0$, the Second Derivative Test does not apply or fail.

It does not necessarily mean that we have neither a relative max or min at these pts.

It just means you need to apply First Derivative Test.

Ex 4: Find all the relative extrema of $f(x) = x^3 - x$ if they exist.

Step 1: Find $f'(x) = 0$

$$f'(x) = 3x^2 - 1 = 0$$

$$x^2 = 1/3$$

$$x = \pm \sqrt{1/3}$$

Step 3: Plug ① into ②

$$f''(\sqrt{1/3}) = 6\sqrt{1/3}$$

$$f''(-\sqrt{1/3}) = 6(-\sqrt{1/3})$$

Step 2: Find $f''(x)$

$$f''(x) = 6x$$

Step 4: Use Second Derivative Test

$$f''(\sqrt{1/3}) = 6\sqrt{1/3} > 0 \quad \cup$$

$$\Rightarrow \text{rel min}$$

$$f''(-\sqrt{1/3}) = 6(-\sqrt{1/3}) < 0 \quad \cap$$

$$\Rightarrow \text{rel max}$$

Ex 5: Find all the relative extrema of $f(x) = \frac{3}{5}x^5 - x^4$ if they exist.

Step 1: Find $f'(x) = 0$

$$f'(x) = \frac{3}{5}(5)x^4 - 4x^3$$

$$3x^4 - 4x^3 = 0$$

$$x^3(3x - 4) = 0$$

$$x^3 = 0 \quad | \quad 3x - 4 = 0$$

$$x = 0 \quad | \quad x = \frac{4}{3}$$

Step 2: Find $f''(x)$

$$f''(x) = 12x^3 - 12x^2$$

$$= 12x^2(x - 1)$$

Step 3: Plug ① into ②

$$f''(0) = 12(0)^2(0 - 1) = 0$$

$$f''\left(\frac{4}{3}\right) = 12\left(\frac{4}{3}\right)^2\left(\frac{4}{3} - 1\right)$$

Step 4: Use Second Derivative Test


$$f''(0) = 12(0)^2(0 - 1) = 0 \implies \text{Fails so we need to use First Derivative Test}$$

$$f''\left(\frac{4}{3}\right) = 12\left(\frac{4}{3}\right)^2\left(\frac{4}{3} - 1\right) > 0 \quad \cup$$

$$\implies \text{rel min @ } x = \frac{4}{3}$$

First Derivative Test for $x = 0$.

$$f'(x) = x^3(3x - 4)$$



$\cap \implies \text{relative max @ } x = 0$