

Please show **all** your work! Answers without supporting work will not be given credit.
Write answers in spaces provided.

Name: _____

1. You are supposed to know and understand the basics about the exponential function, and about the logarithmic function as the inverse of the exponential function. You should be able to solve the equations involving them.

Example Problems

- (a) Solve the following equations.

i. $\ln(x) + \ln(x - 3) = 0$

x = _____

ii. $7^{2x+1} = 21$

x = _____

iii. $3^{\ln(x)} = 5$

x = _____

iv. $e^{2x^2-7x-15} = 1$

x = _____

(b) Find the exact values.

i. $\frac{\log_{15}(9) + \log_{15}(25)}{e^{3\ln(5)+2\ln(7)}}$

$\frac{\log_{15}(9) + \log_{15}(25)}{e^{3\ln(5)+2\ln(7)}}$ = _____

ii. $\log_b \frac{\sqrt{x}}{\sqrt[3]{z}}$ when $\log_b(x) = 2.4$ and $\log_b(z) = 3.6$

$\log_b \frac{\sqrt{x}}{\sqrt[3]{z}}$ = _____

-
2. You are supposed to know the domain and range of each of the inverse trigonometric functions \sin^{-1} , \cos^{-1} , and \tan^{-1} . You are also supposed to know the basic identities involving the inverse trigonometric functions.

Example Problems

- (a) Find the exact values for the following:

i. $\sin^{-1}\left(-\frac{\sqrt{3}}{2}\right)$

$$\sin^{-1}\left(-\frac{\sqrt{3}}{2}\right) = \underline{\hspace{2cm}}$$

ii. $\cos^{-1}\left(-\frac{\sqrt{2}}{2}\right)$

$$\cos^{-1}\left(-\frac{\sqrt{2}}{2}\right) = \underline{\hspace{2cm}}$$

iii. $\tan^{-1}\left(\frac{\sqrt{3}}{3}\right)$

$$\tan^{-1}\left(\frac{\sqrt{3}}{3}\right) = \underline{\hspace{2cm}}$$

iv. $\sin^{-1}\left(\sin\left(\frac{8\pi}{7}\right)\right)$

$$\sin^{-1}\left(\sin\left(\frac{8\pi}{7}\right)\right) = \underline{\hspace{2cm}}$$

v. $\sin^{-1}\left(\cos\left(\frac{3\pi}{5}\right)\right)$

$$\sin^{-1}\left(\cos\left(\frac{3\pi}{5}\right)\right) = \underline{\hspace{2cm}}$$

vi. $\sin^{-1}\left(\cos\left(\frac{4\pi}{7}\right)\right)$

$$\sin^{-1}\left(\cos\left(\frac{4\pi}{7}\right)\right) = \underline{\hspace{2cm}}$$

vii. $\cos\left(\sin^{-1}\left(\frac{1}{5}\right)\right)$

$$\cos\left(\sin^{-1}\left(\frac{1}{5}\right)\right) = \underline{\hspace{2cm}}$$

viii. $\sin\left(\cos^{-1}\left(-\frac{1}{5}\right)\right)$

$$\sin\left(\cos^{-1}\left(-\frac{1}{5}\right)\right) = \underline{\hspace{2cm}}$$

ix. $\tan\left(\sin^{-1}\left(\frac{1}{2}\right)\right)$

$$\tan\left(\sin^{-1}\left(\frac{1}{2}\right)\right) = \underline{\hspace{2cm}}$$

x. $\sin^{-1}\left(\frac{1}{3}\right) + \cos^{-1}\left(\frac{1}{3}\right)$

$$\sin^{-1}\left(\frac{1}{3}\right) + \cos^{-1}\left(\frac{1}{3}\right) = \underline{\hspace{2cm}}$$

xi. $\tan^{-1}\left(\tan\left(\frac{4\pi}{5}\right)\right) + \tan^{-1}\left(\tan\left(\frac{6\pi}{5}\right)\right)$

$$\tan^{-1}\left(\tan\left(\frac{4\pi}{5}\right)\right) + \tan^{-1}\left(\tan\left(\frac{6\pi}{5}\right)\right) = \underline{\hspace{2cm}}$$

-
3. You are supposed to be able to solve the equations involving the trigonometric functions and find solutions on the given interval, using the basic formulas of the trigonometric functions (e.g., double angle formula for sine and cosine, $\sin^2(x) + \cos^2(x) = 1$, etc.).

Example Problems

- (a) Find the values of x on the interval $[0, 2\pi]$ which satisfy the equation.

i. $\cos^2(x) = \cos(2x)$

x = _____

ii. $2\sin^2(x) = \cos(2x)$

x = _____

- (b) Find the values of x on the interval $[0, 2\pi)$ which satisfy the equation

i. $\sin(x) = \sin(2x)$

x = _____

ii. $\sin(x) = \cos(2x)$

x = _____

iii. $-\sin^2(x) = \cos(2x)$

x = _____

(c) Find the values of x on the interval $[0, 2\pi]$ which satisfy the equation

$$\tan^2(x) - 3 = 0$$

x = _____

-
4. You are supposed to express the trigonometric functions, using a right-triangle relationship, as an algebraic function of x , when the angle is given in terms of x via the inverse trigonometric functions.

Example Problems

- (a) Express the following as an algebraic function of x .

i. $\cos\left(\sin^{-1}\left(\frac{x}{3}\right)\right)$

$$\cos\left(\sin^{-1}\left(\frac{x}{3}\right)\right) = \underline{\hspace{2cm}}$$

ii. $\sin\left(\cos^{-1}\left(\frac{x}{5}\right)\right)$

$$\sin\left(\cos^{-1}\left(\frac{x}{5}\right)\right) = \underline{\hspace{2cm}}$$

iii. $\cos\left(\sin^{-1}(7x)\right)$

$$\cos\left(\sin^{-1}(7x)\right) = \underline{\hspace{2cm}}$$

iv. $\tan(\sin^{-1}(7x))$

$$\tan(\sin^{-1}(7x)) = \underline{\hspace{2cm}}$$

v. $\sin(2\cos^{-1}(3x))$

$$\sin(2\cos^{-1}(3x)) = \underline{\hspace{2cm}}$$

vi. $\sin(\tan^{-1}(2x))$

$$\sin(\tan^{-1}(2x)) = \underline{\hspace{2cm}}$$

vii. $\cot\left(\sin^{-1}\left(\frac{x}{5}\right)\right)$

$$\sin(\tan^{-1}(2x)) = \underline{\hspace{2cm}}$$

5. You should know the condition (one-to-one) for a function to have its inverse. Given a function which is one-to-one, you are supposed to be able to find the formula of the inverse function, its domain and range, and draw the graph of the inverse function.

Example Problems

- (a) Find the formula, and state the domain and range of the inverse of the function

$$y = f(x) = \ln\left(\frac{1}{3}e^{2x} - 3\right)$$

$$f^{-1}(x) = \underline{\hspace{2cm}}$$

$$\text{Domain of } f^{-1}(x): \underline{\hspace{2cm}}$$

$$\text{Range of } f^{-1}(x): \underline{\hspace{2cm}}$$

-
- (b) Consider the function $y = f(x) = \frac{x+5}{x-2}$ over the domain $(2, \infty)$. Find the formula and domain of its inverse function.

$$f^{-1}(x) = \underline{\hspace{4cm}}$$

$$\text{Domain of } f^{-1}(x): \underline{\hspace{4cm}}$$

- (c) Consider the function $y = f(x) = \frac{2x-1}{x+5}$ over the domain $(0, \infty)$. Find the formula and domain of its inverse function.

$$f^{-1}(x) = \underline{\hspace{4cm}}$$

$$\text{Domain of } f^{-1}(x): \underline{\hspace{4cm}}$$

-
- (d) Consider the function $y = f(x) = \frac{2x+1}{x+1}$ over the domain $(-1, 0)$. Find the formula and domain of its inverse function.

$$f^{-1}(x) = \underline{\hspace{4cm}}$$

$$\text{Domain of } f^{-1}(x): \underline{\hspace{4cm}}$$

6. You are supposed to be able to compute the (right/left hand side) limit, understanding its proper meaning, and using the Squeeze Theorem. You are also supposed to be able to determine the exact value of the limit who has an indeterminate form (e.g., $0/0$, $\pm\infty//\pm\infty$, $\infty - \infty$) using some proper technique.

Example Problems

- (a) Compute the following limits:

i. $\lim_{x \rightarrow 5^-} \frac{x^2 - 3x - 10}{|x - 5|}$

$$\lim_{x \rightarrow 5^-} \frac{x^2 - 3x - 10}{|x - 5|} = \underline{\hspace{4cm}}$$

ii. $\lim_{x \rightarrow (\pi/2)^-} \tan(x)$

$$\lim_{x \rightarrow (\pi/2)^-} \tan(x) = \underline{\hspace{2cm}}$$

iii. $\lim_{x \rightarrow (\pi/2)^+} e^{\tan(x)}$

$$\lim_{x \rightarrow (\pi/2)^+} e^{\tan(x)} = \underline{\hspace{2cm}}$$

iv. $\lim_{x \rightarrow 2} \left(\frac{x^2 + 5x - 14}{x^2 - 6x + 8} \right)$

$$\lim_{x \rightarrow 2} \left(\frac{x^2 + 5x - 14}{x^2 - 6x + 8} \right) = \underline{\hspace{2cm}}$$

v. $\lim_{x \rightarrow 0} \left(\frac{5}{x^2 - x} + \frac{5}{x} \right)$

$$\lim_{x \rightarrow 0} \left(\frac{5}{x^2 - x} + \frac{5}{x} \right) = \underline{\hspace{2cm}}$$

vi. $\lim_{x \rightarrow \infty} (\sqrt{9x^2 - x} - 3x)$

$$\lim_{x \rightarrow \infty} (\sqrt{9x^2 - x} - 3x) = \underline{\hspace{2cm}}$$

vii. $\lim_{x \rightarrow -\infty} (\sqrt{x^2 + 3x} + x)$

$$\lim_{x \rightarrow -\infty} (\sqrt{x^2 + 3x} + x) = \underline{\hspace{2cm}}$$

viii. $\lim_{x \rightarrow 0} \frac{|3x - 4| - |5x + 4|}{x}$

$$\lim_{x \rightarrow 0} \frac{|3x - 4| - |5x + 4|}{x} = \underline{\hspace{2cm}}$$

ix. $\lim_{x \rightarrow 0} \frac{\sin(3x)}{7x}$

$$\lim_{x \rightarrow 0} \frac{\sin(3x)}{7x} = \underline{\hspace{2cm}}$$

x. $\lim_{x \rightarrow \infty} \frac{\sin(x)}{\ln(x)}$

$$\lim_{x \rightarrow \infty} \frac{\sin(x)}{\ln(x)} = \underline{\hspace{2cm}}$$

xi. $\lim_{x \rightarrow \infty} \frac{\cos(3x)}{\ln(x)}$

$$\lim_{x \rightarrow \infty} \frac{\cos(3x)}{\ln(x)} = \underline{\hspace{2cm}}$$

xii. $\lim_{x \rightarrow 0} \frac{\sqrt{x+1} - 1}{\tan(x/2)}$

$$\lim_{x \rightarrow 0} \frac{\sqrt{x+1} - 1}{\tan(x/2)} = \underline{\hspace{2cm}}$$

xiii. $\lim_{x \rightarrow 0} \frac{\sin(1/x)}{1/x}$

$$\lim_{x \rightarrow 0} \frac{\sin(1/x)}{1/x} = \underline{\hspace{2cm}}$$

$$\text{xiv. } \lim_{x \rightarrow 8} \frac{\sqrt[3]{x} - 2}{x - 8}$$

$$\lim_{x \rightarrow 8} \frac{\sqrt[3]{x} - 2}{x - 8} = \underline{\hspace{2cm}}$$

$$\text{xv. } \lim_{x \rightarrow \infty} \frac{x^3 + 3x^2}{2x^3 + \sqrt{9x^6 + 4x^2}}$$

$$\lim_{x \rightarrow \infty} \frac{x^3 + 3x^2}{2x^3 + \sqrt{9x^6 + 4x^2}} = \underline{\hspace{2cm}}$$

$$\text{xvi. } \lim_{x \rightarrow -\infty} \frac{x^3 + 3x^2}{2x^3 + \sqrt{9x^6 + 4x^4}}$$

$$\lim_{x \rightarrow -\infty} \frac{x^3 + 3x^2}{2x^3 + \sqrt{9x^6 + 4x^4}} = \underline{\hspace{2cm}}$$

$$\text{xvii. } \lim_{x \rightarrow 0^-} \frac{x^3 + 3x^2}{2x^3 + \sqrt{9x^6 + 4x^4}}$$

$$\lim_{x \rightarrow 0^-} \frac{x^3 + 3x^2}{2x^3 + \sqrt{9x^6 + 4x^4}} = \underline{\hspace{2cm}}$$

$$\text{xviii. } \lim_{x \rightarrow 0^-} \frac{2x + 5x^3}{3x + \sqrt{25x^6 + x^2}}$$

$$\lim_{x \rightarrow 0^-} \frac{2x + 5x^3}{3x + \sqrt{25x^6 + x^2}} = \underline{\hspace{2cm}}$$

$$\text{xix. } \lim_{x \rightarrow 0^+} \frac{2x + 5x^3}{3x + \sqrt{25x^6 + x^2}}$$

$$\lim_{x \rightarrow 0^+} \frac{2x + 5x^3}{3x + \sqrt{25x^6 + x^2}} = \underline{\hspace{2cm}}$$

$$\lim_{x \rightarrow \infty} (\sqrt{x+1} - \sqrt{x}) \cdot \sqrt{x} = \underline{\hspace{2cm}}$$

7. You are supposed to understand the meaning of the Intermediate Value Theorem (IVT), and to be able to use it to show that a certain equation has a root in a specified interval.

Example Problems

- (a) Using the IVT determine on which of the following intervals the equation $x^3 - 3x = 5$ has a root?
- (A) (0,1)
 - (B) (1,2)
 - (C) (2,3)
 - (D) (3,4)
 - (E) (4,5)

Answer:_____

(b) From the IVT, which of the following interval, can one conclude, contains a solution of the equation $2x^3 + 3x^2 = 5x - 3$?

- (A) $(-3,-2)$
- (B) $(-2,-1)$
- (C) $(-1,0)$
- (D) $(0,1)$
- (E) $(1,2)$

Answer:_____

(c) From the IVT, which of the following interval, can one conclude, contains a solution of the equation $x^3 - 3x^2 = -2x + 1$?

- (A) $(-1,0)$
- (B) $(0,1)$
- (C) $(1,2)$
- (D) $(2,3)$
- (E) $(3,4)$

Answer:_____

-
8. You are supposed to understand the meaning of the defining formula of the derivative, and being able to determine the values of the related limits.

Example Problems

(a) Suppose we have a function $f(x)$ with $f'(2) = 5$. Determine the following values:

i. $g'(1)$ where $g(x) = f(2x)$

$$g'(1) = \underline{\hspace{2cm}}$$

ii. $\lim_{h \rightarrow 0} \frac{f(2+4h) - f(2)}{3h}$

$$\lim_{h \rightarrow 0} \frac{f(2+4h) - f(2)}{3h} = \underline{\hspace{2cm}}$$

iii. $\lim_{h \rightarrow 0} \frac{f(2+4h) - f(2-5h)}{7h}$

$$\lim_{h \rightarrow 0} \frac{f(2+4h) - f(2-5h)}{7h} = \underline{\hspace{2cm}}$$

$$f'(5) = \lim_{h \rightarrow 0} \frac{f(5+h) - f(5)}{h} = 3$$

then evaluate the following limit

$$\lim_{h \rightarrow 0} \frac{f(5+2h) - f(5-4h)}{h}$$

$$\lim_{h \rightarrow 0} \frac{f(5+2h) - f(5-4h)}{h} = \underline{\hspace{2cm}}$$

(c) Let $f(x)$ be the function defined as follows:

$$f(x) = \begin{cases} -(x-1)^2 + 5 & \text{if } x < 0 \\ 2x + 4 & \text{if } x \geq 0 \end{cases}$$

Does $f'(0)$ exists? If it exists, compute the value of $f'(0)$.

Answer:_____

(d) We have a function $f(x)$ such that

$$\begin{cases} f(4) = 3 \\ f'(4) = 2 \end{cases}$$

Compute the following limit:

$$\lim_{h \rightarrow 0} \frac{f(4+h)\sqrt{4+h} - 6}{h}$$

HINT: What is the relation between the derivative of the product function $g(x) = f(x)\sqrt{x}$ at $x = 4$ and the given limit above?

$$\lim_{h \rightarrow 0} \frac{f(4+h)\sqrt{4+h} - 6}{h} = \underline{\hspace{2cm}}$$

(e) Compute the following limit

$$\lim_{h \rightarrow 0} \frac{\ln(1+3h)}{7h}$$

$$\lim_{h \rightarrow 0} \frac{\ln(1+3h)}{7h} = \underline{\hspace{2cm}}$$

-
9. When a function is defined piecewise and depending on some variables, you are supposed to know how to determine those variables so that the function becomes continuous entirely over the specified interval (e.g., everywhere).

Example Problems

- (a) Find the values of a and b so that the function

$$f(x) = \begin{cases} x^2 - a & \text{if } x \leq 1 \\ \frac{3x^2 + 12x - b}{x^2 + 2x - 3} & \text{if } x > 1 \end{cases}$$

is continuous on $(-\infty, \infty)$.

a = _____

b = _____

- (b) Find the values of a and b so that the function

$$f(x) = \begin{cases} x - b + 7 & \text{if } x \leq 3 \\ \frac{ax + b}{x - 3} & \text{if } x > 3 \end{cases}$$

is continuous on $(-\infty, \infty)$.

a = _____

b = _____

-
10. You are supposed to be able to find the horizontal/vertical asymptote(s) of a given function.

Example Problems

- (a) Find the horizontal/vertical asymptote(s) of the following functions

i. $y = f(x) = \frac{x^3 + 4x^2 + x - 6}{x(x^2 - 1)}$

HA: _____

VA: _____

ii. $y = f(x) = \frac{7 + e^x}{4 - e^x}$

HA: _____

VA: _____

iii. $y = f(x) = \frac{x^3 + x^2}{3x^2 + \sqrt{25x^6 + x^2}}$

HA: _____

VA: _____

iv. $y = f(x) = \frac{4e^{2x} - 16}{e^{2x} - 3e^x - 10}$

HA: _____

VA: _____

v. $y = f(x) = \frac{x-7}{\sqrt{2x^2+7x}}$

HA: _____

VA: _____

vi. $y = f(x) = \frac{e^{2x}-1}{(e^x-1)(e^x-2)}$

HA: _____

VA: _____

-
11. You are supposed to be able to compute the derivative of a polynomial (square root/rational) function, and understand that its value represents the slope of the tangent line to the graph of the function.

Example Problems

- (a) Find the equation of the line that is tangent to the curve $y = \sin(x)$ over $(0, \pi)$ and that is parallel to the line $y = -\frac{1}{2}x + 5$.

Answer:_____

- (b) Find an equation for the line tangent to the graph of $f(x) = \sin(x)$ over the interval $(0, \pi)$, which is perpendicular to the line $y = 2x + 1$.

Answer:_____

-
- (c) Find an equation for the line tangent to the graph of $f(x) = \tan(x)$ over the interval $(-\pi/2, 0)$ which is parallel to the line $y = 2x + 1$.

Answer:_____

- (d) Find the equation of the line(s) which is tangent to the parabola $y = x^2$ and passes through the point $(1, -3)$.

Answer:_____

-
12. You are supposed to understand the meaning of the continuity and differentiability, and their difference.

Example Problems

- (a) Determine when the following function is continuous/differentiable.

i. $f(x) = \begin{cases} \frac{x-2}{|x-2|} & \text{if } x \neq 2 \\ 0 & \text{if } x = 2 \end{cases}$

Continuous? _____

Differentiable? _____

ii. $g(x) = x|x-2|$

Continuous? _____

Differentiable? _____

iii. $h(x) = x(x-2)|x-2|$

Continuous? _____

Differentiable? _____

(b) Consider the following piece-wise defined function:

$$h(x) = \begin{cases} -4x & \text{if } x \leq 0 \\ x(x-2)|x-2| & \text{if } x > 0 \end{cases}$$

Judge continuity/ differentiability of the function (i) at $x = 0$ and (ii) at $x = 2$.

Continuous at $x = 0$?_____

Differentiable at $x = 0$?_____

Continuous at $x = 2$?_____

Differentiable at $x = 2$?_____

What happens if we change the first part of the description of $h(x)$ to " $3x$ if $x \leq 0$ "?

Answer:_____

(c) Judge continuity/differentiability of the following function at $x = 0$.

i. $f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0, \\ 0 & \text{if } x = 0 \end{cases}$

Continuous at $x = 0$? _____

Differentiable at $x = 0$? _____

ii. $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0, \\ 0 & \text{if } x = 0 \end{cases}$

Continuous at $x = 0$? _____

Differentiable at $x = 0$? _____

-
13. You are supposed to know how to compute the derivative of the sum, product, and quotient of two functions. You are suppose to know the following relation between the derivative of the original function and its inverse.

Fact: Suppose that we have the function f and its inverse f^{-1} and that $f(a) = b$ (i.e., $a = f^{-1}(b)$.) Then

$$\{f^{-1}\}'(b) = \frac{1}{f'(a)}$$

Example Problems

- (a) Suppose that f, g and h are functions that satisfy the following conditions:

$$\begin{cases} f(0) = 2, f'(0) = 3, f'(-3) = 2, f(-3) = 0, \\ g(0) = -1, g'(0) = 5, \\ h(0) = 1, h'(0) = -2 \end{cases}$$

Evaluate the derivative of the following function at $x = 0$.

$$F(x) = \frac{f^{-1}(x)}{g(x) - h(x)}$$

$$F'(0) = \underline{\hspace{2cm}}$$

-
- i. Suppose that f is a function that satisfy the following conditions:

$$f(2) = 3 \text{ and } f'(2) = 5$$

Evaluate the derivative of the following function at $x = 3$

$$F(x) = \frac{1}{f^{-1}(x)}$$

$$F'(3) = \underline{\hspace{2cm}}$$

- ii. Suppose that f is a function that satisfy the following conditions:

$$f(5) = 2 \text{ and } f'(5) = 3$$

Evaluate the derivative of the following function at $x = 2$

$$F(x) = \frac{1}{f^{-1}(x)}$$

$$F'(2) = \underline{\hspace{2cm}}$$