

Reminders

- WEDNESDAY QUIZ 5 on
 - Volume of Revolutions
 - Disks (Lesson 14)
 - Washers (Lesson 15)
- NEXT FRIDAY NO CLASS

1

MA 16020: Lesson 16

Volume By Revolution

Rotation around any non-Axis

By Alexandra Cuadra

2

RECAP of Formulas from Lesson 14 and 15

For rotation around x-axis:

- Disk Method:

$$V = \pi \int_a^b [f(x)]^2 dx$$

- Washer Method:

$$V = \pi \int_a^b (R^2 - r^2) dx$$

For rotation around y-axis:

- Disk Method:

$$V = \pi \int_c^d [g(y)]^2 dy$$

- Washer Method:

$$V = \pi \int_c^d (R^2 - r^2) dy$$

3

RECAP: When do we apply Disk Method or Washer Method?

- When the region “hugs” the axis of rotation

⇒ Disk Method

- When there is a “gap” between the region and axis of rotation

⇒ Washer Method

4

Today's Lecture


- In the previous two lessons, we looked at rotations around the x-axis or y-axis.
- Today we are going to rotate about **ANY** arbitrary axis.
 - Don't worry. We are going to limit ourselves to any vertical or horizontal line parallel to the x-axis or y-axis

5

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x-axis

Draw the region.




→ WASHER

<https://www.geogebra.org/m/wrj2euhf>


6

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



Rotation about x-axis




<https://www.geogebra.org/m/wrj2euhf>


7

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



Furthermore, 3-D




<https://www.geogebra.org/m/wrj2euhf>

8

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x-axis



$\Rightarrow y = ?$ and Top, Bottom
 Top $\Rightarrow x^2y = 4 \Rightarrow y = \frac{4}{x^2}$
 Bottom $\Rightarrow y = 1$


$$V = \pi \int \left(\frac{4}{x^2} \right)^2 - 1^2 dx$$

<https://www.geogebra.org/m/wrj2euhf>

9

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x-axis Now the bounds we are




given the smallest on $x=1$ next find the other by putting $y=1$ into $x^2y=4$

<https://www.geogebra.org/m/wrj2euhf>

10

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



$$\text{So } x^2 = 4, x = \pm 2$$

But we are looking for $x > 1$, so $x = 2$ Hence


$$V = \pi \int_1^2 \left(\frac{4}{x^2} \right)^2 - 1^2 dx$$

<https://www.geogebra.org/m/wrj2euhf>

11

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



$$V = \pi \int_1^2 \frac{16}{x^4} - 1 dx$$

$$= \pi \int_1^2 16x^{-4} - 1 dx$$

$$= \pi \left(\frac{16x^{-3}}{-3} - x \right) \Big|_1^2 = \frac{11\pi}{3}$$


<https://www.geogebra.org/m/wrj2euhf>

12

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the y -axis

Draw the region.




WASHER

<https://www.geogebra.org/m/wzbm2xht>


13

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the y -axis



Rotation about y -axis




<https://www.geogebra.org/m/wzbm2xht>


14

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the y -axis



Furthermore, 3-D




<https://www.geogebra.org/m/wzbm2xbt>

15

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the y -axis



$\Rightarrow x = ?$ and Right, Left
 Right $\Rightarrow x^2y = 4 \Rightarrow x = \sqrt{\frac{4}{y}}$
 Left $\Rightarrow x = 1$


$$V = \pi \int \left(\sqrt{\frac{4}{y}} \right)^2 - 1^2 dy$$

<https://www.geogebra.org/m/wzbm2xbt>

16

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the y -axis




Now the bounds we are given
the smallest one $y=1$
Next find the other by
putting
 $x=1$ into $x^2y=4$
So $y=4$

<https://www.geogebra.org/m/wzbm2xbt>

17

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the y -axis



$$\begin{aligned}
 V &= \pi \int_1^4 \left(\left(\sqrt{\frac{4}{y}} \right)^2 - 1^2 \right) dy \\
 &= \pi \int_1^4 \left(\frac{4}{y} - 1 \right) dy \\
 &= \pi \left(4 \ln |y| - y \right) \Big|_1^4 \\
 &\approx 7.9959
 \end{aligned}$$

<https://www.geogebra.org/m/wzbm2xbt>

18


Let's Backtrack a Bit...

Remember when we first described Washers, we talked about farthest and closest.

Consider the case of x-axis rotation.

In terms of distance,


- R is the length of Top Function away from x-axis
 - i.e. $R = f$
- r is the length of Bottom Function away from x-axis
 - i.e. $r = g$



19

When rotating around the line $y = e$...


- So, what is the distance between f (or g) and $y = e$?
 - Distance b/w f and $y = e$ is $R = f - e$
 - Distance b/w g and $y = e$ is $r = g - e$
- Note this formula is also true for the x-axis case, because the x-axis is simply the line $y = 0$



20

GOOD NEWS EVERYBODY: When rotating around the line $x = e$...

- The same formulas, for R and r , from the case of $y = e$ applies.
- So, the distance between f (or g) and $x = e$ are as follows:
 - $R = f - e$
 - $r = g - e$
- Note that though we did all this calculation for the Washer Problems; this also applies for the Disk Problems.



21

Rotation around any non-Axis Formulas

For rotation around the line $y = e$:

- Disk Method:

$$V = \pi \int_a^b [f(x) - e]^2 dx$$

- Washer Method:

$$V = \pi \int_a^b ((R - e)^2 - (r - e)^2) dx$$

For rotation around the line $x = e$:

- Disk Method:

$$V = \pi \int_c^d [g(y) - e]^2 dy$$

- Washer Method:


$$V = \pi \int_c^d ((R - e)^2 - (r - e)^2) dy$$

Note: That these formulas work for the case of x-axis ($y = 0$) and y-axis ($x = 0$).

22

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the line $y = 1$
Draw the region.




<https://www.geogebra.org/m/n2jzwh8f>


23

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the line $y = 1$



Rotation about $y = 1$




<https://www.geogebra.org/m/n2jzwh8f>

24


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the line $y = 1$



DISK PROBLEM

Rotation about $y = 1$




<https://www.geogebra.org/m/n2jzwh8f>

25


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the line $y = 1$



DISK PROBLEM

Furthermore, 3-D




<https://www.geogebra.org/m/n2jzwh8f>

26

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the line $y = 1$



dx problem

But we saw this is now a disk problem so

$$V = \pi \int_1^2 \left(\frac{4}{x^2} - 1 \right)^2 dx$$


Note our bounds are the same as (a)

<https://www.geogebra.org/m/n2jzwh8f>

27

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the line $y = 1$



$$V = \pi \int_1^2 \left(\frac{16}{x^4} - \frac{8}{x^2} + 1 \right) dx$$

$$= \pi \int_1^2 (16x^{-4} - 8x^{-2} + 1) dx$$

$$= \pi \left(\frac{16x^{-3}}{-3} - \frac{8x^{-1}}{-1} + x \right) \Big|_1^2$$


$$= 5\pi/3$$

<https://www.geogebra.org/m/n2jzwh8f>

28

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$
Draw the region.




<https://www.geogebra.org/m/cppyhngk>


29

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



Rotation about $x = 1$




<https://www.geogebra.org/m/cppyhngk>

30


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



DISK PROBLEM

Furthermore, 3-D




<https://www.geogebra.org/m/cppyhngk>

31

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



dy problem
 But we saw this is now
 a disk problem so


$$V = \pi \int_1^4 \left(\sqrt{\frac{4}{y}} - 1 \right)^2 dy$$
 Note our bounds are the
 same as (b)

<https://www.geogebra.org/m/cppyhngk>

32

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$




$$\begin{aligned}
 V &= \pi \int_1^4 \left(\frac{4}{y} - 2\sqrt{\frac{4}{y}} + 1 \right) dy \\
 &= \pi \int_1^4 \left(\frac{4}{y} - \frac{2 \cdot 2}{y^{1/2}} + 1 \right) dy \\
 &= \pi \int_1^4 \left(\frac{4}{y} - 4y^{-1/2} + 1 \right) dy \\
 &= \pi \left(4 \ln|y| - 4 \cdot \frac{2}{1} y^{1/2} + y \right) \Big|_1^4 \\
 &\approx 1.7127
 \end{aligned}$$

<https://www.geogebra.org/m/cppyhngk>

33

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$




<https://www.geogebra.org/m/cppyhngk>

34

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

Draw the region.





WASHER

<https://www.geogebra.org/m/eyabfyva>

35

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$



Right

Rotation about $x = 6$

Left


<https://www.geogebra.org/m/eyabfyva>

36

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

$x=6 \Rightarrow dy$ problem



Right $\Rightarrow y = \sqrt{x} \Rightarrow x = y^2$

Left $\Rightarrow x = 4$

BUT we are going around $x=6$

Right $\Rightarrow x = y^2 - 6$ Left $\Rightarrow x = 4 - 6$


<https://www.geogebra.org/m/eyabfyqa>

37

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

So $V = \pi \int (y^2 - 6)^2 - (4 - 6)^2 dy$



To find the bounds set Right = Left

$$y^2 = 4 \Rightarrow y = 2$$


B/c I want greater than $y=0$ b/c that a bound

<https://www.geogebra.org/m/eyabfyqa>

38

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$




$$\begin{aligned} V &= \pi \int_0^2 (y-6)^2 - (4-6)^2 dy \\ &= \pi \int_0^2 (y^2 - 12y + 36 - 4) dy \\ &= \pi \int_0^2 (y^2 - 12y + 32) dy \end{aligned}$$

<https://www.geogebra.org/m/eyabfyva>

39

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$



$$\begin{aligned} V &= \pi \left(\frac{y^3}{3} - \frac{12y^2}{2} + 32y \right) \Big|_0^2 \\ &= \frac{128\pi}{3} \end{aligned}$$


<https://www.geogebra.org/m/eyabfyva>

40

Example 3: Find the volume of the solid generated by revolving the given region about the line $y = 2$:

$$y = 2x, \quad x = 0, \quad y = 2$$

Draw the region.




<https://www.geogebra.org/m/z6tjgnn9>

41


Example 3: Find the volume of the solid generated by revolving the given region about the line $y = 2$:

$$y = 2x, \quad x = 0, \quad y = 2$$



Disk

Rotation about $y = 2$




<https://www.geogebra.org/m/z6tjgnn9>

42

Example 3: Find the volume of the solid generated by revolving the given region about the line $y = 2$:

$$y = 2x, \quad x = 0, \quad y = 2$$

$y = 2 \Rightarrow dx$ -problem



$$\begin{aligned} V &= \pi \int_0^1 (2x - 2)^2 dx \\ &= \pi \int_0^1 (4x^2 - 8x + 4) dx \\ &= \pi \left(\frac{4x^3}{3} - \frac{8x^2}{2} + 4x \right) \Big|_0^1 \\ &= \frac{4\pi}{3} \end{aligned}$$

<https://www.geogebra.org/m/z6tjan9>

43

GeoGebra Link for Lesson 16

○ <https://www.geogebra.org/m/y4pqm3mr>

○ Note click on the play buttons on the left-most screen and the animation will play/pause.

46