

# MA16020 Exam 3 Study Guide: Cal II

## Differential Equations

- Growth & Decay:  $y' = ky \Rightarrow y = Ce^{kt}$  where  $k$  is a constant.

- Separation of Variables: Solve the differential equations of the type

$$\frac{dy}{dx} = \frac{f(x)}{g(y)}$$

The idea is to try to get terms w/  $y$  on one side and  $x$ -terms on the other. Integrate and solve for  $y$ .

- First-Order Linear Differential Equations: Are equations of the form  $a(t)y' + b(t)y = c(t)$

### How to solve:

- ① Using simple algebra, rewrite your equation to be

$$y' + P(t)y = Q(t)$$

- ② Determine  $P(t)$  and  $Q(t)$

- ③ Find integrating factor:  $u(t) = \exp[\int P(t)dt]$

- ④ Plug  $u(t)$  and  $Q(t)$  in

$$y \cdot u(t) = \int Q(t)u(t)dt + C$$

- ⑤ Integrate the RHS of ④

- ⑥ Divide both sides of the equation from ⑤ by  $u(t)$ .

## Sums / Series

- Geometric Series: Are of the form  $\sum_{n=0}^{\infty} ar^n$

↳ Converge if  $|r| < 1$  and  $\sum_{n=0}^{\infty} ar^n = \frac{a}{1-r}$

↳ Diverges if  $|r| \geq 1$

- Power Series: Are of the form  $\sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$  where  $|x| < 1$

↳ Radius of convergence is  $R$  when  $|x| < R$ .

e.g.  $\sum_{n=0}^{\infty} (2x)^n \Rightarrow |2x| < 1$   
 $|x| < \frac{1}{2} \Rightarrow R = \frac{1}{2}$

- Maclaurin Series: Are of the form

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n \quad \text{where } |x| < R$$

$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ ; $\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}$	$\left[ \begin{array}{l} \text{Will be provided} \\ \text{on the exam} \end{array} \right]$
$\sin x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!}$ ; $\cos x = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$	

## Using Series to Estimate Definite Integrals

- ① Convert the function into a series
- ② Integrate the series (remember  $x$  is the variable)
- ③ Write out the number of terms to be used.
- ④ Substitute the bounds,

# Functions of Several Variables

Domain: All points  $(x, y)$  in the  $xy$ -plane for which  $f(x, y)$  is defined

Range: All values that the function  $f(x, y)$  produces

## Techniques for Finding the Domain

- Given  $\sqrt{?} \Rightarrow ? \geq 0$
- Given  $\ln(?) \Rightarrow ? > 0$
- Given  $\frac{1}{?} \Rightarrow ? \neq 0$
- Given  $\frac{1}{\sqrt{?}} \Rightarrow ? > 0$

Level Curves:  $f(x, y) = K$  where  $K$  is a constant.

Descriptions of these curves can be found on the next page.

---

## Partial & Higher Order Partial Derivatives

$f_x$   $\Rightarrow$  Find the derivative w/ respect to  $x$  and treat  $y$  as a constant.

$f_y$   $\Rightarrow$  Find the derivative w/ respect to  $y$  and treat  $x$  as a constant.

$$f_{xx} = \frac{d}{dx}(f_x) \quad f_{xy} = \frac{d}{dy}(f_x) \quad f_{yy} = \frac{d}{dy}(f_y)$$

# Descriptions of Curves

## USEFUL DEFINITIONS

1. Point at the origin  $\Rightarrow (0,0)$

2. Lines  $\Rightarrow y = mx + b$

where  $m$  is the slope  
and  $b$  is the y-intercept

3. Parabolas  $\Rightarrow y = a(x - h)^2 + k$

where  $(h, k)$  is the vertex  
of the parabola

4. Exponential Functions

a. Increasing  $\Rightarrow$  example  $y = e^x$

b. Decreasing  $\Rightarrow$  example  $y = e^{-x}$

5. Logarithmic Functions

a. Increasing  $\Rightarrow$  example  $y = \ln x$

b. Decreasing  $\Rightarrow$  example  $y = -\ln x$



6. Rational Functions are functions of the form:  $y = \frac{p(x)}{q(x)}$

a. x-axis symmetry

$$\Rightarrow f(x) = -f(x)$$

b. y-axis symmetry

$$\Rightarrow f(x) = f(-x)$$



7. Circles  $\Rightarrow (x - h)^2 + (y - k)^2 = r^2$

where  $r$  is radius and  $(h, k)$  is the center

8. Ellipses  $\Rightarrow \frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$

where  $(h, k)$  is the center

9. Hyperbolas  $\Rightarrow \frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$

where  $(h, k)$  is the center

To find the foci for 8 and 9, we use the equation  $c^2 = a^2 + b^2$ , and solve for  $c$ .