

Reminders

- WEDNESDAY QUIZ 6 on
 - Volume of Revolutions
 - Disks (Lesson 14)
 - Washers (Lesson 15)

1

MA 16020: Lesson 16 Volume By Revolution Rotation around any non-Axis

By Alexandra Cuadra

2

RECAP of Formulas from Lesson 14 and 15

For rotation around x-axis:

- Disk Method:

$$V = \pi \int_a^b [f(x)]^2 dx$$

- Washer Method:

$$V = \pi \int_a^b [R^2 - r^2] dx$$

For rotation around y-axis:

- Disk Method:

$$V = \pi \int_c^d [g(y)]^2 dy$$

- Washer Method:

$$V = \pi \int_c^d [R^2 - r^2] dy$$

3

RECAP: When do we apply Disk Method or Washer Method?

- When the region “hugs” the axis of rotation

⇒ Disk Method

- When there is a “gap” between the region and axis of rotation

⇒ Washer Method

4

Today's Lecture

- In the previous two lessons, we looked at rotations around the x-axis or y-axis.
- Today we are going to rotate about **ANY** arbitrary axis.
- Don't worry. We are going to limit ourselves to any vertical or horizontal line parallel to the x-axis or y-axis

5


Let's Backtrack a Bit...

Remember when we first described Washers, we talked about **farthest** and **closest**.

Consider the case of x-axis rotation.

In terms of distance,

- R is the length that is **FARTHEST** from x-axis
 - i.e. $R = f$
- r is the length that is **CLOSEST** to x-axis
 - i.e. $r = g$




6

When rotating around the line $y = \#$...

- Since f is the **FARTHEST**,
 - Distance b/w f and $y = \#$ is
 $R = f - \#$
- Since g is the **CLOSEST**,
 - Distance b/w g and $y = \#$ is
 $r = g - \#$
- Washer Method for around $y = \#$:

$$V = \pi \int_a^b [(R - \#)^2 - (r - \#)^2] dx$$




Note this formula is also true for the x-axis case, because the x-axis is simply the line $y = 0$

7

GOOD NEWS EVERYBODY: When rotating around the line $x = \#$...

- The same formulas, for R and r , from the case of $y = \#$ applies.
- Washer Method for around $x = \#$:

$$V = \pi \int_a^b [(R - \#)^2 - (r - \#)^2] dy$$



Note this formula is also true for the y-axis case, because the y-axis is simply the line $x = 0$

8

Note that though we did all these calculations for the Washer Problems; this idea also applies for the Disk Problems.

9

Rotation around any non-Axis Formulas

For rotation around the line $y = \#$:

○ Disk Method:

$$V = \pi \int_a^b [f(x) - \#]^2 dx$$

○ Washer Method:

$$V = \pi \int_a^b [(R - \#)^2 - (r - \#)^2] dx$$

For rotation around the line $x = \#$:

○ Disk Method:

$$V = \pi \int_c^d [g(y) - \#]^2 dy$$

○ Washer Method:

$$V = \pi \int_c^d [(R - \#)^2 - (r - \#)^2] dy$$

Note: That these formulas work for the case of x-axis ($y = 0$) and y-axis ($x = 0$).

10

Note that

- If you replace # with 0, and
- Remember that
 - x-axis $\Rightarrow y = 0$
 - y-axis $\Rightarrow x = 0$

you get the formulas from Lessons 14 and 15 which are...

11

Rotation around any Axis Formulas

For rotation around x-axis:

- Disk Method:

$$V = \pi \int_a^b [f(x)]^2 dx$$

- Washer Method:

$$V = \pi \int_a^b [R^2 - r^2] dx$$

For rotation around y-axis:

- Disk Method:

$$V = \pi \int_c^d [g(y)]^2 dy$$

- Washer Method:

$$V = \pi \int_c^d [R^2 - r^2] dy$$

12

AGAIN: When do we apply Disk Method or Washer Method?


- When the region “hugs” the axis of rotation
⇒ Disk Method
- When there is a “gap” between the region and axis of rotation
⇒ Washer Method

13

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis

Draw the region.




<https://www.geogebra.org/m/wrj2euhf>

14


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



WASHER
PROBLEM

Rotation about x-axis




<https://www.geogebra.org/m/wrj2euhf>

15


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



WASHER
PROBLEM

Furthermore, 3-D




<https://www.geogebra.org/m/wrj2euhf>

16

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x-axis $\Rightarrow dx$ problem



$$\text{Far} \Rightarrow x^2y = 4 \Rightarrow y = \frac{4}{x^2}$$

$$\text{Close} \Rightarrow y = 1$$


$$V = \pi \int \left(\frac{4}{x^2} \right)^2 - 1^2 dx$$

<https://www.geogebra.org/m/wrj2euhf>

17

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x-axis Now the bounds we are




given the smallest on $x=1$ next find the other by putting $y=1$ into $x^2y=4$

<https://www.geogebra.org/m/wrj2euhf>

18

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



$$\text{So } x^2 = 4, x = \pm 2$$

But we are looking for $x > 1$, so $x = 2$ Hence


$$V = \pi \int_1^2 \left(\frac{4}{x^2} \right)^2 - 1^2 dx$$

<https://www.geogebra.org/m/wrj2euhf>

19

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

A) the x -axis



$$V = \pi \int_1^2 \frac{16}{x^4} - 1 dx$$

$$= \pi \int_1^2 16x^{-4} - 1 dx$$


$$= \pi \left(\frac{16x^{-3}}{-3} - x \right) \Big|_1^2 = \frac{11\pi}{3}$$

<https://www.geogebra.org/m/wrj2euhf>

20

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the line $y = 1$
Draw the region.




<https://www.geogebra.org/m/n2jzwh8f>

21


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the line $y = 1$



DISK PROBLEM

Rotation about $y = 1$




<https://www.geogebra.org/m/n2jzwh8f>

22


Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the line $y = 1$



DISK PROBLEM

Furthermore, 3-D




<https://www.geogebra.org/m/n2jzwh8f>

23

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the line $y = 1$



\rightarrow dx problem

But now it is a disk problem so

$$V = \pi \int_1^2 \left(\frac{4}{x^2} - 1 \right)^2 dx$$


Note our bounds are the same as (a)

<https://www.geogebra.org/m/n2jzwh8f>

24

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

B) the line $y = 1$



$$\begin{aligned}
 V &= \pi \int_1^2 \left(\frac{16}{x^4} - \frac{8}{x^2} + 1 \right) dx \\
 &= \pi \int_1^2 (16x^{-4} - 8x^{-2} + 1) dx \\
 &= \pi \left(\frac{16x^{-3}}{-3} - \frac{8x^{-1}}{-1} + x \right) \Big|_1^2 \\
 &= 5\pi/3
 \end{aligned}$$


<https://www.geogebra.org/m/n2jzwh8f>

25

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the y -axis

Draw the region.




<https://www.geogebra.org/m/wzbn2xbt>


26

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the y -axis



Rotation about y -axis




<https://www.geogebra.org/m/wzbm2xbt>


27

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the y -axis



Furthermore, 3-D




<https://www.geogebra.org/m/wzbm2xbt>

28

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the y -axis



\Rightarrow dy problem

$$\text{Far} \Rightarrow x^2y = 4 \Rightarrow x = \sqrt{\frac{4}{y}}$$

$$\text{Close} \Rightarrow x = 1$$


$$V = \pi \int \left(\sqrt{\frac{4}{y}} \right)^2 - 1^2 dy$$

<https://www.geogebra.org/m/wzbm2xbt>

29

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the y -axis



Now the bounds we are given

the smallest one $y = 1$

Next find the other by putting

$$x = 1 \text{ into } x^2y = 4$$


$$\text{So } y = 4$$

<https://www.geogebra.org/m/wzbm2xbt>

30

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

C) the y -axis



$$\begin{aligned}
 V &= \pi \int_1^4 \left(\sqrt{\frac{4}{y}} \right)^2 - 1^2 dy \\
 &= \pi \int_1^4 \left(\frac{4}{y} - 1 \right) dy \\
 &= \pi \left(4 \ln |y| - y \right) \Big|_1^4 \\
 &\approx 7.9959
 \end{aligned}$$


<https://www.geogebra.org/m/wzbm2xbt>

31

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$

Draw the region.




<https://www.geogebra.org/m/cppyhnaq>


32

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



Rotation about $x = 1$




<https://www.geogebra.org/m/cppyhngk>


33

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



Furthermore, 3-D




<https://www.geogebra.org/m/cppyhngk>

34

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



$d y$ problem
 But now it is a disk problem so


$$V = \pi \int_1^4 \left(\sqrt{\frac{4}{y}} - 1 \right)^2 dy$$
 Note our bounds are the same as (c)

<https://www.geogebra.org/m/cppyhqk>

35

Example 1: Let R be the region of the xy -plane bounded by the curves $x^2y = 4$ below by the line $y = 1$, on the left by the line $x = 1$. Find the volume of the solid obtained by rotating R around

D) the line $x = 1$



$$\begin{aligned} V &= \pi \int_1^4 \left(\frac{4}{y} - 2\sqrt{\frac{4}{y}} + 1 \right) dy \\ &= \pi \int_1^4 \left(\frac{4}{y} - \frac{2 \cdot 2}{y^{1/2}} + 1 \right) dy \\ &= \pi \int_1^4 \left(\frac{4}{y} - 4y^{-1/2} + 1 \right) dy \\ &= \pi \left(4 \ln|y| - 4 \frac{2}{1} y^{1/2} + y \right) \Big|_1^4 \\ &\approx 1.7127 \end{aligned}$$


<https://www.geogebra.org/m/cppyhqk>

36

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

Draw the region.





<https://www.geogebra.org/m/eyabfyva>

37

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

WASHER
PROBLEM



Rotation about $x = 6$


<https://www.geogebra.org/m/eyabfyva>

38

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

$x=6 \Rightarrow dy$ problem



Far $\Rightarrow y = \sqrt{x} \Rightarrow x = y^2$
Close $\Rightarrow x = 4$

BUT we are going around $x=6$

Far $\Rightarrow x = y^2 - 6$ Close $\Rightarrow x = 4 - 6$


<https://www.geogebra.org/m/eyabfyqa>

39

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$

So $V = \pi \int (y^2 - 6)^2 - (4 - 6)^2 dy$



To find the bounds set Far = Close

$$y^2 = 4 \Rightarrow y = 2$$


B/c I want greater than $y=0$ b/c that a bound

<https://www.geogebra.org/m/eyabfyqa>

40

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$




$$\begin{aligned} V &= \pi \int_0^2 (y-6)^2 - (4-6)^2 dy \\ &= \pi \int_0^2 (y^2 - 12y + 36 - 4) dy \\ &= \pi \int_0^2 (y^2 - 12y + 32) dy \end{aligned}$$

<https://www.geogebra.org/m/eyabfyva>

41

Example 2: Find the volume of the solid generated by revolving the given region about the line $x = 6$:

$$y = \sqrt{x}, \quad y = 0, \quad x = 4$$



$$\begin{aligned} V &= \pi \left(\frac{y^3}{3} - \frac{12y^2}{2} + 32y \right) \Big|_0^2 \\ &= \frac{128\pi}{3} \end{aligned}$$


<https://www.geogebra.org/m/eyabfyva>

42

Example 3: Find the volume of the solid generated by revolving the given region about the line $y = 2$:

$$y = 2x, \quad x = 0, \quad y = 2$$

Draw the region.




<https://www.geogebra.org/m/z6tjgnn9>

43


Example 3: Find the volume of the solid generated by revolving the given region about the line $y = 2$:

$$y = 2x, \quad x = 0, \quad y = 2$$



DISK PROBLEM

Rotation about $y = 2$




<https://www.geogebra.org/m/z6tjgnn9>

44

Example 3: Find the volume of the solid generated by revolving the given region about the line $y = 2$:

$$y = 2x, \quad x = 0, \quad y = 2$$

$y = 2 \Rightarrow dx$ -problem



$$\begin{aligned} V &= \pi \int_0^1 (2x - 2)^2 dx \\ &= \pi \int_0^1 (4x^2 - 8x + 4) dx \\ &= \pi \left(\frac{4x^3}{3} - \frac{8x^2}{2} + 4x \right) \Big|_0^1 \\ &= \frac{4\pi}{3} \end{aligned}$$

<https://www.geogebra.org/m/z6tjan9>

45

GeoGebra Link for Lesson 16

○ <https://www.geogebra.org/m/y4pqm3mr>

○ Note click on the play buttons on the left-most screen and the animation will play/pause.

46