

# MA 16020 LESSON 13: VOLUME BY REVOLUTION – SHELL METHOD (SUPPEMENTAL HOMEWORK)

## Solutions

### Formulas:

- Rotating around y-axis:

$$V = 2\pi \int_a^b x \cdot (\text{Top} - \text{Bottom}) dx$$

- Rotating around  $x = \#$

- o If  $a \geq \#$ , then

$$V = 2\pi \int_a^b (x - \#) \times (\text{Top} - \text{Bottom}) dx$$

- o If  $b \leq \#$ , then

$$V = 2\pi \int_a^b (\# - x) \times (\text{Top} - \text{Bottom}) dx$$

- Rotating around x-axis:

$$V = 2\pi \int_c^d y \cdot (\text{Right} - \text{Left}) dy$$

- Rotating around  $y = \#$

- o If  $a \geq \#$ , then

$$V = 2\pi \int_a^b (y - \#) \times (\text{Right} - \text{Left}) dy$$

- o If  $b \leq \#$ , then

$$V = 2\pi \int_a^b (\# - y) \times (\text{Right} - \text{Left}) dy$$

**(OPTIONAL HOMEWORK):** Set up the integral using the **Shell Method** that represents the volume of the following solids about the given line:


1.  $x = y$ ,

$x =$

$y = 0$

about the **x-axis**

$\Rightarrow dy$



$$V = 2\pi \int y (\text{Right} - \text{Left}) dy$$

$$V = 2\pi \int_0^2 y (y - 0) dy$$

$$V = 2\pi \int_0^2 y^2 dy$$

2.  $x = 2y - y^2, \quad x = 0$

about the x-axis  $\Rightarrow dy$


Bounds.  $0 = 2y - y^2$   
 $0 = y(2 - y)$   
 $y = 0, 2$

$$V = 2\pi \int_0^2 y(2y - y^2) dy$$

3.  $y = \sqrt{x}, \quad y = -x$

about the y-axis  $\Rightarrow dx$

Bounds.  $\sqrt{x} = -x$   
 $(\sqrt{x})^2 = x^2$   
 $x = x^2$   
 $x - x^2 = 0$   
 $x(1 - x) = 0$   
 $x = 0, 1$



$$V = 2\pi \int_0^1 x(\sqrt{x} - x) dx$$

4.  $y = 2 - x^2, \quad y = x^2$

about the y-axis  $\Rightarrow dx$

Bounds:  $2 - x^2 = x^2$   
 $2 = 2x^2$   
 $1 = x^2$   
 $x = \pm 1$

$$V = 2\pi \int_{-1}^1 x(2 - x^2 - x^2) dx$$

Test Pt:  $x = 0$

$y = 2 - x^2 \rightarrow y = 2 \rightarrow \text{Top}$   
 $y = x^2 \rightarrow y = 0 \rightarrow \text{Bottom}$

$$V = 2\pi \int_{-1}^1 x(2 - 2x^2) dx$$

5.  $y = x,$


$y = x^2$

Bounds:  $x = x^2$   
 $x - x^2 = 0$   
 $x(1-x) = 0$   
 $x = 0, 1$

about  $x = -2 \rightarrow dx$   
 Since  $x = -2$  is smaller than the bounds,

$$V = 2\pi \int_0^1 (x - (-2)) [x - x^2] dx$$

$$V = 2\pi \int_0^1 (x+2)(x-x^2) dx$$



6.  $y = \sqrt{x},$


$y = 0,$

$x = 1$

about  $x = -1 \rightarrow dx$

$$V = 2\pi \int_0^1 (x - (-1)) \sqrt{x} dx$$

$$V = 2\pi \int_0^1 (x+1) \sqrt{x} dx$$




7.  $y = x,$

$y = x^2$

Bounds:  $x = x^2$   
 $x - x^2 = 0$   
 $x(1-x) = 0$   
 $x = 0, 1$

about  $x = 3 \rightarrow dx$



Since  $x = 3$  is larger than the bounds,

$$V = 2\pi \int_0^1 (3-x)(x-x^2) dx$$

8.  $y = 4x - x^2, y = 3$

Bounds:  $4x - x^2 = 3$   
 $0 = x^2 - 4x + 3$   
 $0 = (x-3)(x-1)$   
 $x = 1, 3$

Test Pt:  $x = 2$   
 $y = 4x - x^2 \rightarrow y = 4 \rightarrow \text{Top}$   
 $y = 3 \rightarrow y = 3 \rightarrow \text{Bottom}$

about  $x = 1 \Rightarrow dx$   
 Since  $x=1$  is equal to the bottom bound

$$V = 2\pi \int_1^3 (x-1)(4x-x^2-3) dx$$

9.  $x = 2y - y^2, x = 1,$

Bounds:  $2y - y^2 = 1$   
 $0 = y^2 - 2y + 1$   
 $0 = (y-1)^2$   
 $0 = y - 1$   
 $y = 1$

The other bound is given  $y = 0$

$y = 0$  about  $y = -1 \Rightarrow dy$

Test Pt:  $y = 1/2$   
 $x = 2y - y^2 \rightarrow x = 3/4 \rightarrow \text{Left}$   
 $x = 1 \rightarrow x = 1 \rightarrow \text{Right}$

Since  $y = -1$  is smaller than bounds,

$$V = 2\pi \int_0^1 (y - (-1))(1 - (2y - y^2)) dy$$

$$V = 2\pi \int_0^1 (y+1)(1-2y+y^2) dy$$

10.  $x = y^2 + 1, x = 2$

Bounds:  $y^2 + 1 = 2$   
 $y^2 = 1$   
 $y = \pm 1$

Test Pt:  $y = 0$   
 $x = y^2 + 1 \rightarrow x = 1 \rightarrow \text{Left}$   
 $x = 2 \rightarrow x = 2 \rightarrow \text{Right}$

about  $y = -2 \Rightarrow dy$

Since  $y = -2$  is smaller than the bounds ↗

$$V = 2\pi \int_{-1}^1 (y - (-2))(2 - (y^2 + 1)) dy$$

$$V = 2\pi \int_{-1}^1 (y+2)(2 - (y^2 + 1)) dy$$

$$11. x = 4y^2 - y^3, \quad x = 0$$

about  $y = 6 \Rightarrow dy$

Bounds:  $4y^2 - y^3 = 0$   
 $y^2(4 - y) = 0$   
 $y = 0, 4$

Since  $y = 6$  is larger than the bounds,

$$V = 2\pi \int_0^4 (6 - y)(4y^2 - y^3) dy$$

$$12. x = (y - 3)^2, \quad x = 4$$

about  $y = 1 \Rightarrow dy$

Bounds:  $(y - 3)^2 = 4$   
 $y - 3 = \pm 2$   
 $y = 3 \pm 2$   
 $y = 1, 5$

Test Pt:  $y = 2$   
 $x = (y - 3)^2 \rightarrow x = 1 \rightarrow \text{left}$   
 $x = 4 \rightarrow x = 4 \rightarrow \text{right}$

Since  $y = 1$  is equal to the lower bound,

$$V = 2\pi \int_1^5 (y - 1)(4 - (y - 3)^2) dy$$