

APPROXIMATION OF DEFINABLE SETS BY COMPACT FAMILIES, AND UPPER BOUNDS ON HOMOTOPY AND HOMOLOGY

ANDREI GABRIELOV AND NICOLAI VOROBJOV

ABSTRACT. We prove new upper bounds on homotopy and homology groups of o-minimal sets in terms of their approximations by compact o-minimal sets. In particular, we improve the known upper bounds on Betti numbers of semi-algebraic sets defined by quantifier-free formulae, and obtain for the first time a singly exponential bound on Betti numbers of sub-Pfaffian sets.

INTRODUCTION

We study upper bounds on topological complexity of sets definable in o-minimal structures over the reals. The fundamental case of algebraic sets in \mathbb{R}^n was first considered around 1950 by Petrovskii and Oleinik [13, 14], and then in 1960s by Milnor [12] and Thom [16]. They gave explicit upper bounds on total Betti numbers in terms of degrees and numbers of variables of the defining polynomials.

There are two natural approaches to generalizing and expanding these results. First, noticing that not much of algebraic geometry is used in the proofs, one can obtain similar upper bounds for polynomials with the “description complexity” measure different from the degree, and for non-algebraic functions, such as Khovanskii’s fewnomials and Pfaffian functions [11]. A bound for algebraic sets defined by quadratic polynomials was proved in [1].

Second, the bounds can be expanded to semialgebraic and semi-Pfaffian sets defined by formulae more general than just conjunctions of equations. Basu [2] proved an asymptotically tight upper bound on Betti numbers in the case of semialgebraic sets defined by conjunctions and disjunctions of non-strict inequalities. The proof can easily be extended to special classes of non-algebraic functions. For fewnomials and Pfaffian functions, this was done by Zell [17]. For quadratic polynomials an upper bound was proved in [3]. The principal difficulty arises when neither the set itself nor its complement is locally closed.

Until recently, the best available upper bound for the Betti numbers of a semi-algebraic set defined by an arbitrary Boolean combination of equations and inequalities remained doubly exponential in the number of variables. The first singly exponential upper bound was obtained by the authors in [9] based on a construction which replaces a given semialgebraic set by a homotopy equivalent compact semi-algebraic set. This construction extends to semi-Pfaffian sets and, more generally, to the sets defined by Boolean combinations of equations and inequalities between continuous functions definable in an o-minimal structure over \mathbb{R} . It cannot be applied to sets defined by formulae with quantifiers, such as sub-Pfaffian sets, but can

The first author was supported in part by NSF grant.

be used in conjunction with effective quantifier elimination in the semialgebraic situation.

In [10] we obtained a spectral sequence converging to the homology of the projection of an o-minimal set under the closed continuous surjective definable map. It gives an upper bound on Betti number of the projection which, in the semialgebraic case, is better than the one based on quantifier elimination. The requirement for the map to be closed can be relaxed but not completely removed, which left the upper bound problem unresolved in the general Pfaffian case, where quantifier elimination is not applicable.

In this paper we suggest a new construction for approximating a large class of definable sets, including the sets defined by arbitrary Boolean combinations of equations and inequalities, by compact sets. The construction is applicable to images of such sets under a large class of definable maps, e.g., projections. Based on this construction we refine the results from [9, 10], and prove similar upper bounds, individual for different Betti numbers, for images under arbitrary continuous definable maps.

In the semialgebraic case the bound from [9] is squaring the number of different polynomials occurring in the formula, while the bounds proved in this paper multiply the number of polynomials by a typically smaller coefficient that does not exceed the dimension. This is especially relevant for applications to problems of subspace arrangements, robotics and visualization, where the dimension and degrees usually remain small, while the number of polynomials is very large. Applied to projections, the bounds are stronger than the ones obtained by the effective quantifier elimination.

In the non-algebraic case, for the first time the bounds, singly exponential in the number of variables, are obtained for projections of semi-Pfaffian sets, as well as projections of sets defined by Boolean formulae with polynomials from special classes.

Notations. In this paper we use the following (standard) notations. For a topological space X , $H_i(X)$ is its singular homology group with coefficients in some fixed Abelian group, $\pi_i(X)$ is the homotopy group (provided that X is connected), the symbol \simeq denotes the homotopy equivalence, and the symbol \cong stands for the group isomorphism. If $Y \subset X$, then \overline{Y} denotes its closure in X .

1. MAIN RESULT

In what follows we fix an o-minimal structure over \mathbb{R} and consider sets, families of sets, maps, etc., *definable* in this structure.

Definition 1.1. Let G be a definable compact set. Consider a definable family $\{S_\delta\}_{\delta>0}$ of compact subsets of G , such that for all $\delta', \delta \in (0, 1)$, if $\delta' > \delta$, then $S_{\delta'} \subset S_\delta$. Denote $S := \bigcup_{\delta>0} S_\delta$.

For each $\delta > 0$, let $\{S_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ be a definable family of compact subsets of G such that:

- (i) for all $\varepsilon, \varepsilon' \in (0, 1)$, if $\varepsilon' > \varepsilon$, then $S_{\delta,\varepsilon} \subset S_{\delta,\varepsilon'}$;
- (ii) $S_\delta = \bigcap_{\varepsilon>0} S_{\delta,\varepsilon}$;
- (iii) for all $\delta' > 0$ sufficiently smaller than δ , and for all $\varepsilon' > 0$, there exists an open in G set $U \subset G$ such that $S_\delta \subset U \subset S_{\delta',\varepsilon'}$.

We say that S is *represented* by the families $\{S_\delta\}_{\delta>0}$ and $\{S_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ in G .

Let S' (respectively, S'') be represented by $\{S'_\delta\}_{\delta>0}$ and $\{S'_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ (respectively, by $\{S''_\delta\}_{\delta>0}$ and $\{S''_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$) in G .

Lemma 1.2. $S' \cap S''$ is represented by the families $\{S'_\delta \cap S''_\delta\}_{\delta>0}$ and $\{S'_{\delta,\varepsilon} \cap S''_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ in G , while $S' \cup S''$ is represented by $\{S'_\delta \cup S''_\delta\}_{\delta>0}$ and $\{S'_{\delta,\varepsilon} \cup S''_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ in G .

Proof. Straightforward checking of Definition 1.1. \square

Let S be represented by $\{S_\delta\}_{\delta>0}$ and $\{S_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ in G , and let $F : D \rightarrow H$ be a continuous definable map, where D and H are definable, $S \subset D \subset G$, and H is compact.

Lemma 1.3. Let D be open in G , and F be an open map. Then $F(S)$ is represented by families $\{F(S_\delta)\}_{\delta>0}$ and $\{F(S_{\delta,\varepsilon})\}_{\delta,\varepsilon>0}$ in H .

Proof. Straightforward checking of Definition 1.1 (openness is required for (iii) to hold). \square

Consider projections $\rho_1 : G \times H \rightarrow G$ and $\rho_2 : G \times H \rightarrow H$. Let $\Gamma \subset G \times H$ be the graph of F . Suppose that Γ is represented by families $\{\Gamma_\delta\}_{\delta>0}$ and $\{\Gamma_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$ in $G \times H$.

Lemma 1.4. The set $F(S)$ is represented by the families

$$\{\rho_2(\rho_1^{-1}(S_\delta) \cap \Gamma_\delta)\}_{\delta>0} \quad \text{and} \quad \{\rho_2(\rho_1^{-1}(S_{\delta,\varepsilon}) \cap \Gamma_{\delta,\varepsilon})\}_{\delta,\varepsilon>0}$$

in H .

Proof. The set $\rho_1^{-1}(S)$ is represented by the families

$$\{\rho_1^{-1}(S_\delta) \cap \Gamma_\delta\}_{\delta>0} \quad \text{and} \quad \{\rho_1^{-1}(S_{\delta,\varepsilon}) \cap \Gamma_{\delta,\varepsilon}\}_{\delta,\varepsilon>0}$$

in $G \times H$, and the projection ρ_2 satisfies Lemma 1.3. \square

Along with this general case we will be considering the following important particular cases.

Let $S = \{\mathbf{x} | \mathcal{F}(\mathbf{x})\} \subset \mathbb{R}^n$ be a bounded definable set of points satisfying a Boolean combination \mathcal{F} of equations of the kind $h(\mathbf{x}) = 0$ and inequalities of the kind $h(\mathbf{x}) > 0$, where $h : \mathbb{R}^n \rightarrow \mathbb{R}$ are continuous definable functions (e.g., polynomials). As G take a closed ball of a sufficiently large radius centered at 0. We now define the representing families $\{S_\delta\}$ and $\{S_{\delta,\varepsilon}\}$.

Definition 1.5. For a given finite set $\{h_1, \dots, h_k\}$ of functions $h_i : \mathbb{R}^n \rightarrow \mathbb{R}$ define its *sign set* as a non-empty subset in \mathbb{R}^n of the kind

$$h_{i_1} = \dots = h_{i_{k_1}} = 0, h_{i_{k_1+1}} > 0, \dots, h_{i_{k_2}} > 0, h_{i_{k_2+1}} < 0, \dots, h_{i_k} < 0,$$

where $i_1, \dots, i_{k_1}, \dots, i_{k_2}, \dots, i_k$ is a permutation of $1, \dots, k$.

Let now $\{h_1, \dots, h_k\}$ be the set of all functions in the Boolean formula defining S . Then S is a disjoint union of some sign sets of $\{h_1, \dots, h_k\}$. The set S_δ is the result of the replacement independently in each sign set in this union of all inequalities $h > 0$ and $h < 0$ by $h \geq \delta$ and $h \leq -\delta$ respectively. The set $S_{\delta,\varepsilon}$ is obtained by replacing independently in each sign set all expressions $h > 0$, $h < 0$ and $h = 0$ by $h \geq \delta$, $h \leq -\delta$ and $-\varepsilon \leq h \leq \varepsilon$, respectively. According to Lemma 1.2, the set S , being the union of sign sets, is represented by families $\{S_\delta\}$ and $\{S_{\delta,\varepsilon}\}$ in G .




FIGURE 1. The set $S_{\delta, \varepsilon}$ (right) for the closed quadrant S (left).

Example 1.6. Let the closed quadrant S be defined as the union of sign sets $\{x > 0, y > 0\} \cup \{x > 0, y = 0\} \cup \{x = 0, y > 0\} \cup \{x = y = 0\}$. Fig. 1 shows the corresponding set $S_{\delta, \varepsilon}$ for $\varepsilon < \delta$.

Now suppose that the set $S \subset \mathbb{R}^n$, defined as above by a Boolean formula \mathcal{F} , is not necessarily bounded. In this case as G take the definable one-point (Alexandrov) compactification of \mathbb{R}^n . Note that each function h is continuous in $G \setminus \{\infty\}$. Define sets S_δ and $S_{\delta, \varepsilon}$ as in the bounded case, replacing equations and inequalities independently in each sign set of $\{h_1, \dots, h_k\}$, and then taking the conjunction of the resulting formula with $|\mathbf{x}|^2 \leq 1/\delta$. Again, S is represented by $\{S_\delta\}$ and $\{S_{\delta, \varepsilon}\}$ in G , and in the sequel we will refer to this instance as the *constructible case*.

Definition 1.7. Let $\mathcal{P} := \mathcal{P}(\varepsilon_0, \dots, \varepsilon_\ell)$ be a predicate (property) over $(0, 1)^{\ell+1}$. We say that the property \mathcal{P} holds for

$$0 < \varepsilon_0 \ll \varepsilon_1 \ll \dots \ll \varepsilon_\ell \ll 1,$$

if there exist definable functions $f_k : (0, 1)^{\ell-k} \rightarrow (0, 1)$, $k = 0, \dots, \ell$ (with f_ℓ being a positive constant) such that \mathcal{P} holds for any sequence $\varepsilon_0, \dots, \varepsilon_\ell$ satisfying

$$0 < \varepsilon_k < f_k(\varepsilon_{k+1}, \dots, \varepsilon_\ell) \quad \text{for } k = 0, \dots, \ell.$$

Now we return to the general case in the Definition 1.1, which we will refer to, in what follows, as the *definable case*.

Definition 1.8. For a sequence $\varepsilon_0, \delta_0, \varepsilon_1, \delta_1, \dots, \varepsilon_m, \delta_m$, where $m \geq 0$, introduce the compact set

$$T(S) := S_{\delta_0, \varepsilon_0} \cup S_{\delta_1, \varepsilon_1} \cup \dots \cup S_{\delta_m, \varepsilon_m}.$$

From Definition 1.1 it is easy to see that for any $m \geq 0$, and for

$$(1.1) \quad 0 < \varepsilon_0 \ll \delta_0 \ll \varepsilon_1 \ll \delta_1 \ll \dots \ll \varepsilon_m \ll \delta_m \ll 1,$$

there is a surjective map $C : \mathbf{T} \rightarrow \mathbf{S}$ from the finite set \mathbf{T} of all connected components of $T(S)$ onto the set \mathbf{S} of all connected components of S , such that for any $S' \in \mathbf{S}$,

$$\bigcup_{T' \in C^{-1}(S')} T' = T(S').$$

Lemma 1.9. *If $m > 0$ then C is bijective.*

Proof. Let S be connected and $m > 0$. We prove that $T(S)$ is connected. Let $\mathbf{x}, \mathbf{y} \in S_{\delta_i, \varepsilon_i} \subset T(S)$. Let $\mathbf{x}_\varepsilon, \mathbf{y}_\varepsilon$ be a definable connected curves such that $\mathbf{x}_{\varepsilon_i} = \mathbf{x}$, $\mathbf{x}_0 := \lim_{\varepsilon \searrow 0} \mathbf{x}_\varepsilon \in S_{\delta_i}$, $\mathbf{y}_{\varepsilon_i} = \mathbf{y}$, and $\mathbf{y}_0 := \lim_{\varepsilon \searrow 0} \mathbf{y}_\varepsilon \in S_{\delta_i}$. Let $\Gamma \subset S$ be a connected compact definable curve containing \mathbf{x}_0 and \mathbf{y}_0 . Then Γ is represented by the families $\{S_\delta \cap \Gamma\}$ and $\{S_{\delta, \varepsilon} \cap \Gamma\}$ in Γ , hence $T(\Gamma) \subset T(S)$. It is easy to see that, under the condition $m > 0$, the one-dimensional $T(\Gamma)$ is connected. It follows that \mathbf{x} and \mathbf{y} belong to a connected definable curve in $T(S)$. \square

In what follows we denote $T := T(S)$, and let $m > 0$. We assume that S is connected in order to make the homotopy groups $\pi_k(S)$ and $\pi_k(T)$ independent of a base point.

Theorem 1.10. (i) *For (1.1) and every $1 \leq k \leq m$, there are epimorphisms*

$$\psi_k : \pi_k(T) \rightarrow \pi_k(S),$$

$$\varphi_k : H_k(T) \rightarrow H_k(S),$$

in particular, $\text{rank } H_k(S) \leq \text{rank } H_k(T)$.

(ii) *In the constructible case, for (1.1) and every $1 \leq k \leq m - 1$, ψ_k and φ_k are isomorphisms, in particular, $\text{rank } H_k(S) = \text{rank } H_k(T)$. Moreover, if $m \geq \dim(S)$, then $T \simeq S$.*

The plan of the proof of Theorem 1.10 is as follows. We consider a simplicial complex R in \mathbb{R}^n such that it is a triangulation of G , and S is a union of some open simplices of R . For any sequence $\varepsilon_0, \delta_0, \varepsilon_1, \delta_1, \dots, \varepsilon_m, \delta_m$ we construct a subset V of the complex R , which is a combinatorial analogy of T , and prove that there are isomorphisms of k -homotopy groups of V and S for $k \leq m - 1$ and an epimorphism for $k = m$. We prove the same for homology groups. We then show that for (1.1) there are epimorphisms $\psi_k : \pi_k(T) \rightarrow \pi_k(V)$ and $\varphi_k : H_k(T) \rightarrow H_k(V)$ for every $k \leq m$. We prove that if the pair $(R, \{S_\delta\}_{\delta > 0})$ satisfies a certain ‘‘separability’’ property (Definition 5.7), then ψ_k and φ_k are isomorphisms for every $k < m$. In particular, in the constructible case $(R, \{S_\delta\}_{\delta > 0})$ is always separable. This completes the proof.

Remark 1.11. We conjecture that in the *definable* case the statement (ii) of Theorem 1.10 is also true, i.e., for (1.1) and every $1 \leq k \leq m - 1$, the homomorphisms ψ_k, φ_k are isomorphisms, and $T \simeq S$ when $m \geq \dim(S)$.

2. TOPOLOGICAL BACKGROUND

In this section we formulate some topological definitions and statements which we will use in further proofs.

Recall that a continuous map between topological spaces $f : X \rightarrow Y$ is called a *weak homotopy equivalence* if for every $j > 0$ the induced homomorphism of homotopy groups $f_{\#j} : \pi_j(X) \rightarrow \pi_j(Y)$ is an isomorphism.

Theorem 2.1 (Whitehead Theorem on weak homotopy equivalence, [15], 7.6.24). *A map between connected CW-complexes is a weak homotopy equivalence iff it is a homotopy equivalence.*

Let $f : X \rightarrow Y$ be a continuous map between path-connected topological spaces.

Theorem 2.2 (Whitehead Theorem on homotopy and homology, [15], 7.5.9). *If there is $k > 0$ such that the induced homomorphism of homotopy groups $f_{\#j} : \pi_j(X) \rightarrow \pi_j(Y)$ is an isomorphism for $j < k$ and an epimorphism for $j = k$, then the induced homomorphism of homology groups $f_{*j} : H_j(X) \rightarrow H_j(Y)$ is an isomorphism for $j < k$ and an epimorphism for $j = k$.*

Definition 2.3 ([5]). A map $f : P \rightarrow Q$, where P and Q are posets with order relations \preceq_P and \preceq_Q respectively, is called *poset map* if, for $\mathbf{x}, \mathbf{y} \in P$, $\mathbf{x} \preceq_P \mathbf{y}$ implies $f(\mathbf{x}) \preceq_Q f(\mathbf{y})$. With a poset P is associated the simplicial complex $\Delta(P)$, called *order complex*, whose simplices are *chains* (totally ordered subsets) of P . Each poset map f induces the simplicial map $f : \Delta(P) \rightarrow \Delta(Q)$.

Theorem 2.4 ([5], Th. 2). *Let P and Q be connected posets and $f : P \rightarrow Q$ a poset map. Suppose that the fibre $f^{-1}(\Delta(Q_{\preceq q}))$ is k -connected for all $q \in Q$. Then the induced homomorphism $f_{\#j} : \pi_j(\Delta(P)) \rightarrow \pi_j(\Delta(Q))$ is an isomorphism for all $j \leq k$ and an epimorphism for $j = k + 1$.*

Remark 2.5. In the formulation and proof of this theorem in [5] the statement that $f_{\#k+1}$ is an epimorphism, is missing. Here is how it follows from the proof of Theorem 2 in [5]. In the proof, a map $g : \Delta^{(k+1)}(Q) \rightarrow \Delta(P)$ is defined, where $\Delta^{(k+1)}(Q)$ is the $(k+1)$ -dimensional skeleton of $\Delta(Q)$, such that $f \circ g : \Delta^{(k+1)}(Q) \rightarrow \Delta(Q)$ is homotopic to the identity map id . Then the induced homomorphism

$$f_{\#k+1} \circ g_{\#k+1} = (f \circ g)_{\#k+1} = id_{\#k+1} : \pi_{k+1}(\Delta^{(k+1)}(Q)) \rightarrow \pi_{k+1}(\Delta(Q))$$

is an epimorphism, since any map of a j -dimensional sphere to $\Delta(Q)$ is homotopic to a map of the sphere to $\Delta^{(j)}(Q)$. It follows that $f_{\#k+1}$ is also an epimorphism.

Corollary 2.6 (Vietoris-Begle Theorem). *Let X and Y be connected simplicial complexes and $f : X \rightarrow Y$ a simplicial map.*

- (i) *If the fibre $f^{-1}(B)$ is k -connected for every closed simplex B in Y , then the induced homomorphism $f_{\#j} : \pi_j(X) \rightarrow \pi_j(Y)$ is an isomorphism for all $j \leq k$ and an epimorphism for $j = k + 1$.*
- (ii) *If the fibre $f^{-1}(B)$ is contractible, then $X \simeq Y$.*

Proof. (i) Consider barycentric subdivisions \widehat{X} and \widehat{Y} of complexes X and Y respectively. Note that $\widehat{X} = \Delta(P)$ and $\widehat{Y} = \Delta(Q)$ where P and Q are *simplex posets* of X and Y respectively (i.e., closed simplices ordered by containment). For a closed simplex $B \in Q$ the subcomplex $\Delta(Q_{\preceq B})$ of \widehat{Y} is the union of all simplices of the barycentric subdivision of B . Now (i) follows from Theorem 2.4.

(ii) Since the fibre $f^{-1}(B)$ is contractible, according to (i), the induced homomorphisms $f_{\#j}$ are isomorphisms for all $j > 0$, hence, by Whitehead theorem on weak homotopy equivalence (Theorem 2.1), f induces the homotopy equivalence $X \simeq Y$. \square

Definition 2.7. Let Δ be a simplicial complex and X be a topological space. A map C taking simplices B to subspaces $C(B)$ of X is called *carrier* if $C(B) \subset C(K)$ for all simplices B, K in Δ such that B is a subsimplex of K . A continuous map $f : \Delta \rightarrow X$ is *carried* by C if $f(B) \subset C(B)$ for all simplices B in Δ .

Theorem 2.8 (Carrier Lemma, [5], Lemma 1). *Fix $k \geq 0$, and let $\Delta^{(k)}$ be the k -skeleton of Δ .*

- (i) If $C(B)$ is $\dim(B)$ -connected for all simplices B in $\Delta^{(k)}$, then every two maps $f, g : |\Delta^{(k)}| \rightarrow X$ that are both carried by C are homotopic, $f \sim g$.
- (ii) If $C(B)$ is $(\dim(B) - 1)$ -connected for all simplices B in $\Delta^{(k)}$, then there exists a map $|\Delta^{(k)}| \rightarrow X$ carried by C .

Definition 2.9. The *nerve* of a family $\{X_i\}_{i \in I}$ of sets is the (abstract) simplicial complex \mathcal{N} defined on the vertex set I so that a simplex $\sigma \subset I$ is in \mathcal{N} iff $\bigcap_{i \in \sigma} X_i \neq \emptyset$.

Let X be a connected regular CW-complex and $\{X_i\}_{i \in I}$ be a family of its sub-complexes such that $X = \bigcup_{i \in I} X_i$. Let $|\mathcal{N}|$ denote the geometric realization of the nerve \mathcal{N} of $\{X_i\}_{i \in I}$.

Theorem 2.10 (Nerve Theorem, [5], Th. 6). (i) *If every nonempty finite intersection $X_{i_1} \cap \dots \cap X_{i_t}$, $t \geq 1$, is $(k - t + 1)$ -connected, then there is a map $f : X \rightarrow |\mathcal{N}|$ such that the induced homomorphism $f_{\#j} : \pi_j(X) \rightarrow \pi_j(|\mathcal{N}|)$ is an isomorphism for all $j \leq k$ and an epimorphism for $j = k + 1$.*
(ii) *If every nonempty finite intersection $X_{i_1} \cap \dots \cap X_{i_t}$, $t \geq 1$, is contractible, then $X \simeq |\mathcal{N}|$.*

Remark 2.11. As with Theorem 2.4, in the formulation and proof of this theorem in [5] the statement that $f_{\#k+1}$ is an epimorphism, is missing. This statement follows from the proof of Theorem 6 in [5] by the same argument as described in Remark 2.5.

Remark 2.12. Let X be a connected triangulated set, $\{X_i\}_{i \in I}$ be a family of all of its (open) simplices, and the nerve \mathcal{N}_X is defined on the index set I so that a simplex $\sigma \subset I$ is in \mathcal{N}_X iff the family $\{X_i\}_{i \in \sigma}$, after a suitable ordering, forms a $|\sigma|$ -flag (see Definition 3.1 below). For this version of the nerve the Theorem 2.10 also holds true. Indeed, it is applicable to the union $X' := \bigcup_i X_i$ of simplices X_i in X that are contained in X with their closures (hence X' may be a proper subset of X). Since $X' \simeq X$ and $|\mathcal{N}_{X'}| \simeq |\mathcal{N}_X|$, Theorem 2.10 is also applicable to X and $\{X_i\}_{i \in I}$.

Definition 2.13. For two continuous maps $f_1 : X_1 \rightarrow Y$ and $f_2 : X_2 \rightarrow Y$, the *fibred product* is defined as

$$X_1 \times_Y X_2 := \{(\mathbf{x}_1, \mathbf{x}_2) \in X_1 \times X_2 \mid f_1(\mathbf{x}_1) = f_2(\mathbf{x}_2)\}.$$

Theorem 2.14 ([10], Th. 1). *Let $f : X \rightarrow Y$ be a continuous closed surjective o -minimal map. Then there is a spectral sequence $E_{p,q}^r$ converging to $H_*(Y)$ with $E_{p,q}^1 = H_q(W_p)$, where $W_p := \underbrace{X \times_Y \dots \times_Y X}_{p+1 \text{ times}}$.*

Corollary 2.15. *For $f : X \rightarrow Y$ as in Theorem 2.14 and for any $k \geq 0$*

$$b_k(Y) \leq \sum_{p+q=k} b_q(W_p),$$

where $b_k := \text{rank } H_k$ is the k -th Betti number.

3. SIMPLICIAL CONSTRUCTION

Since G and S are definable, they are triangulable ([6], Th. 4.4), i.e., there exists a finite simplicial complex $R = \{\Delta_j\}$ and a definable homeomorphism $\Phi : |R| \rightarrow G$,

where $|R|$ is the geometric realization of R , such that S is a union of images under Φ of some simplices of R . By a *simplex* we always mean an *open simplex*. If Δ is a simplex, then $\overline{\Delta}$ denotes its closure. In what follows we will ignore the distinction between simplices of $|R|$ and their images in G .

Definition 3.1. For a simplex Δ of S , its *subsimplex* is a simplex $\Delta' \neq \Delta$ such that $\Delta' \subset \overline{\Delta}$. A *k-flag* of simplices of R is a sequence $\Delta_{i_0}, \dots, \Delta_{i_k}$ such that Δ_{i_ν} is a subsimplex of $\Delta_{i_{\nu-1}}$ for $\nu = 1, \dots, k$.

Definition 3.2. The set S is *marked* if for every pair (Δ', Δ) of simplices of S , such that Δ' is a subsimplex of Δ , the simplex Δ' is classified as either *hard* or *soft* subsimplex of Δ . If Δ' is not in S , it is always soft.

In what follows we assume that S is marked.

Let \widehat{R} be the barycentric subdivision of R . Then each vertex v_j of \widehat{R} is the center of a simplex Δ_j of R . Let $B = B(j_0, \dots, j_k)$ be a k -simplex of \widehat{R} having vertices v_{j_0}, \dots, v_{j_k} . Assume that the vertices of B are ordered so that $\dim \Delta_{j_0} > \dots > \dim \Delta_{j_k}$. Then B corresponds to a k -flag $\Delta_{j_0}, \dots, \Delta_{j_k}$ of simplices of R . Let \widehat{S} be the set of simplices of \widehat{R} which belong to S . Then S is the union of all simplices of \widehat{S} .

Definition 3.3. The *core* $C(B)$ of a simplex $B = B(j_0, \dots, j_k)$ of \widehat{S} is the maximal subset $\{j_0, \dots, j_p\}$ of the set $\{j_0, \dots, j_k\}$ such that Δ_{j_ν} is a *hard* subsimplex of Δ_{j_μ} for all $\mu < \nu \leq p$. Note that j_0 is always in $C(B)$, in particular, $C(B) \neq \emptyset$. Assume that for a simplex B not in \widehat{S} , the core $C(B)$ is empty.

Lemma 3.4. Let $B = B(i_0, \dots, i_k)$ be a simplex in \widehat{S} , and $K = K(j_0, \dots, j_\ell)$ be a simplex in \widehat{R} , with $B \subset \overline{K}$, i.e., $I = \{i_0, \dots, i_k\} \subset J = \{j_0, \dots, j_\ell\}$. Then $I \setminus C(B) \subset J \setminus C(K)$.

Proof. Straightforward consequence of the definitions. \square

Definition 3.5. For two simplices B and B' of \widehat{S} , let $B' \succeq B$ if either B' is a subsimplex of B (reverse inclusion) and $C(B') \cap C(B) = \emptyset$, or $B' = B$. If $B' \succeq B$ and $B' \neq B$, then we write $B' \succ B$. Lemma 3.4 implies that \succeq is a partial order on the set of all simplices of \widehat{S} . The *rank* $r(\widehat{S})$ of \widehat{S} is the maximal length r of a chain $\Delta_0 \succ \dots \succ \Delta_r$ of simplices in \widehat{S} . Let B be a simplex in \widehat{S} . The set S_B of simplices $B' \subset \overline{B} \cap \widehat{S}$ is a poset with partial order induced from (\widehat{S}, \succeq) . The *rank* $r(S_B)$ of S_B is the maximal length of its chain.

Definition 3.6. Let simplices B and K be as in Lemma 3.4. For $0 < \delta < 1$, define

$$B(\delta) := \left\{ \sum_{i_\nu \in I} t_{i_\nu} v_{i_\nu} \in B(i_0, \dots, i_k) \mid \sum_{i_\nu \in C(B)} t_{i_\nu} > \delta \right\}.$$

For $0 < \varepsilon < 1$ and $0 < \delta < 1$, define

$$K_B(\delta, \varepsilon) := \left\{ \sum_{j_\nu \in J} t_{j_\nu} v_{j_\nu} \in K(j_0, \dots, j_\ell) \mid \sum_{i_\nu \in C(B)} t_{i_\nu} > \delta, \sum_{i_\nu \in I} t_{i_\nu} > 1 - \varepsilon, \right. \\ \left. \forall i_\nu \in I \forall j_\mu \in (J \setminus I) (t_{i_\nu} > t_{j_\mu}) \right\}.$$

Definition 3.7. Let B be a simplex in \widehat{S} . Fix some $m \geq 0$ and a sequence $\varepsilon_0, \delta_0, \varepsilon_1, \delta_1, \dots, \varepsilon_m, \delta_m$. Define V_B as the union of sets $K_{B'}(\delta_i, \varepsilon_i)$ over all simplices $B' \in S_B$, simplices K of \widehat{R} such that $B \subset \overline{K}$, and $i = 0, \dots, m$. Define V as the union of the sets V_B over all simplices B of \widehat{S} .

4. TOPOLOGICAL RELATIONS BETWEEN V AND S

Lemma 4.1. Let $B = B(i_0, \dots, i_k)$ be a simplex in \widehat{S} , and $K = K(j_0, \dots, j_\ell)$ be a simplex in \widehat{R} , with $B \subset \overline{K}$. Then

$$K_B(\delta, \varepsilon) \cap K_B(\delta', \varepsilon') = K_B(\max\{\delta, \delta'\}, \min\{\varepsilon, \varepsilon'\}),$$

for all $0 < \delta, \varepsilon, \delta', \varepsilon' < 1$.

Proof. Straightforward consequence of the definitions. \square

Lemma 4.2. For any two simplices B and B' of \widehat{S} , a simplex K of \widehat{R} such that B and B' are subsimplices of K , and all $0 < \delta, \varepsilon, \delta', \varepsilon' < 1$,

- (i) if $K_B(\delta, \varepsilon) \cap K_{B'}(\delta', \varepsilon') \neq \emptyset$, then either $B \subset \overline{B'}$ or $B' \subset \overline{B}$;
- (ii) $K_B(\delta, \varepsilon) \cap K_{B'}(\delta', \varepsilon')$ is convex.

Proof. Straightforward consequence of the definitions. \square

Lemma 4.3. Let K be a simplex of \widehat{R} , and let B_0, \dots, B_k be a flag of simplices of \widehat{S} , with $B_0 \subset \overline{K}$. Then for (1.1) and a sequence $i_0, j_0, \dots, i_k, j_k$ of integers in $\{0, 1, \dots, m\}$, the intersection

$$Z_K(i_0, j_0, \dots, i_k, j_k) := K_{B_0}(\delta_{i_0}, \varepsilon_{j_0}) \cap \dots \cap K_{B_k}(\delta_{i_k}, \varepsilon_{j_k})$$

is non-empty if and only if $B_\mu \succ B_\nu$ implies $j_\mu > i_\nu$ for any $\mu, \nu \in \{0, 1, \dots, k\}$.

Proof. The necessity of the condition is straightforward. To show that it is sufficient we will construct a point $v := \sum t_j v_j$, where the sum is taken over all vertices v_j of K , such that $v \in Z_K(i_0, j_0, \dots, i_k, j_k)$. This will be done in three steps.

(a) Define ℓ_ν as the last index in $C(B_\nu)$ (i.e., v_{ℓ_ν} is the center of the smallest simplex Δ_j of R such that $j \in C(B_\nu)$). Set $t_{\ell_\nu} := \delta_{i_\nu}$. If ℓ_ν is the same index for several ν , set t_{ℓ_ν} to be the maximum of the corresponding δ_{i_ν} .

(b) Fix a sequence $\gamma_0, \dots, \gamma_{k+1}$ such that $0 < \gamma_0 < \dots < \gamma_{k+1} \ll \varepsilon_0$. For a vertex v_j of $B_{\nu-1}$ which is not one of v_{ℓ_μ} and not a vertex of B_ν , set $t_j := \gamma_\nu + \max \delta_{i_\mu}$, where the maximum is taken over all μ such that $B_\nu \succ B_\mu$ (or equals 0 if there is no such μ). For any vertex v_j of K that does not belong to B_0 , set $t_j := \gamma_0$. For a vertex v_j of B_k that is not one of v_{ℓ_ν} set $t_j := \gamma_{k+1} + \max \delta_{i_\mu}$, where the maximum is taken over $\mu = 0, \dots, k$.

(c) For the last vertex v_ω of B_k set $t_\omega := 1 - \sum_{v_j} t_j$, where the sum is taken over all vertices v_j of K other than v_ω . If $\omega = \ell_k$, this overrides the setting in (a). If $\omega \neq \ell_k$, this overrides the setting in (b).

It is easy to check that $v \in Z_K(i_0, j_0, \dots, i_k, j_k)$. \square

Lemma 4.4. Let B_0, \dots, B_k be a flag of simplices of \widehat{S} , and $i_0, j_0, \dots, i_k, j_k$ be a sequence of integers in $\{0, 1, \dots, m\}$. For (1.1), if $B_\mu \succ B_\nu$ implies $j_\mu > i_\nu$ for any $\mu, \nu \in \{0, 1, \dots, k\}$, then the set

$$Z(i_0, j_0, \dots, i_k, j_k) := \bigcup_K Z_K(i_0, j_0, \dots, i_k, j_k),$$

where the union is taken over all simplices K of \widehat{R} with $B_0 \subset \overline{K}$, is an open contractible subset of G . Otherwise $Z(i_0, j_0, \dots, i_k, j_k) = \emptyset$.

Proof. By Lemma 4.3, $Z(i_0, j_0, \dots, i_k, j_k) \neq \emptyset$ if and only if $B_\mu \succ B_\nu$ implies $j_\mu > i_\nu$ for any $\mu, \nu \in \{0, 1, \dots, k\}$. So, suppose that $Z(i_0, j_0, \dots, i_k, j_k) \neq \emptyset$, and consider two simplices, K and K' , such that $B_0 \subset \overline{K'} \subset \overline{K}$. Then the intersection of the closure of the complement in K of $Z_K(i_0, j_0, \dots, i_k, j_k)$ with K' coincides with the complement in K' of $Z_{K'}(i_0, j_0, \dots, i_k, j_k)$. Hence, $Z_{K'}(i_0, j_0, \dots, i_k, j_k) \cup Z_K(i_0, j_0, \dots, i_k, j_k)$ is open in \overline{K} . It follows that $\overline{Z(i_0, j_0, \dots, i_k, j_k)}$ is open in G , and has a closed covering by convex sets $\overline{Z_K(i_0, j_0, \dots, i_k, j_k)} \cap Z(i_0, j_0, \dots, i_k, j_k)$ over all simplices K of \widehat{R} . This covering has the same nerve as the star of B_0 in the complex \widehat{R} , this star is contractible. An intersection of any number of elements of the covering of $Z(i_0, j_0, \dots, i_k, j_k)$ is convex, and therefore contractible. By the Nerve Theorem (Theorem 2.10 (ii)), both $Z(i_0, j_0, \dots, i_k, j_k)$ and the star are homotopy equivalent to the geometric realization of the nerve, and hence to one another. It follows that $Z(i_0, j_0, \dots, i_k, j_k)$ is contractible. \square

Lemma 4.5. *For (1.1) and for each simplex B in \widehat{S} and every $m \geq 1$ the set V_B (see Definition 3.7) is open in G and $(m - 1)$ -connected.*

Proof. For every simplex $B' \in S_B$ consider the set $U_{B',i} := \bigcup_K K_{B'}(\delta_i, \varepsilon_i)$, where the union is taken over all simplices K of \widehat{R} with $B \subset \overline{K}$. Obviously, the family $\{U_{B',i} \mid B' \in S_B, 0 \leq i \leq m\}$ is an open covering of V_B . Let M_B denote the nerve of this covering. From Lemmas 4.1, 4.4, M_B is the simplicial complex whose k -simplices can be identified with all sequences of the kind $((p_0, i_0), \dots, (p_k, i_k))$, where p_ν are indices of the simplices $B'_{p_\nu} \in S_B$, such that

- (a) $B'_{p_\nu} \subset \overline{B'_{p_{\nu-1}}}$,
- (b) $0 \leq i_\nu \leq m$,
- (c) if $B'_{p_\mu} \succeq B'_{p_\nu}$ and $\mu > \nu$, then $i_\mu > i_\nu$.

By Lemma 4.4 any non-empty intersection of sets $U_{B',i}$ is contractible. Therefore, due to the Nerve Theorem (Theorem 2.10 (ii)), V_B is homotopy equivalent to M_B , and in order to prove that V_B is $(m - 1)$ -connected it is sufficient to show that M_B is an $(m - 1)$ -connected simplicial complex. This follows from Proposition 4.6 below. \square


Let \succeq be a poset on $\{0, \dots, N\}$ such that if $p \succeq q$ and $p \neq q$, then $p > q$. For each $p \in \{0, \dots, N\}$, let $r(p)$ be the maximal length of a poset chain with the maximal element p (i.e., the rank of the order ideal generated by p). Let m_0, \dots, m_N be nonnegative integers. Let $M(m_0, \dots, m_N)$ be the simplicial complex containing all k -simplices $((p_0, i_0), \dots, (p_k, i_k))$ such that

- (a) $p_\nu \in \{0, \dots, N\}$, $p_0 \leq \dots \leq p_k$,
- (b) $i_\nu \in \{0, \dots, m_\nu\}$,
- (c) if $p_\mu \succeq p_\nu$ and $\mu > \nu$, then $i_\mu > i_\nu$.

Let $m := \min\{m_1, \dots, m_N\}$.

An example of the complex $M(2, 2)$ with $1 \succ 0$ is shown on Fig 2.

Proposition 4.6. *The simplicial complex $M(m_0, \dots, m_N)$ is $(m - 1)$ -connected.*


 FIGURE 2. The complex $M(2,2)$ with $1 \succ 0$.

Proof. Let Δ^N be the N -simplex, and $\overline{\Delta^N(m)}$ be the m -dimensional skeleton of its closure. There is a natural simplicial map

$$\begin{aligned} \rho : M(m_0, \dots, m_N) &\rightarrow \overline{\Delta^N}, \\ (p, i) &\mapsto p. \end{aligned}$$

It is easy to see that $\overline{\Delta^N(m)} \subset \rho(M(m_0, \dots, m_N))$, hence $\rho(M(m_0, \dots, m_N))$ is $(m-1)$ -connected.

Consider any face Δ^L of Δ^N , $L \leq N$, which has nonempty pre-image under ρ . Without loss of generality assume that its vertices are $0, \dots, L$. Let $M(m_0, \dots, m_L)$ be the simplicial complex defined over the poset on $\{0, \dots, L\}$ induced by \succeq . We prove inductively on L that for any point $\mathbf{x} \in \Delta^L$ the fibre $\rho^{-1}(\mathbf{x})$ is contractible. The proposition then follows from Vietoris-Begle Theorem (Corollary 2.6 (ii)). The base of induction, for $L = 0$, is obvious. Assume that the statement is true for $L-1$. For any simplex $K = ((p_0, i_0), \dots, (p_\ell, i_\ell))$ of $M(m_0, \dots, m_L)$ that projects surjectively onto Δ^L , if $p_\nu = L$ then $i_\nu \geq r(L)$. Let $s = i_\ell$ be the minimal of these i_ν in K , so that $p_\nu < L$ for $\nu < \ell$, while $p_\ell = L$. Then $((p_0, i_0), \dots, (p_{\ell-1}, i_{\ell-1}))$ is a simplex of the simplicial complex $M(s) := M(m'_0, \dots, m'_{L-1})$, defined over the poset on $\{0, \dots, L-1\}$ induced by \succeq , where $m'_p := \min\{m_p, r(p) + s - r(L)\}$ if $L \succeq p$, and $m'_p := m_p$ if L is incomparable with p . It follows that K is a simplex of the join of $M(s)$ and $\overline{\Delta^{m_L-s}}$, where Δ^{m_L-s} is the simplex with vertices s, \dots, m_L . Since the complex $M(s)$ is contractible due to the induction hypothesis, its join with $\overline{\Delta^{m_L-s}}$ has a contractible fibre over any $\mathbf{x} \in \Delta^L$. The fibre over \mathbf{x} of $M(m_0, \dots, m_L)$ is the union of these contractible fibres for $s = r(N), \dots, m_L$. The intersection of any number of these fibres is nonempty and contractible, being a fibre of the join of M_{\min} and $\overline{\Delta^{m_L-s_{\max}}}$. Due to the Nerve Theorem (Theorem 2.10 (ii)), their union is homotopy equivalent to its nerve, a simplex, and thus is contractible. \square

Corollary 4.7. *In the definition of the simplicial complex $M(m_0, \dots, m_N)$ assume additionally that $m_j \geq r(j)$ for every $j = 0, \dots, N$. Then $M(m_0, \dots, m_N)$ is contractible.*

Proof. The condition $m_j \geq r(j)$ guarantees that the map ρ is surjective, hence $\rho(M(m_0, \dots, m_N))$ is contractible. \square

Theorem 4.8. *For (1.1) there are homomorphisms $\chi_k : H_k(V) \rightarrow H_k(S)$ and $\tau_k : \pi_k(V) \rightarrow \pi_k(S)$ such that χ_k, τ_k are isomorphisms for every $k \leq m-1$, and χ_m, τ_m are epimorphisms. Moreover, if $m \geq \dim(S)$, then $V \simeq S$.*

Proof. Due to Lemma 4.2(i), for any three simplices B_0, B_1, B_2 in \widehat{S} , the equality $\overline{B_0} = \overline{B_1} \cap \overline{B_2}$ is equivalent to $V_{B_0} = V_{B_1} \cap V_{B_2}$. Hence, a nonempty intersection

of any number of sets V_B is a set of the same type, and therefore is $(m - 1)$ -connected. Moreover, there is an isomorphism $\xi : |\mathcal{N}_V| \rightarrow |\mathcal{N}_{\widehat{S}}|$ between the geometric realization of the nerve $\mathcal{N}_{\widehat{S}}$ of the covering of \widehat{S} by its simplices and the geometric realization of the nerve \mathcal{N}_V of the open covering of V by sets V_B .

Since intersections of any number of elements of the covering of \widehat{S} (i.e., simplices) are contractible if non-empty, the Nerve Theorem (Theorem 2.10 (ii) and Remark 2.12) implies that $\widehat{S} \simeq |\mathcal{N}_{\widehat{S}}|$, i.e., there is a continuous map $\psi_{\widehat{S}} : \widehat{S} \rightarrow |\mathcal{N}_{\widehat{S}}|$ which induces isomorphisms of homotopy groups $\psi_{\widehat{S}\#} : \pi_k(\widehat{S}) \rightarrow \pi_k(|\mathcal{N}_{\widehat{S}}|)$ for all integers $k \geq 0$.

On the other hand, by the Nerve Theorem (Theorem 2.10 (i)), there is a continuous map $\psi_V : V \rightarrow |\mathcal{N}_V|$ inducing isomorphisms of homotopy groups $\psi_{V\#} : \pi_k(V) \rightarrow \pi_k(|\mathcal{N}_V|)$ for every $k \leq m - 1$ and an epimorphism $\psi_{V\#} : \pi_m(V) \rightarrow \pi_m(|\mathcal{N}_V|)$. As τ_k take

$$\psi_{\widehat{S}\#}^{-1} \circ \xi \circ \psi_{V\#} : \pi_k(V) \rightarrow \pi_k(\widehat{S}).$$

By Whitehead Theorem on homotopy and homology (Theorem 2.2), $\psi_{\widehat{S}}$ induces isomorphisms of homology groups $\psi_{\widehat{S}*} : H_k(\widehat{S}) \rightarrow H_k(|\mathcal{N}_{\widehat{S}}|)$ for all $k \geq 0$, while ψ_V induces isomorphisms of homology groups $\psi_{V*} : H_k(V) \rightarrow H_k(|\mathcal{N}_V|)$ for every $k \leq m - 1$, and an epimorphism $\psi_{V*} : H_m(V) \rightarrow H_m(|\mathcal{N}_V|)$. As χ_k take

$$\psi_{\widehat{S}*}^{-1} \circ \xi \circ \psi_{V*} : H_k(V) \rightarrow H_k(\widehat{S}).$$

If $m \geq \dim(S)$ then, by Corollary 4.7, a nonempty intersection of any number of sets V_B is contractible. Then, according to Nerve Theorem, sets V and \widehat{S} are homotopy equivalent to geometric realizations of the respective nerves, and therefore $V \simeq S$. \square

5. PROOF OF THEOREM 1.10

We now need to re-define the simplicial complex R so that it would satisfy additional properties. Recall that definable functions are triangulable [6], Th. 4.5. Consider a finite simplicial complex R' such that R' is a triangulation of the projection

$$\rho : G \times [0, 1] \rightarrow [0, 1],$$

and R' is compatible with

$$\bigcup_{\delta \in (0,1)} (S_\delta, \delta) \subset G \times [0, 1].$$

Define R as the triangulation induced by R' on the fibre $\rho^{-1}(0)$.

Definition 5.1. Along with the sequence $\varepsilon_0, \dots, \delta_m$, consider another sequence $\varepsilon'_0, \delta'_0, \varepsilon'_1, \delta'_1 \dots \varepsilon'_m, \delta'_m$. Let T' be the set defined as in Definition 1.8 replacing all δ_i, ε_i by $\delta'_i, \varepsilon'_i$. Let V' be the set defined as in Definition 3.7 replacing all δ_i, ε_i by $\delta'_i, \varepsilon'_i$.

5.1. Definable case. In the definable case we specify the hard-soft relation for the set V' as follows. For any pair (Δ_1, Δ_2) of S such that Δ_1 is a subsimplex of Δ_2 , we assume that Δ_1 is *soft* in Δ_2 .

Definition 5.2. Let B and K be as in Lemma 3.4. For $0 < \varepsilon < 1$ define

$$K_B(\varepsilon) := \left\{ \sum_{j_\nu \in J} t_{j_\nu} v_{j_\nu} \in K(j_0, \dots, j_\ell) \mid \sum_{i_\nu \in I} t_{i_\nu} > 1 - \varepsilon, \right. \\ \left. \forall i_\nu \in I \forall j_\mu \in (J \setminus I) (t_{i_\nu} > t_{j_\mu}) \right\}.$$

Introduce a new parameter ε'' , and define V'' as the union of $K_B(\varepsilon'')$ over all simplices B of \widehat{S} , and all simplices K of \widehat{R} such that $B \subset \overline{K}$.

In each of the following Lemmas 5.3, 5.4, 5.5, and 5.6, the statement holds for

$$0 < \varepsilon'_0 \ll \dots \ll \varepsilon'_i \ll \varepsilon_i \ll \delta_i \ll \delta'_i \ll \dots \ll \delta'_m \ll \varepsilon'' \quad (i = 0, \dots, m).$$

Lemma 5.3. $S \simeq V''$.

Proof. Let B be a simplex of \widehat{S} , and let $U_B := \bigcup K_B(\varepsilon'')$, where the union is taken over all simplices K of \widehat{R} such that $B \subset \overline{K}$. Then the family of all sets U_B forms an open covering of V'' whose nerve we denote by $\mathcal{N}_{V''}$. Each U_B is contractible, since B is a deformation retract of U_B . Any intersection $U := U_{B_0} \cap \dots \cap U_{B_k}$ is nonempty iff, after the suitable reordering, the sequence B_{i_1}, \dots, B_{i_k} is a k -flag of simplices. If $U \neq \emptyset$, then B_{i_k} is its deformation retract, hence U is contractible. By Nerve Theorem (Theorem 2.10 (ii)), $V'' \simeq |\mathcal{N}_{V''}|$. On the other hand, the simplices of \widehat{S} form a covering of S with nerve \mathcal{N}_S (in the sense of Remark 2.12), therefore, $S \simeq |\mathcal{N}_S|$ by Nerve Theorem. Then $S \simeq V''$, since nerves $\mathcal{N}_{V''}$ and \mathcal{N}_S are isomorphic. \square

Lemma 5.4. $V' \subset T \subset V''$.

Proof. Let $V_{\delta, \varepsilon}$ be the union of sets $K_B(\delta, \varepsilon)$ over all simplices K of \widehat{R} and simplices B of \widehat{S} such that $B \subset \overline{K}$. We first show that $V_{\delta', \varepsilon'} \subset S_{\delta, \varepsilon}$ which immediately implies $V' \subset T$. Fix δ' , and let $\mathbf{x}_{\varepsilon'} \in V_{\delta', \varepsilon'}$ be a definable curve. Then $\mathbf{x}_0 := \lim_{\varepsilon' \searrow 0} \mathbf{x}_{\varepsilon'} \in \overline{B(\delta')}$, where B is a simplex in \widehat{S} (this follows from Definition 2.6). Let Δ be the simplex in S containing \mathbf{x}_0 . Since every subsimplex of Δ is soft in Δ , $\mathbf{x}_0 \in S_\delta$ for $\delta \ll \delta'$. Also an open neighborhood of \mathbf{x}_0 in G , of the size independent of ε' , is contained in $S_{\delta, \varepsilon}$ for $\varepsilon \ll \delta \ll \delta'$. Hence $\mathbf{x}_{\varepsilon'} \in S_{\delta, \varepsilon}$ for $\varepsilon' \ll \varepsilon \ll \delta \ll \delta'$.

Next, we show that $S_{\delta, \varepsilon} \subset V''$, and therefore $T \subset V''$. Fix δ , and let $\mathbf{x}_\varepsilon \in S_{\delta, \varepsilon}$ be a definable curve. Then $\mathbf{x}_0 := \lim_{\varepsilon \searrow 0} \mathbf{x}_\varepsilon \in S_\delta$. Hence \mathbf{x}_0 belongs to a simplex B of \widehat{S} . According to Definition 5.2, an open neighbourhood of \mathbf{x}_0 of the radius larger than ε is contained in $K_B(\varepsilon'')$ for any simplex K of \widehat{R} such that $B \subset \overline{K}$. In particular, $\mathbf{x}_\varepsilon \in K_B(\varepsilon'')$ and therefore $\mathbf{x}_\varepsilon \in V''$. \square

Lemma 5.5. *The inclusion map $\iota : V' \hookrightarrow V''$ induces isomorphisms of homotopy groups $\iota_{k\#} : \pi_k(V') \rightarrow \pi_k(V'')$ for every $k \leq m - 1$, and an epimorphism $\iota_{m\#}$.*

Proof. Recall that V' admits an open covering by sets of the kind $V'_B := V_B$ (Definition 3.7), over all simplices B in \widehat{S} , such that every non-empty intersection of sets V'_B is $(m - 1)$ -connected (Lemma 4.5). Similarly, the set V'' has an open covering by sets V''_B , where V''_B is the union of sets $K_{B'}(\varepsilon'')$ over all simplices $B' \in S_B$, and simplices K of \widehat{R} such that $B \subset \overline{K}$. Every non-empty intersection of sets V''_B is contractible (cf. the proof of Lemma 5.3).

The inclusion relation $V'_B \subset V''_B$ implies that these two coverings have the same nerve \mathcal{N} , up to isomorphism. Let $\mathcal{N}^{(m)}$ be the m -skeleton of \mathcal{N} . Following the proof of Theorem 6 in [5], we now describe a *carrier* (see Definition 2.7) C' assigning certain intersections of the sets V'_B to simplices σ of the barycentric subdivision of $\mathcal{N}^{(m)}$.

Let Q be the face poset of $\mathcal{N}^{(m)}$ (i.e., simplices ordered by subsimplex relation), and $\Delta(Q)$ its order complex (see Definition 2.3). Then $\Delta(Q)$ is homeomorphic to $|\mathcal{N}^{(m)}|$ being its barycentric subdivision. For σ in $\Delta(Q)$ let

$$C'(\sigma) = \bigcap_{V'_B \in \min \sigma} V'_B.$$

In a similar way a carrier C'' from simplices in the barycentric subdivision of \mathcal{N} to intersections of sets V''_B is defined.

According to Carrier Lemma (ii) (Theorem 2.8), there exist continuous maps $g' : \mathcal{N}^{(m)} \rightarrow V'$ and $g'' : \mathcal{N} \rightarrow V''$ such that g' is carried by C' and g'' is carried by C'' . On the other hand, g' is also carried by C'' because $V'_B \subset V''_B$ implies $g'(\sigma) \subset C'(\sigma) \subset C''(\sigma)$. Since all non-empty intersections of V''_B are contractible, in particular m -connected, Carrier Lemma (i) implies that

$$(5.1) \quad \iota \circ g' \sim g''|_{\mathcal{N}^{(m)}}.$$

Due to Nerve Theorem (Theorem 2.10 (ii), details in [5]) g'' is a homotopy equivalence. Passing to homomorphisms of homotopy groups, we have that $g''_{\#k} : \pi_k(\mathcal{N}) \rightarrow \pi_k(V'')$ is an isomorphism for all k hence, by (5.1), $\iota_{\#k}$ is an epimorphism for all $k \leq m$. According to Nerve Theorem, $g'_{\#k} : \pi_k(\mathcal{N}^{(m)}) \rightarrow \pi_k(V')$ is an isomorphism for $k \leq m-1$, thus, $\iota_{\#k}$ is also a monomorphism for $k \leq m-1$. \square

Lemma 5.6. *For every $k \leq m$, there are epimorphisms $\zeta_k : \pi_k(T) \rightarrow \pi_k(V'')$ and $\eta_k : H_k(T) \rightarrow H_k(V'')$.*

Proof. Due to Lemmas 5.4, 5.5,

$$V' \xrightarrow{p} T \xrightarrow{q} V'',$$

where \hookrightarrow are the inclusion maps, and $q \circ p$ induces isomorphisms $(q \circ p)_{\#} = q_{\#} \circ p_{\#}$ of homotopy groups $\pi_k(V') \cong \pi_k(V'')$ for every $k \leq m-1$, and an epimorphism $\pi_m(V') \rightarrow \pi_m(V'')$. Then $\zeta_k := q_{\#}$ is an epimorphism for every $k \leq m$.

By Whitehead Theorem on homotopy and homology (Theorem 2.2), $q \circ p$ also induces isomorphisms $(q \circ p)_* = q_* \circ p_*$ of homology groups $H_k(V') \cong H_k(V'')$ for every $k \leq m-1$, and an epimorphism $H_m(V') \rightarrow H_m(V'')$. Hence, $\eta_k := q_*$ is an epimorphism for every $k \leq m$. \square

Theorem 1.10(i) immediately follows from Lemmas 5.3 and 5.6.

5.2. Separability and constructible case.

Definition 5.7. For the simplicial complex R and the family $\{S_\delta\}_{\delta>0}$ we call the pair $(R, \{S_\delta\}_{\delta>0})$ *separable* if for any pair (Δ_1, Δ_2) of simplices of S such that Δ_1 is a subsimplex of Δ_2 , the equality $\Delta_2 \cap S_\delta \cap \Delta_1 = \emptyset$ is equivalent to the inclusion $\Delta_1 \subset \Delta_2 \setminus S_\delta$ for all sufficiently small $\delta > 0$.

Recall that in the constructible case we assume that S is defined by a Boolean combination of equations and inequalities with continuous definable functions, and the set S_δ is defined using sign sets of these functions (see Section 1).

Lemma 5.8. *In the constructible case $(R, \{S_\delta\}_{\delta>0})$ is separable.*

Proof. Observe that R is compatible with the sign set decomposition of S .

Consider a pair (Δ_1, Δ_2) of simplices of S such that Δ_1 is a subsimplex of Δ_2 . If both Δ_1 and Δ_2 lie in the same sign set, then $\overline{\Delta_2 \cap S_\delta} \cap \Delta_1 = \overline{\Delta_1 \cap S_\delta} \neq \emptyset$ and $\Delta_1 \not\subset \overline{\Delta_2 \setminus S_\delta}$.

If Δ_1 and Δ_2 lie in two different sign sets, then there is a function h in the Boolean combination defining S such that $h(\mathbf{x}) = 0$ for every point $\mathbf{x} \in \Delta_1$, while $h(\mathbf{y})$ satisfies a strict inequality, say $h(\mathbf{y}) > 0$, for every point $\mathbf{y} \in \Delta_2$. Then $\overline{\Delta_2 \cap S_\delta} \subset \overline{\Delta_2 \cap \{h \geq \delta\}}$ and $\overline{\Delta_2 \cap \{h \geq \delta\}} \cap \{h = 0\} = \emptyset$. Hence $\overline{\Delta_2 \cap S_\delta} \cap \Delta_1 = \emptyset$. On the other hand, $\overline{\Delta_2 \setminus S_\delta} \supset \overline{\Delta_2 \cap \{h < \delta\}} \supset \overline{\Delta_2 \cap \{h = 0\}} \supset \Delta_1$. \square

Now we return to the general definable case, and assume for the rest of this section that $(R, \{S_\delta\})$ is separable. For any pair (Δ_1, Δ_2) of simplices of S such that Δ_1 is a subsimplex of Δ_2 , we assume that Δ_1 is *soft* in Δ_2 if $\overline{\Delta_2 \cap S_\delta} \cap \Delta_1 = \emptyset$ (equivalently, $\Delta_1 \subset \overline{\Delta_2 \setminus S_\delta}$) for all sufficiently small $\delta > 0$. Otherwise, Δ_1 is *hard* in Δ_2 .

Lemma 5.9. *If Δ_1 is hard in Δ_2 , then for every $\mathbf{x} \in \Delta_1$ there is a neighbourhood $U_{\mathbf{x}}$ of \mathbf{x} in $\overline{\Delta_2}$ such that for all sufficiently small $\delta \in (0, 1)$, $U_{\mathbf{x}} \subset \overline{\Delta_2 \cap S_\delta}$.*

Proof. Suppose that contrary to the claim, for some $\mathbf{x} \in \Delta_1$, $U_{\mathbf{x}} \setminus \overline{\Delta_2 \cap S_\delta} \neq \emptyset$ for any neighbourhood $U_{\mathbf{x}}$ of \mathbf{x} in Δ_1 , for arbitrarily small $\delta > 0$.

Since the set S_δ grows (with respect to inclusion) as $\delta \searrow 0$, and Δ_1 is hard in Δ_2 , the intersection $\overline{\Delta_2 \cap S_\delta} \cap \Delta_1$ is non-empty and also grows. If for any neighbourhood $W_{\mathbf{x}}$ of \mathbf{x} in Δ_1 , $W_{\mathbf{x}} \not\subset \overline{\Delta_2 \cap S_\delta} \cap \Delta_1$ for arbitrarily small $\delta > 0$, then the limits of both $\overline{\Delta_2 \cap S_\delta} \cap \Delta_1$ and its complement in Δ_1 , as $\delta \searrow 0$, have non-empty intersections with Δ_1 . This contradicts to the assumption that Δ_1 is a simplex in the complex R compatible with R' , thus there is a neighbourhood $W_{\mathbf{x}}$ in Δ_1 such that $W_{\mathbf{x}} \subset \overline{\Delta_2 \cap S_\delta} \cap \Delta_1$ for sufficiently small $\delta > 0$. It follows that $U_{\mathbf{x}} \setminus \overline{\Delta_2 \cap S_\delta} \subset \Delta_2$. Since $\mathbf{x} \in \overline{\Delta_2 \setminus S_\delta}$, and using again the compatibility of the complex R with R' , we conclude that $\Delta_1 \subset \overline{\Delta_2 \setminus S_\delta}$, i.e., Δ_1 is soft in Δ_2 , which is a contradiction. \square

In each of the following Lemmas 5.10, 5.11, and Theorem 5.12 the statement holds for

$$(5.2) \quad 0 < \varepsilon'_0 \ll \dots \ll \varepsilon'_i \ll \varepsilon_i \ll \delta_i \ll \delta'_i \ll \dots \ll \delta'_m \ll 1 \quad (i = 0, \dots, m).$$

Lemma 5.10. *$T' \subset V$ and $V' \subset T$.*

Proof. We show first that $S_{\delta', \varepsilon'} \subset V_{\delta, \varepsilon}$, for $\varepsilon' \ll \varepsilon \ll \delta \ll \delta'$, where $V_{\delta, \varepsilon}$ is the union of $K_B(\delta, \varepsilon)$ over all simplices K of \widehat{R} and simplices B of \widehat{S} such that $B \subset \overline{K}$.

Let us fix δ' , and let $\mathbf{x}_{\varepsilon'} \in S_{\delta', \varepsilon'}$ be any definable curve. It is enough to show that $\mathbf{x}_{\varepsilon'} \in V_{\delta, \varepsilon}$ for $\varepsilon' \ll \varepsilon \ll \delta \ll \delta'$. Clearly, $\mathbf{x}_0 =: \lim_{\varepsilon' \searrow 0} \mathbf{x}_{\varepsilon'}$ belongs to $S_{\delta'}$. Hence \mathbf{x}_0 belongs to a simplex $B = B(j_0, \dots, j_\ell)$ of \widehat{S} . Suppose that $\mathbf{x}_0 \notin B(\delta)$. Let $\mathbf{x}_{0, \delta} \in B \setminus B(\delta)$ be a definable curve. Then $\mathbf{x}_{0, 0} =: \lim_{\delta \searrow 0} \mathbf{x}_{0, \delta}$ belongs to a subsimplex $B' = B(i_0, \dots, i_k)$ of B . It follows that $\mathbf{x}_{0, 0} \in \overline{\Delta_{j_0} \cap S_{\delta'} \cap \Delta_{i_0}}$, therefore Δ_{i_0} is *hard* in Δ_{j_0} . On the other hand, by the definition of $B(\delta)$ (Definition 3.6), Δ_{i_0} is *soft* in Δ_{j_0} . This contradiction shows that $\mathbf{x}_0 \in B(\delta)$.

For $\varepsilon' \ll \varepsilon$, the distance from $\mathbf{x}_{\varepsilon'}$ to $\mathbf{x}_0 \in S_{\delta'} \cap B$ is much smaller than ε . From Definition 2.6, for $\varepsilon \ll \delta \ll \delta'$, the union of $K(\delta, \varepsilon)$ over all simplices K of \widehat{R} such

that $B \subset \overline{K}$ contains an open in G neighborhood of $\mathbf{x}_0 \in B$ of the size that is independent of ε' . Hence $\mathbf{x}_{\varepsilon'} \in V_{\delta, \varepsilon}$ for $\varepsilon' \ll \varepsilon \ll \delta \ll \delta'$.

Next, we want to show that $V_{\delta', \varepsilon'} \subset S_{\delta, \varepsilon}$. As before, fix δ' . Let $\mathbf{x}_{\varepsilon'} \in V_{\delta', \varepsilon'}$ be a definable curve. Then $\mathbf{x}_0 := \lim_{\varepsilon' \searrow 0} \mathbf{x}_{\varepsilon'} \in B(\delta')$, where $B = B(j_0, \dots, j_\ell)$ is a simplex in \widehat{S} (this follows from Definition 3.6). Suppose that $\mathbf{x}_0 \notin S_\delta$, then $\mathbf{x}_0 \in B \setminus S_\delta$. Let $\mathbf{x}_{0, \delta} \in B \setminus S_\delta$ be a definable curve. Therefore $\mathbf{x}_{0,0} := \lim_{\delta \searrow 0} \mathbf{x}_{0, \delta}$ belongs to a subsimplex $B' = B(i_0, \dots, i_k)$ of B . Then by Lemma 5.9, Δ_{i_0} is *soft* in Δ_{j_0} , and thus $\mathbf{x}_{0,0} \notin V_{\delta', \varepsilon'}$. The same is true for $\mathbf{x}_{\varepsilon'}$ as well, namely $\mathbf{x}_{\varepsilon'} \notin V_{\delta', \varepsilon'}$ for $\varepsilon' \ll \delta \ll \delta'$. This contradiction shows that $\mathbf{x}_0 \in S_\delta$.

Since $\mathbf{x}_0 \in S$, an open neighborhood of \mathbf{x}_0 in G , of the size independent of ε' , is contained in $S_{\delta, \varepsilon}$ for $\varepsilon \ll \delta \ll \delta'$. Hence $\mathbf{x}_{\varepsilon'} \in S_{\delta, \varepsilon}$ for $\varepsilon' \ll \varepsilon \ll \delta \ll \delta'$. \square

Lemma 5.11. *The inclusion maps $T' \hookrightarrow T$ and $V' \hookrightarrow V$ are homotopy equivalences.*

Proof. Proofs of homotopy equivalences are similar for the both inclusions, so we will consider only the case of $T' \hookrightarrow T$.

Consider $\varepsilon_0, \delta_0, \dots, \varepsilon_m, \delta_m$ as variables, then $T \subset \mathbb{R}^{n+2m+2}$. From the o-minimal version of Hardt's triviality, applied to the projection $\rho : T \rightarrow \mathbb{R}^{2m+2}$ on the subspace of coordinates $\varepsilon_0, \delta_0, \dots, \varepsilon_m, \delta_m$, follows the existence of a partition of \mathbb{R}^{2m+2} into a finite number of connected definable sets $\{A_i\}$ such that T is definably trivial over each A_i , i.e., for any point $(\bar{\varepsilon}, \bar{\delta}) := (\varepsilon_0, \delta_0, \dots, \varepsilon_m, \delta_m) \in A_i$ the pre-image $\rho^{-1}(A_i)$ is definably homeomorphic to $\rho^{-1}(\bar{\varepsilon}, \bar{\delta}) \times A_i$ by a fibre preserving homeomorphism.

There exists an element A_{i_0} of the partition which is an open connected set in \mathbb{R}^{2m+2} and contains both points $(\bar{\varepsilon}, \bar{\delta})$ and $(\bar{\varepsilon}', \bar{\delta}')$ for (5.2). Let $\gamma : [0, 1] \rightarrow A_{i_0}$ be a definable simple curve such that $\gamma(0) = (\bar{\varepsilon}, \bar{\delta})$ and $\gamma(1) = (\bar{\varepsilon}', \bar{\delta}')$. Then $\rho^{-1}(\gamma(0)) = T$, $\rho^{-1}(\gamma(1)) = T'$ and $\rho^{-1}(\gamma([0, 1]))$ is definably homeomorphic to $T \times \gamma([0, 1])$. Let $\Phi_{t, t'} : \rho^{-1}(\gamma(t')) \rightarrow \rho^{-1}(\gamma(t))$ for $0 \leq t \leq t' \leq 1$ be the homeomorphism of fibres. Replacing if necessary $(\bar{\varepsilon}, \bar{\delta})$ by a point closer to $(\bar{\varepsilon}', \bar{\delta}')$ along the curve γ , we can assume that $\rho^{-1}(\gamma(t')) \subset \rho^{-1}(\gamma(t))$ for all $0 \leq t \leq t' \leq 1$. Then T' is a strong deformation retract of T defined by the homotopy $F : T \times [0, 1] \rightarrow T$ as follows. If $\mathbf{x} \in \rho^{-1}(\gamma(t'))$ for some $t' \leq t$ and $\mathbf{x} \notin \rho^{-1}(\gamma(t''))$ for any $t'' > t'$, then $F(\mathbf{x}, t) = \Phi_{t', t}(\mathbf{x})$. If $\mathbf{x} \in \rho^{-1}(\gamma(t'))$ with $t' > t$, then $F(\mathbf{x}, t) = \mathbf{x}$. \square

Theorem 5.12. $T \simeq V$

Proof. Consider four sequences $(\varepsilon^{(j)}, \delta^{(j)}) := (\varepsilon_0^{(j)}, \delta_0^{(j)}, \dots, \varepsilon_m^{(j)}, \delta_m^{(j)})$, $1 \leq j \leq 4$. Let $T(\varepsilon^{(j)}, \delta^{(j)})$ (respectively, $V(\varepsilon^{(j)}, \delta^{(j)})$) be the set defined as in Definition 1.8 (resp., Definition 3.7) replacing all δ_i, ε_i by $\delta_i^{(j)}, \varepsilon_i^{(j)}$.

Due to Lemmas 5.10, the following chain of inclusions holds

$$T(\varepsilon^{(1)}, \delta^{(1)}) \xrightarrow{p} V(\varepsilon^{(2)}, \delta^{(2)}) \xrightarrow{q} T(\varepsilon^{(3)}, \delta^{(3)}) \xrightarrow{r} V(\varepsilon^{(4)}, \delta^{(4)}),$$

for

$$0 < \varepsilon_0^{(j)} \ll \delta_0^{(j)} \ll \dots \ll \varepsilon_m^{(j)} \ll \delta_m^{(j)} \ll 1,$$

where

$$\delta_{i-1}^{(j-1)} \ll \varepsilon_i^{(j-1)} \ll \varepsilon_i^{(j)} \ll \delta_i^{(j)} \ll \delta_i^{(j-1)}$$

for all $i = 1, \dots, m$, $j = 2, 3, 4$.

According to Lemma 5.11, $q \circ p$ and $r \circ q$ are homotopy equivalences. Passing to induced homomorphisms of homotopy groups, we have that $(q \circ p)_\# = q_\# \circ p_\#$ is an

isomorphism, hence $q_{\#}$ is epimorphism. Similarly, since $(r \circ q)_{\#} = r_{\#} \circ q_{\#}$ is an isomorphism, $q_{\#}$ is a monomorphism. It follows that $q_{\#}$ is an isomorphism, therefore $T \simeq V$ by Whitehead Theorem on weak homotopy equivalence (Theorem 2.1). \square

Theorem 1.10(ii) immediately follows from Theorems 5.12 and 4.8.

6. UPPER BOUNDS ON BETTI NUMBERS

The method described in this section can be applied to obtain upper bounds on Betti numbers for sets defined by Boolean formulae with functions from various classes which admit a natural measure of “description complexity” and a suitable version of “Bezout Theorem”, most notably for semialgebraic and semi- and sub-Pfaffian sets (see, e.g., [8]). We give detailed proofs for the semialgebraic case. The proofs can be extended to the Pfaffian case straightforwardly.

Definition 6.1. Let $f, g, h : \mathbb{N}^{\ell} \rightarrow \mathbb{N}$ be three functions, $n \in \mathbb{N}$. The expression $f \leq O(g)^n$ means: there exists $c \in \mathbb{N}$ such that $f \leq (cg)^n$ everywhere on \mathbb{N}^{ℓ} . The expression $f \leq g^{O(h)}$ means: there exists $c \in \mathbb{N}$ such that $f \leq g^{ch}$ everywhere on \mathbb{N}^{ℓ} .

6.1. Semialgebraic sets defined by quantifier-free formulae. Consider the constructible case with $S = \{\mathbf{x} \mid \mathcal{F}(\mathbf{x})\} \subset \mathbb{R}^n$, where \mathcal{F} is a Boolean combination of polynomial equations and inequalities of the kind $h(\mathbf{x}) = 0$ or $h(\mathbf{x}) > 0$, $h \in \mathbb{R}[x_1, \dots, x_n]$. Suppose that the number of different polynomials h is s and their degrees do not exceed d . The following upper bounds on the total Betti number $b(S)$ of the set S originate from the classic works of [13, 14, 12, 16]. Their proofs can be found in [4].

- (i) If \mathcal{F} is a conjunction of any number equations, then $b(S) \leq d(2d - 1)^{n-1}$.
- (ii) If \mathcal{F} is a conjunction of s *non-strict* inequalities, then $b(S) \leq (sd + 1)^n$.
- (iii) If \mathcal{F} is a conjunction of s equations and *strict* inequalities, then $b(S) \leq O(sd)^n$.

The following statement applies to more general semialgebraic sets.

Theorem 6.2 ([2], Th. 1; [4], Th. 7.38). *If \mathcal{F} is a monotone Boolean combination (i.e., exclusively connectives \wedge, \vee are used, no negations) of only strict or only non-strict inequalities, then $b(S) \leq O(sd)^n$.*

In [9], Th. 1 the authors proved the bound $b(S) \leq O(s^2d)^n$ for an arbitrary Boolean formula \mathcal{F} . Theorem 1.10 implies the following refinement of this bound.

Theorem 6.3. *Let $\nu := \min\{k + 1, n - k, s\}$. Then the k -th Betti number*

$$b_k(S) \leq O(\nu sd)^n.$$

Proof. Assume first that $k > 0$. For $m = k$ construct $T(S)$ in the compactification of \mathbb{R}^n , as described in Section 1. $T(S)$ is a compact set defined by a Boolean formula with $4(k + 1)s$ polynomials in $\mathbb{R}[x_1, \dots, x_n]$ of the kind $h + \delta_i, h - \delta_i, h + \varepsilon_i$ or $h - \varepsilon_i$, $0 \leq i \leq k$, having degrees at most d . According to Lemma 1.9, there is a bijection C from the set \mathbf{T} of all connected components of $T(S)$ to the set \mathbf{S} of all connected components of S such that $C^{-1}(S') = T(S')$ for every $S' \in \mathbf{S}$. By Theorem 1.10 (i), $b_k(S') \leq b_k(T(S'))$. It follows that

$$b_k(S) = \sum_{S' \in \mathbf{S}} b_k(S') \leq \sum_{S' \in \mathbf{S}} b_k(T(S')) = b_k(T(S)).$$

Then, applying the bound from Theorem 6.2 to $T(S)$,

$$(6.1) \quad b_k(S) \leq b_k(T(S)) \leq O((k+1)sd)^n.$$

On the other hand, since $T(S)$ is compact, $b_k(T(S)) = b_{n-k-1}(\mathbb{R}^n \setminus T(S))$ by Alexander's duality. The semialgebraic set $\mathbb{R}^n \setminus T(S)$ is defined by a monotone Boolean combination of only strict inequalities, hence, due to Theorem 6.2,

$$(6.2) \quad b_k(S) \leq b_{n-k-1}(\mathbb{R}^n \setminus T(S)) \leq O((n-k)sd)^n.$$

The theorem now follows from (6.1), (6.2) and the bound $b(S) \leq O(s^2d)^n$ from [9].

In the case $k = 0$, $b_0(S) \leq b_0(T(S))$ since the map C is surjective, hence by Theorem 6.2,

$$b_0(S) \leq b_0(T(S)) \leq O(sd)^n.$$

□

6.2. Projections of semialgebraic sets. Let $\rho : \mathbb{R}^{n+r} \rightarrow \mathbb{R}^n$ be the projection map, and $S = \{(\mathbf{x}, \mathbf{y}) \mid \mathcal{F}(\mathbf{x}, \mathbf{y})\} \subset \mathbb{R}^{n+r}$ be a semialgebraic set, where \mathcal{F} is a Boolean combination of polynomial equations and inequalities of the kind $h(\mathbf{x}, \mathbf{y}) = 0$ or $h(\mathbf{x}, \mathbf{y}) > 0$, $h \in \mathbb{R}[x_1, \dots, x_n, y_1, \dots, y_r]$. Suppose that the number of different polynomials h is s and their degrees do not exceed d .

Effective quantifier elimination algorithm ([4], Ch. 14) produces a Boolean combination \mathcal{F}_ρ of polynomial equations and inequalities, with polynomials in $\mathbb{R}[x_1, \dots, x_n]$, defining the projection $\rho(S)$. The number of different polynomials in \mathcal{F}_ρ is $(sd)^{O(nr)}$, and their degrees are bounded by $d^{O(r)}$. Then Theorem 6.3 (or Theorem 1 in [9]) implies that

$$(6.3) \quad b_k(\rho(S)) \leq (sd)^{O(n^2r)}$$

for any $k \geq 0$. We now improve this bound as follows.

Theorem 6.4. *The k -th Betti number of $\rho(S)$ satisfies the inequality*

$$b_k(\rho(S)) \leq \sum_{0 \leq p \leq k} O((p+1)(k+1)sd)^{n+(p+1)r} \leq ((k+1)sd)^{O(n+kr)}.$$

Proof. For $k = 0$ the bound immediately follows from Theorem 6.3, so assume that $k > 0$. The set S is represented by families $\{S_\delta\}_\delta$, $\{S_{\delta,\varepsilon}\}_{\delta,\varepsilon}$ in the compactification of \mathbb{R}^{n+r} as described in Section 1. According to Lemma 1.3, the projection $\rho(S)$ is represented by families $\{\rho(S_\delta)\}_\delta$, $\{\rho(S_{\delta,\varepsilon})\}_{\delta,\varepsilon}$ in the compactification of \mathbb{R}^n . Fix $m = k$, then the set $T(\rho(S)) = \rho(T(S))$ is defined. According to Corollary 2.15,

$$b_k(\rho(T(S))) \leq \sum_{p+q=k} b_q(W_p),$$

where

$$W_p = \underbrace{T(S) \times_{\rho(T(S))} \cdots \times_{\rho(T(S))} T(S)}_{p+1 \text{ times}}.$$

The fibred product $W_p \subset \mathbb{R}^{n+(p+1)r}$ is definable by a Boolean formula with

$$4(p+1)(k+1)s$$

polynomials of degrees not exceeding d . Hence, by Theorem 6.2,

$$b_q(W_p) \leq O((p+1)(k+1)sd)^{n+(p+1)r}.$$

It follows that

$$(6.4) \quad b_k(T(\rho(S))) \leq \sum_{0 \leq p \leq k} O((p+1)(k+1)sd)^{n+(p+1)r} \leq ((k+1)sd)^{O(n+kr)}.$$

Finally, by Theorem 1.10 (i), $b_k(\rho(S)) \leq b_k(T(\rho(S)))$, which, in conjunction with (6.4), completes the proof. \square

6.3. Semi- and sub-Pfaffian sets. Necessary definitions regarding semi-Pfaffian and sub-Pfaffian sets can be found in [8, 7] (see also [11]).

Let $S = \{\mathbf{x} \mid \mathcal{F}(\mathbf{x})\} \subset (0, 1)^n$ be a semi-Pfaffian set, where \mathcal{F} is a Boolean combination of equations and inequalities with s different Pfaffian functions (here and in the sequel $(0, 1)$ can be replaced by any, bounded or unbounded, interval). Assume that all functions are defined in $(0, 1)^n$, have a common Pfaffian chain of order ℓ , and degree (α, β) . A straightforward generalization of Theorem 6.2 gives the following upper bound.

Theorem 6.5 ([17], Th. 1; [8], Th. 3.4). *If \mathcal{F} is a monotone Boolean combination of only strict or only non-strict inequalities such that $\overline{S} \subset (0, 1)^n$, then*

$$b(S) \leq s^n 2^{\ell(\ell-1)/2} O(n\beta + \min\{n, \ell\}\alpha)^{n+\ell}.$$

In conjunction with Theorem 1.10 this implies the following bound for the set S defined by an *arbitrary* Boolean formula \mathcal{F} .

Theorem 6.6. *Let $\nu := \min\{k+1, n-k, s\}$. Then the k -th Betti number*

$$b_k(S) \leq (\nu s)^n 2^{\ell(\ell-1)/2} O(n\beta + \min\{n, \ell\}\alpha)^{n+\ell}.$$

Proof. Analogous to the proof of Theorem 6.3. \square

Remark 6.7. Unlike Theorem 6.5, the condition $\overline{S} \subset (0, 1)^n$ is not required in Theorem 6.6, since taking the conjunction of inequalities $0 < x_i < 1$, for $i = 1, \dots, n$, with \mathcal{F} , guarantees that the closed set $T(S) \subset (0, 1)^n$.

Now we consider the sub-Pfaffian case. Let $\rho : \mathbb{R}^{n+r} \rightarrow \mathbb{R}^n$ be the projection map, and $S = \{(\mathbf{x}, \mathbf{y}) \mid \mathcal{F}(\mathbf{x}, \mathbf{y})\} \subset (0, 1)^{n+r}$ be a semi-Pfaffian set, where \mathcal{F} is a Boolean combination of Pfaffian equations and inequalities. Suppose that all different Pfaffian functions occurring in \mathcal{F} are defined in $(0, 1)^{n+r}$, have a common Pfaffian chain of order ℓ , their number is s , and their degree is (α, β) . Since the Pfaffian o-minimal structure does not admit quantifier elimination (i.e., the projection of a semi-Pfaffian set may not be semi-Pfaffian, see [8]), it is not possible to apply in the Pfaffian case the same method that we used to obtain the bound (6.3). On the other hand, the method employed in the proof of Theorem 6.4 extends straightforwardly to projections of semi-Pfaffian sets, and produces the following first general singly exponential upper bound for Betti numbers of sub-Pfaffian sets.

Theorem 6.8. *The k -th Betti number of $\rho(S)$ satisfies the inequality*

$$b_k(\rho(S)) \leq (ks)^{O(n+(k+1)r)} 2^{O(k\ell)^2} ((n+(k+1)r)(\alpha+\beta))^{n+(k+1)r+k\ell}.$$

Proof. Analogous to the proof of Theorem 6.4. \square

Acknowledgements. We thank J. McClure for useful discussions. Part of this research was carried out during our joint visit in Spring 2007 to Institute for Mathematics and Its Applications at University of Minnesota, under the program Applications of Algebraic Geometry, to which we are very grateful.

REFERENCES

- [1] A. Barvinok, On the Betti numbers of semi-algebraic sets defined by few quadratic inequalities, *Math. Z.*, 225 (1997), 231–244.
- [2] S. Basu, On bounding the Betti numbers and computing the Euler characteristics of semialgebraic sets, *Discrete Comput. Geom.*, 22 (1999), 1–18.
- [3] S. Basu, D. Pasechnik, and M.-F. Roy, Betti numbers of semi-algebraic sets defined by partly quadratic systems of polynomials, arXiv: 0708.3522.
- [4] S. Basu, R. Pollack, and M.-F. Roy, *Algorithms in Real Algebraic Geometry*, Second Edition, Springer-Verlag, Berlin, 2006.
- [5] A. Björner, Nerves, fibers and homotopy groups, *J. Combinatorial Theory*, Ser. A, 102 (2003), 88–93.
- [6] M. Coste, *An Introduction to O-minimal Geometry*, Dip. Mat. Univ. Pisa, Dottorato di Ricerca in Matematica, Istituti Editoriale e Poligrafici Internazionali, Pisa, 2000.
- [7] A. Gabrielov, N. Vorobjov, Complexity of cylindrical decompositions of sub-Pfaffian sets, *Journal of Pure and Applied Algebra*, 164 (2001), 179–197.
- [8] A. Gabrielov, N. Vorobjov, Complexity of computations with Pfaffian and Noetherian functions, in: *Normal Forms, Bifurcations and Finiteness Problems in Differential Equations*, 211–250, Kluwer, 2004.
- [9] A. Gabrielov, N. Vorobjov, Betti numbers of semialgebraic sets defined by quantifier-free formulae, *Discrete Comput. Geom.*, 33 (2005), 395–401.
- [10] A. Gabrielov, N. Vorobjov, and T. Zell, Betti numbers of semialgebraic and sub-Pfaffian sets, *J. London Math. Soc.*, 69-1 (2004), 27–43.
- [11] A. Khovanskii, *Fewnomials*, Translations of Mathematical Monographs 88, AMS, Providence, RI, 1991.
- [12] J. Milnor, On the Betti numbers of real varieties, *Proc. Amer. Math. Soc.*, 15 (1964), 275–280.
- [13] O. A. Oleinik, Estimates of the Betti numbers of real algebraic hypersurfaces (Russian), *Mat. Sb.*, 28 (1951), 635–640.
- [14] I. G. Petrovskii, O. A. Oleinik, On the topology of real algebraic hypersurfaces (Russian), *Izv. Acad. Nauk SSSR*, 13 (1949), 389–402. English transl.: *Amer. Math. Soc. Transl.*, 7 (1962), 399–417.
- [15] E. Spanier, *Algebraic Topology*, Springer-Verlag, New York-Berlin, 1981.
- [16] R. Thom, Sur l’homologie des variétés algébriques réelles, in: *Differential and Combinatorial Topology*, 255–265, Princeton University Press, Princeton, 1965.
- [17] T. Zell, Betti numbers of semi-Pfaffian sets, *Journal of Pure and Applied Algebra*, 139 (1999), 323–338.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA
E-mail address: `agabriel@math.purdue.edu`

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF BATH, BATH BA2 7AY, ENGLAND, UK
E-mail address: `nnv@cs.bath.ac.uk`