


MIXED POLYNOMIAL/POWER SERIES RINGS AND RELATIONS AMONG THEIR SPECTRA

William Heinzer

joint with Christel Rotthaus and Sylvia Wiegand

Department of Mathematics

Purdue University




OVERVIEW 1

At first glance the rings

$$B := k[[y]] [x] \quad \text{and} \quad C := k[x] [[y]]$$

look similar. One has



OVERVIEW 1

At first glance the rings

$$B := k[[y]] [x] \quad \text{and} \quad C := k[x] [[y]]$$

look similar. One has

$$B = k[[y]] [x] \hookrightarrow k[x] [[y]] = C,$$

but this is a strict inclusion.

OVERVIEW 1

At first glance the rings

$$B := k[[y]][x] \quad \text{and} \quad C := k[x][[y]]$$

look similar. One has


$$B = k[[y]][x] \hookrightarrow k[x][[y]] = C,$$

but this is a strict inclusion.

For example, $1 - xy$ is a nonunit of B , and


$$\frac{1}{1 - xy} = \sum_{i=0}^{\infty} x^i y^i \in C.$$

So $1 - xy$ is a unit of C .



CONCLUSION

Indeed, the rings $B = k[[y]][x]$ and $C = k[x][[y]]$ are not isomorphic: the intersection of the maximal ideals of B is (0) , while y is in every maximal ideal of C .




Overview 2

Consider the mixed polynomial/power series rings

$$k[x, y] \hookrightarrow k[[y]][x] \hookrightarrow k[x][[y]] \hookrightarrow k[[x, y]],$$

where k is a field. The inclusion maps here are all flat homomorphisms. The prime ideal structure of these rings is well understood. The above inclusions induce maps



Overview 2


Consider the mixed polynomial/power series rings

$$k[x, y] \hookrightarrow k[[y]][x] \hookrightarrow k[x][[y]] \hookrightarrow k[[x, y]],$$

where k is a field. The inclusion maps here are all flat homomorphisms. The prime ideal structure of these rings is well understood. The above inclusions induce maps

$$\mathrm{Spec} A \longleftarrow \mathrm{Spec} B \longleftarrow \mathrm{Spec} C \longleftarrow \mathrm{Spec} D.$$

We are interested in describing these Spec maps.



Overview 3

Consider

$$k[x][[y]] = C \hookrightarrow C[1/x] \hookrightarrow k[x, 1/x][[y]] := E,$$

Overview 3

Consider

$$k[x][[y]] = C \hookrightarrow C[1/x] \hookrightarrow k[x, 1/x][[y]] := E,$$

At first glance, it appears that E is a localization of C , but it is not. There are elements in E that are not in the fraction field of C . However, E is obtained from C by the localization $C[1/x]$ followed by the (y) -adic completion of $C[1/x]$. Thus E is flat over C . The map $C \hookrightarrow E$ induces $\text{Spec } C \leftarrow \text{Spec } E$, and again we are interested in describing this Spec map.

Overview 4

Also consider

$$C_1 := k[x] \left[\left[\frac{y}{x} \right] \right] \hookrightarrow \dots \hookrightarrow C_n := k[x] \left[\left[\frac{y}{x^n} \right] \right] \hookrightarrow \dots \hookrightarrow E.$$

Overview 4


Also consider

$$C_1 := k[x] \left[\left[\frac{y}{x} \right] \right] \hookrightarrow \cdots \hookrightarrow C_n := k[x] \left[\left[\frac{y}{x^n} \right] \right] \hookrightarrow \cdots \hookrightarrow E.$$

The maps $C \hookrightarrow C_n$ and $C_i \hookrightarrow C_n$ for $i < n$ are not flat, but $C_n \hookrightarrow E = k[x, 1/x] \left[\left[y \right] \right]$ is the localization $C_n[1/x]$ followed by the (y) -adic completion of $C_n[1/x]$. Thus $C_n \hookrightarrow E$ is flat. These inclusion maps induce maps

$$\text{Spec } C \leftarrow \text{Spec } C_1 \leftarrow \cdots \leftarrow \text{Spec } C_n \leftarrow \cdots \leftarrow \text{Spec } E.$$

We are interested in describing these Spec maps.



Generic fiber rings

Let $R \hookrightarrow S$ be an injective homomorphism of commutative rings with R an integral domain.


DEFINITION. The **generic fiber ring** of the map $R \hookrightarrow S$ is the localization $(R \setminus \{0\})^{-1}S$ of S .

Generic fiber rings

Let $R \hookrightarrow S$ be an injective homomorphism of commutative rings with R an integral domain.

DEFINITION. The **generic fiber ring** of the map $R \hookrightarrow S$ is the localization $(R \setminus \{0\})^{-1}S$ of S .

With $A := k[x, y] \hookrightarrow B := k[[y]][x] \hookrightarrow C := k[x][[y]] \hookrightarrow D := k[[x, y]]$, the generic fiber ring of $A \hookrightarrow R$ is one-dim. for $R \in \{B, C, D\}$, while the generic fiber ring of $R \hookrightarrow S$ is zero-dim for $R \subseteq S$ in $\{B, C, D\}$.



Trivial generic fiber extensions

Let R be a subring of an integral domain S .

Definition. $R \hookrightarrow S$ is a **trivial generic fiber** extension or a **TGF** extension if

$$(0) \neq P \in \operatorname{Spec} S \implies P \cap R \neq (0).$$

Trivial generic fiber extensions

Let R be a subring of an integral domain S .


Definition. $R \hookrightarrow S$ is a **trivial generic fiber** extension or a **TGF** extension if

$$(0) \neq P \in \text{Spec } S \implies P \cap R \neq (0).$$

A TGF extension S of R is gotten via

$$R \hookrightarrow T \rightarrow T/P := S,$$

where T is an extension ring of R and $P \in \text{Spec } T$ is maximal with respect to $P \cap R = (0)$. Thus the generic fiber ring of $R \hookrightarrow T$ is relevant to constructing TGF extensions S of R .



A TGF Extension

Let x and y be indeterminates over a field k . Then

$$R := k[[x, y]] \hookrightarrow S := k[[x]] \left[\left[\frac{y}{x} \right] \right] \quad \text{is TGF.}$$

A TGF Extension

Let x and y be indeterminates over a field k . Then

$$R := k[[x, y]] \hookrightarrow S := k[[x]] \left[\left[\frac{y}{x} \right] \right] \quad \text{is TGF.}$$

Proof. It suffices to show $P \cap R \neq (0)$ for each $P \in \text{Spec } S$ with $\text{ht } P = 1$. This is clear if $x \in P$, while if $x \notin P$, then $k[[x]] \cap P = (0)$, so $k[[x]] \hookrightarrow R/(P \cap R) \hookrightarrow S/P$. Now S/P is one-dim local with residue field k . Hence by Cohen's Theorem 8, S/P is finite over $k[[x]]$. Thus $\dim R/(P \cap R) = 1$, so $P \cap R \neq (0)$.

Cohen's Theorem 8

Theorem (Classical) Let I be an ideal of a ring R and let M be an R -module. Assume that R is complete in the I -adic topology and $\bigcap_{n=1}^{\infty} I^n M = (0)$. If M/I is generated over R/I by elements $\bar{w}_1, \dots, \bar{w}_s$ and w_i is a preimage in M of \bar{w}_i for $1 \leq i \leq s$, then M is generated over R by w_1, \dots, w_s .


Cohen's Theorem 8

Theorem (Classical) Let I be an ideal of a ring R and let M be an R -module. Assume that R is complete in the I -adic topology and $\bigcap_{n=1}^{\infty} I^n M = (0)$. If M/I is generated over R/I by elements $\bar{w}_1, \dots, \bar{w}_s$ and w_i is a preimage in M of \bar{w}_i for $1 \leq i \leq s$, then M is generated over R by w_1, \dots, w_s .

This is useful for proving that with

$$B := k[[y]][x] \hookrightarrow C := k[x][[y]] \hookrightarrow D := k[[x, y]],$$

then $R \hookrightarrow S$ is TGF for $R \subseteq S$ in $\{B, C, D\}$.



TGF Extensions


PROP. 1. Let $R \hookrightarrow S$ and $S \hookrightarrow T$ be injective maps, where R , S and T are integral domains.

- (1) If $R \hookrightarrow S$ and $S \hookrightarrow T$ are TGF extensions, then so is $R \hookrightarrow T$. Equivalently if $R \hookrightarrow T$ is not TGF, then at least one of the extensions $R \hookrightarrow S$ or $S \hookrightarrow T$ is not TGF.

TGF Extensions

PROP. 1. Let $R \hookrightarrow S$ and $S \hookrightarrow T$ be injective maps, where R , S and T are integral domains.

- (1) If $R \hookrightarrow S$ and $S \hookrightarrow T$ are TGF extensions, then so is $R \hookrightarrow T$. Equivalently if $R \hookrightarrow T$ is not TGF, then at least one of the extensions $R \hookrightarrow S$ or $S \hookrightarrow T$ is not TGF.
- (2) If $R \hookrightarrow T$ is TGF, then $S \hookrightarrow T$ is TGF.
- (3) If the map $\text{Spec } T \rightarrow \text{Spec } S$ is surjective, then $R \hookrightarrow T$ is TGF implies $R \hookrightarrow S$ is TGF.



A NON-TGF EXTENSION

PROP. 2. $R = k[[x]][y, z] \hookrightarrow k[y, z][[x]] = S$ is not TGF.

A NON-TGF EXTENSION


PROP. 2. $R = k[[x]][y, z] \hookrightarrow k[y, z][[x]] = S$ is not TGF.

Proof. There exists $\sigma \in k[y][[x]]$ that is transcendental over $k[[x]][y]$. Let $\mathfrak{q} = (z - \sigma x)k[y, z][[x]]$ and define $\pi : k[y, z][[x]] \rightarrow k[y, z][[x]] / \mathfrak{q} \cong k[y][[x]]$. Thus $\pi(z) = \sigma x$. If $h \in \mathfrak{q} \cap (k[[x]][y, z])$, then $\exists s, t \in \mathbb{N}$ so that $h = \sum_{i=0}^s \sum_{j=0}^t (\sum_{\ell \in \mathbb{N}} a_{ij\ell} x^\ell) y^i z^j$, where $a_{ij\ell} \in k$.

A NON-TGF EXTENSION

PROP. 2. $R = k[[x]][y, z] \hookrightarrow k[y, z][[x]] = S$ is not TGF.

Proof. There exists $\sigma \in k[y][[x]]$ that is transcendental over $k[[x]][y]$. Let $\mathfrak{q} = (z - \sigma x)k[y, z][[x]]$ and define $\pi : k[y, z][[x]] \rightarrow k[y, z][[x]] / \mathfrak{q} \cong k[y][[x]]$. Thus $\pi(z) = \sigma x$. If $h \in \mathfrak{q} \cap (k[[x]][y, z])$, then $\exists s, t \in \mathbb{N}$ so that $h = \sum_{i=0}^s \sum_{j=0}^t (\sum_{\ell \in \mathbb{N}} a_{ij\ell} x^\ell) y^i z^j$, where $a_{ij\ell} \in k$. Hence $0 = \pi(h) = \sum_{i=0}^s \sum_{j=0}^t (\sum_{\ell \in \mathbb{N}} a_{ij\ell} x^\ell) y^i (\sigma x)^j$. Since σ is transcendental over $k[[x]][y]$, each $a_{ij\ell} = 0$. Therefore $\mathfrak{q} \cap (k[[x]][y, z]) = (0)$, and $R \hookrightarrow S$ is not TGF.



Power Series Rings 1


Lemma. Let $R[[y]]$ denote the power series ring in the variable y over the commutative ring R . Then

- (1) Each maximal ideal of $R[[y]]$ has the form $(\mathfrak{m}, y)R[[y]]$, where \mathfrak{m} is a maximal ideal of R . Thus y is in every maximal ideal of $R[[y]]$.

Power Series Rings 1

Lemma. Let $R[[y]]$ denote the power series ring in the variable y over the commutative ring R . Then

- (1) Each maximal ideal of $R[[y]]$ has the form $(\mathfrak{m}, y)R[[y]]$, where \mathfrak{m} is a maximal ideal of R . Thus y is in every maximal ideal of $R[[y]]$.
- (2) If R is Noetherian with $\dim R[[y]] = n$ and x_1, \dots, x_m are indeterminates over $R[[y]]$, then y is in every maximal ideal of height $n + m$ of the polynomial ring $R[[y]][x_1, \dots, x_m]$.



Power Series Rings 2


Lemma. Let R be an n -dim. Noetherian domain and let \mathfrak{q} be a prime ideal of height n in the power series ring $R[[y]]$. If $y \notin \mathfrak{q}$, then \mathfrak{q} is contained in a unique maximal ideal of $R[[y]]$.

Power Series Rings 2

Lemma. Let R be an n -dim. Noetherian domain and let \mathfrak{q} be a prime ideal of height n in the power series ring $R[[y]]$. If $y \notin \mathfrak{q}$, then \mathfrak{q} is contained in a unique maximal ideal of $R[[y]]$.


Proof. Let $S := R[[y]]/\mathfrak{q}$. The assertion is clear if \mathfrak{q} is maximal. Otherwise, $\dim S = 1$. Moreover, S is complete in its yS -adic topology and every maximal ideal of S is a minimal prime of the principal ideal yS . Hence S is a complete semilocal ring. Since S is also an integral domain, it is local by [Mat., Theorem 8.15]. Thus \mathfrak{q} is contained in a unique maximal ideal of $R[[y]]$.

Spec $k[[y]][x]$




β is the cardinality of $k[[y]]$, and α is the cardinality of the set of maximal ideals of $k[x]$; the boxed β means there are cardinality β height-one primes in that position with respect to the partial ordering.

$\text{Spec } k[x][[y]]$



Here α is the cardinality of the set of maximal ideals of $k[x]$, and β is the uncountable cardinal equal to the cardinality of $k[[y]]$.

$\text{Spec } R[[y]]$



$\text{Spec } R[[y]]$ for R a one-dim Noetherian domain

Here κ_i and κ_j are uncountable cardinals.

Isomorphic Spectra

REMARK. Let F be a field that is algebraic over a finite field. Roger Wiegand proved that as partially ordered sets or topological spaces

$$\operatorname{Spec} \mathbb{Q}[x, y] \not\cong \operatorname{Spec} F[x, y] \cong \operatorname{Spec} \mathbb{Z}[y].$$

The spectra of power series extensions in y behave differently: we have

$$\operatorname{Spec} \mathbb{Z}[[y]] \cong \operatorname{Spec} \mathbb{Q}[x][[y]] \cong \operatorname{Spec} F[x][[y]].$$

Higher dimension

We display several extensions involving three variables:

$$k[[z]] [x, y] \xrightarrow{\beta} k[x] [[z]] [y] \xrightarrow{\gamma} k[x, y] [[z]] \xrightarrow{\delta} k[x] [[y, z]],$$
$$k[[z]] [x, y] \xrightarrow{\varepsilon} k[[y, z]] [x] \xrightarrow{\zeta} k[x] [[y, z]] \xrightarrow{\eta} k[[x, y, z]],$$

We have been able to show many of these extensions are not TGF.

ANOTHER NON TGF EXTENSION

PROP. 3. $k[[z]][x, y] \xrightarrow{\beta} k[x][[z]][y]$ is not TGF.

ANOTHER NON TGF EXTENSION

PROP. 3. $k[[z]][x, y] \xrightarrow{\beta} k[x][[z]][y]$ is not TGF.

Proof. Fix $\sigma \in k[x][[z]]$ that is transcendental over $k[[z]][x]$. Define $\pi : k[x][[z]][y] \rightarrow k[x][[z]]$ to be the identity map on $k[x][[z]]$ and $\pi(y) = \sigma z$. Let $\mathfrak{q} = \ker \pi$. Then $y - \sigma z \in \mathfrak{q}$. If $h \in \mathfrak{q} \cap (k[[z]][x, y])$, then $h =$

ANOTHER NON TGF EXTENSION

PROP. 3. $k[[z]][x, y] \xrightarrow{\beta} k[x][[z]][y]$ is not TGF.

Proof. Fix $\sigma \in k[x][[z]]$ that is transcendental over $k[[z]][x]$. Define $\pi : k[x][[z]][y] \rightarrow k[x][[z]]$ to be the identity map on $k[x][[z]]$ and $\pi(y) = \sigma z$. Let $\mathfrak{q} = \ker \pi$. Then $y - \sigma z \in \mathfrak{q}$. If $h \in \mathfrak{q} \cap (k[[z]][x, y])$, then $h =$

$$\sum_{j=0}^s \sum_{i=0}^t \left(\sum_{\ell \in \mathbb{N}} a_{ij\ell} z^\ell \right) x^i y^j, \text{ for } s, t \in \mathbb{N} \text{ and } a_{ij\ell} \in k.$$

THEREFORE

$$0 = \pi(h) = \sum_{j=0}^s \sum_{i=0}^t \left(\sum_{\ell \in \mathbb{N}} a_{ij\ell} z^\ell \right) x^i (\sigma z)^j =$$

$$\sum_{j=0}^s \sum_{i=0}^t \left(\sum_{\ell \in \mathbb{N}} a_{ij\ell} z^{\ell+j} \right) x^i \sigma^j.$$

Since σ is trans. over $k[[z]][x]$, x and σ are alg. indep. over $k((z))$. Thus each $a_{ij\ell} = 0$. Therefore $\mathfrak{q} \cap (k[[z]][x, y]) = (0)$, and the embedding β is not TGF.

Question

Is $k[x, y] [[z]] \xrightarrow{\theta} k[x, y, 1/x] [[z]]$ TGF?

REMARK. For k a field and x, y, u and z indeterminates over k , the extension

$k[x, y, u] [[z]] \hookrightarrow k[x, y, u, 1/x,] [[z]]$ is not TGF.