

Outlines of sol's.

1. Use Laplace transform : $X(s) = \mathcal{L}\{x(t)\}$

$$s^3 X(s) + 4s^2 X(s) + 4s X(s) = \frac{1}{s+2}$$

$$\Rightarrow X(s) = \frac{1}{(s+2)^3 s}$$


partial fractions:

$$\frac{s+3}{(s+2)^4 s} = \frac{A}{s} + \frac{B}{s+2} + \frac{C}{(s+2)^2} + \frac{D}{(s+2)^3}$$

$$\Rightarrow \dots \quad \Rightarrow \text{tables}$$


$$x(t) = \frac{1}{8} e^{-2t} (-1 - 2t - 2t^2) + \frac{1}{8}$$

2.



$$f * g(t) = \int_0^t f(\tau) g(t-\tau) d\tau = \begin{cases} \int_0^t \cos(t-\tau) d\tau & t \leq \pi \\ \int_0^\pi \cos(t-\tau) d\tau & t > \pi \end{cases}$$

$$= \begin{cases} \sin(t) & t \leq \pi \\ 2\sin(t) & t > \pi \end{cases}$$



$$\begin{aligned}
 3.a) f * g &= \int_0^t \cos(\alpha\tau) \cos(t-\tau) d\tau \\
 &= \int_0^t \cos(\alpha\tau) (\cos(t) \cos(\tau) - \sin(t) \sin(\tau)) d\tau \\
 &= \cos(t) \int_0^t \cos(\alpha\tau) \cos(\tau) d\tau \\
 &\quad - \sin(t) \int_0^t \cos(\alpha\tau) \sin(\tau) d\tau
 \end{aligned}$$

If $\alpha \neq 0$

$$\begin{aligned}
 \int_0^t \cos(\alpha\tau) \cos(\tau) d\tau &= \frac{1}{\alpha} \sin(\alpha\tau) \cos(\tau) \Big|_0^t \\
 &\quad + \frac{1}{\alpha} \int_0^t \sin(\alpha\tau) \sin(\tau) d\tau
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\alpha} \sin(\alpha t) \cos(t) - \frac{1}{\alpha^2} \cos(\alpha t) \sin(t) \Big|_0^t \\
 &\quad + \tau \frac{1}{\alpha^2} \int_0^t \cos(\alpha \tau) \cos(t) d\tau \\
 \Rightarrow & \int_0^t \cos(\alpha \tau) \cos(t) dt \left(1 - \frac{1}{\alpha^2} \right) \\
 &= \frac{1}{\alpha} \sin(\alpha t) \cos(t) - \frac{1}{\alpha^2} \cos(\alpha t) \sin(t)
 \end{aligned}$$

$$\begin{aligned}
 \alpha \neq 1 \\
 \Rightarrow & \int_0^t \cos(\alpha \tau) \cos(t) d\tau = \\
 & \frac{\alpha}{\alpha^2 - 1} \sin(\alpha t) \cos(t) - \frac{1}{\alpha^2 - 1} \cos(\alpha t) \sin(t)
 \end{aligned}$$

Similarly for the other integral. So

If $\alpha \neq 0, 1$

$$\begin{aligned}
 f_{\alpha \neq 0, 1} &= \\
 \cos(t) & \left[\frac{\alpha}{\alpha^2 - 1} \sin(\alpha t) \cos(t) - \frac{1}{\alpha^2 - 1} \cos(\alpha t) \sin(t) \right] \\
 - \sin^2(t) & \frac{\sin(\alpha t)}{\alpha^2 - 1} - \sin(t) \cos(t) \frac{\cos(\alpha t)}{\alpha^2 - 1} - \frac{1}{\alpha^2 - 1}
 \end{aligned}$$

If $\alpha = 0$:

$$f_\alpha * g = \sin(t)$$

If $\alpha = 1$:

$$f_\alpha * g = \frac{1}{2}(t\cos(t) + \sin(t))$$

b) $f_\alpha * g$ bounded for all $\alpha \neq 1$

c) Not periodic for $\alpha = 1$

Periods of $\sin(\alpha t)$: $\frac{2\pi}{\alpha} n, n \in \mathbb{Z}$.

Periods of $\sin(t)$: $2\pi m, m \in \mathbb{Z}$.

In order to have $f_\alpha * g$ periodic
we need

$$\frac{2\pi}{\alpha} n = 2\pi m \text{ for some } m, n \in \mathbb{Z}$$

$\Rightarrow \alpha = \frac{n}{m}$ a rational number.

4. a)

$$\log\left(\frac{1+s}{s-1}\right) \quad (\text{use Thm 3, p. 471})$$

$$b) \frac{2s(s^2-12)}{(s^2+4)^3} \quad (\text{use Thm 2, p. 469})$$

$$c) \text{ Write as } h(t) = t^3(u(t-1) - u(t-2)) \\ = ((t-1)+1)^3 u(t-1) + ((t-2)+2)^3 u(t-2)$$

$$H(s) = \frac{1}{s^4} \left[(s^3 + 3s^2 + 6s + 6)e^{-s} - (8s^3 + 12s^2 + 12s + 6)e^{-2s} \right]$$

$$5. \text{ Use rule } f(t) = -\frac{1}{t} \mathcal{L}^{-1}\{F'(s)\} \quad (\text{Thm 2, p. 469})$$

$$f(t) = \frac{e^{-2t} \sin(3t)}{t}$$

6. Take Laplace:

$$\mathcal{L}\{e(t)\} = 100t(1-u(t-1))$$

So:

$$sI(s) + 150I(s) + 5000 \quad \frac{I(s)}{s} = \frac{100}{s^2} - 100e^{-s}\left(\frac{1}{s} + \frac{1}{s^2}\right)$$

 $\Rightarrow \dots \Rightarrow$

$$I(s) = \frac{100}{s(s+50)(s+100)} + e^{-s} \frac{100(s+1)}{s(s+50)(s+100)}$$

use p-fractions

$$i(t) = \frac{1}{50} \left[1 - 2e^{-50t} + e^{-100t} \right] \\ - \frac{1}{50} u(t-1) \left[1 + 98e^{-50(t-1)} - 98e^{-100(t-1)} \right]$$

7. Laplace:

$$(s^2 X(s) - 2s - 2) + 2(sX(s) - 2) + X(s) = 1 - e^{-2s}$$

$$(s^2 + 2s + 1) X(s) = 7 + 2s - e^{-2s}$$

$$\Rightarrow X(s) = \frac{7 + 2s - e^{-2s}}{s^2 + 2s + 1}$$

$$\Rightarrow X(s) = \frac{7 + 2s}{(s+1)^2} - e^{-2s} \frac{1}{(s+1)^2}$$

$$\text{Write } \frac{7+2s}{(s+1)^2} = \frac{A}{s+1} + \frac{B}{(s+1)^2}$$

$$\text{Find } A = 2, B = 5$$

$$\Rightarrow X(s) = \frac{2}{s+1} + \frac{5}{(s+1)^2} - \frac{e^{-2s}}{(s+1)^2}$$

$$\Rightarrow x(t) = (2 + 5t)e^{-t} - u(t-2)(t-2)e^{-(t-2)}$$

$$8. \quad x'' + 6x' + 9x = f(t)$$

$$\Rightarrow X(s) (s^2 + 6s + 9) = F(s)$$

$$\Rightarrow X(s) = \frac{1}{s^2 + 6s + 9} F(s)$$

$$\Rightarrow x(t) = \int_0^t w(\tau) f(t-\tau) d\tau, \text{ where}$$

$$w(\tau) = \mathcal{L}^{-1} \left\{ \frac{1}{s^2 + 6s + 9} \right\} = e^{-3\tau} t \quad \text{is the weight function.}$$

$$\text{So } x(t) = \int_0^t e^{-3\tau} \tau f(t-\tau) d\tau$$

9. Only a) is periodic:

$$\sinh(2t) = \frac{1}{2}(e^{2t} - e^{-2t}) \text{ is not periodic}$$

$t \sin(2t)$ is cont. and increases unboundedly

$\arctan(t+\ell)$ is strictly increasing for all t .

$\sin(\pi t) + \sin(t)$:

↳ periods $2\pi, 4\pi, 6\pi$

↳ periods: $2, 4, 6, \dots$

} no common period.

$$(10. \text{ a}) \quad a_0 = \frac{1}{2} \int_0^2 t^2 dt = \frac{4}{3}$$

$$a_n = \frac{1}{2} \int_0^2 t^2 \cos \frac{n\pi t}{2} dt = \frac{8(-1)^n}{n^2 \pi^2}$$

$$b_n = \frac{1}{2} \int_0^2 t^2 \sin \frac{n\pi t}{2} dt$$

$$= - \frac{4((n^2\pi^2 - 2)\cos(n\pi) - 2n\pi \sin(n\pi)) + 2}{n^3 \pi^3}$$

$$= \begin{cases} -\frac{4}{\pi n} & n \text{ even} \\ \frac{4}{n\pi} - \frac{16}{n^3 \pi^3} & n \text{ odd} \end{cases}$$

Plug these into series

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \left(\frac{n\pi}{2} t \right) + b_n \sin \left(\frac{n\pi}{2} t \right) \right)$$

b) See textbook, ex. 2 p. 577.

$$c) \quad a_0 = \frac{2}{\pi}, \quad a_n = -\frac{1+(-1)^n}{\pi(n^2-1)}, \quad n \geq 1$$

$$a_1 = 0.$$


$$b_n = -\frac{\sin n\pi}{\pi(n^2-1)} \quad n \geq 1$$

$$b_1 = \frac{1}{2}$$

Plug in into

$$f = \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos(n\pi t) + b_n \sin(n\pi t))$$


None of those functions is even or odd.
 Term-by-term differentiation valid only
 for the last one.



F. cosine series:


$$f_1(t) = \cos(t)$$

F. sine series:




$$f = \sum_{n=2}^{\infty} \frac{2n(1+(-1)^n)}{(-1+n^2)\pi} \sin(nt)$$

↑
notice $b_1 = 0$.



$$\cos(t) = -\frac{8}{3\pi} + \sum_{n=1}^{\infty} \frac{12(-1)^n}{-9\pi + 4n^2\pi} \cos\left(\frac{2n}{3}t\right)$$



$$\cos(t) = \sum_{n=1}^{\infty} \frac{b_n}{-9\pi + 4n^2\pi} \sin\left(\frac{2n}{3}t\right)$$

12. Write $x(t) = \sum_{n=1}^{\infty} b_n \sin(nt)$ to satisfy condit condition

$$1 = \frac{4}{\pi} \sum_{n \text{ odd}} \frac{\sin(nt)}{n}$$

Find $\begin{cases} b_n = -\frac{4}{\pi n(n^2+4)} & n \text{ odd} \\ b_n = 0 & n \text{ even} \end{cases}$