


2. Set up an integral $\iiint_E f(x, y, z) dV$, where E is bounded above by the plane $z = 3 - x - y$, below by the xy plane, and also bounded by the planes $x = -1$, $x = 1$, $y = 0$ and $y = 1$ in the order $dx dz dy$.



① $x = 3 - z - y$

② $x = -1$

③ $x = 1$


④ $y = 0$

⑤ $y = 1$

Need sum of int's
Find projection on yz plane

①, ② $\Rightarrow z + y = 4$

①, ③ $\Rightarrow z + y = 2$



Over D_1 : $-1 \leq x \leq 3 - z - y$
and

$$D_1 = \{(y, z) : 2 - y \leq z \leq 4 - y, 0 \leq y \leq 1\}$$

Over D_2 : $-1 \leq x \leq 1$

and

$$D_2 = \{(y, z) : 0 \leq z \leq 2 - y \text{ and } 0 \leq y \leq 1\}$$

So

$$\iiint_E f dV = \int_0^1 \int_0^{2-y} \int_{-1}^1 f dx dz dy + \int_0^1 \int_{2-y}^{4-y} \int_{-1}^{3-z-y} f dx dz dy$$

3. Compute the volume of the bounded domain lying between the cones $z^2 = x^2 + y^2$ and $z^2 = 3(x^2 + y^2)$, and under the plane $z = 3$.



there is a domain here but it's not bounded.

Do it in spherical coords
 $z = \sqrt{x^2 + y^2} \Rightarrow$

$$\rho \cos \varphi = \sqrt{\rho^2 \sin^2 \varphi}$$

$$\Rightarrow \varphi = \frac{\pi}{4}$$

$$z = \sqrt{3} \sqrt{x^2 + y^2} \Rightarrow$$

$$\rho \cos \varphi = \sqrt{3} \sqrt{\rho^2 \sin^2 \varphi}$$

$$\Rightarrow \varphi = \frac{\pi}{6}$$

$$\text{So } \frac{\pi}{6} \leq \varphi \leq \frac{\pi}{4}$$

$$0 \leq \vartheta \leq 2\pi$$

$$z = 3 \Rightarrow \rho \cos \varphi = 3 \Rightarrow \rho = \frac{3}{\cos \varphi}$$

$$V = \int_0^{2\pi} \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \int_0^{\frac{3}{\cos \varphi}} \rho^2 \sin \varphi \, d\rho \, d\varphi \, d\vartheta$$

$$= \int_0^{2\pi} \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin \varphi \left[\frac{\rho^3}{3} \right]_0^{\frac{3}{\cos \varphi}} \, d\varphi \, d\vartheta = \int_0^{2\pi} \int_{\frac{\pi}{6}}^{\frac{\pi}{4}} \sin \varphi \frac{9}{\cos^3 \varphi} \, d\varphi \, d\vartheta$$

$$= \int_0^{2\pi} \frac{9}{2} \cos^{-2} \varphi \Big|_{\frac{\pi}{6}}^{\frac{\pi}{4}} \, d\vartheta = \int_0^{2\pi} \frac{9}{2} \left(\frac{1}{\left(\frac{\sqrt{2}}{2}\right)^2} - \frac{1}{\left(\frac{\sqrt{3}}{2}\right)^2} \right) \, d\vartheta$$

$$= 9\pi \left(2 - \frac{4}{3} \right) = 6\pi$$

Done in Cylindrical coords:

3. Compute the volume of the bounded domain lying between the cones $z^2 = x^2 + y^2$ and $z^2 = 3(x^2 + y^2)$, and under the plane $z = 3$.

Write down equations solving for z :

① $z = \sqrt{x^2 + y^2}$

② $z = \sqrt{3(x^2 + y^2)}$

③ $z = 3$

z appears 3 times, so if we want z to be the innermost variable we need a sum of 2 integrals

Find projection on xy plane:

①, ② $\Rightarrow \sqrt{x^2 + y^2} = \sqrt{3}\sqrt{x^2 + y^2} \Rightarrow x = y = 0$ (only a point)

①, ③ $\Rightarrow \sqrt{x^2 + y^2} = 3 \Rightarrow x^2 + y^2 = 9$ (a circle)

②, ③ $\Rightarrow \sqrt{3}\sqrt{x^2 + y^2} = 3 \Rightarrow x^2 + y^2 = 3$ (circle)

Over $D_1 = \{(x, y) : 3 \leq x^2 + y^2 \leq 9\}$

$= \{(r, \theta) : \sqrt{3} \leq r \leq 3, \theta \in [0, 2\pi]\}$

we have



$\sqrt{x^2 + y^2} \leq z \leq 3$ or $r \leq z \leq 3$

Over $D_2 = \{(x, y) : 0 \leq x^2 + y^2 \leq 3\}$

$= \{(r, \theta) : 0 \leq r \leq \sqrt{3}, 0 \leq \theta \leq 2\pi\}$

we have

$\sqrt{x^2 + y^2} \leq z \leq \sqrt{3(x^2 + y^2)}$ or $r \leq z \leq \sqrt{3}r$



So: $V = \int_0^{2\pi} \int_0^{\sqrt{3}} \int_r^{\sqrt{3}r} 1 \cdot r \, dz \, dr \, d\theta$

+ $\int_0^{2\pi} \int_{\sqrt{3}}^3 \int_r^3 1 \cdot r \, dz \, dr \, d\theta = 6\pi$

4. If a transformation T is written as $x = x(u, v)$ and $y = y(u, v)$ and given a point (u_0, v_0) on the uv plane, we can produce an affine transformation dT called the **differential** or **pushforward** of T at (u_0, v_0) that is the best affine approximation of T near (u_0, v_0) and is given by

$$x = \frac{\partial x}{\partial u}|_{(u,v)=(u_0,v_0)}(u - u_0) + \frac{\partial x}{\partial v}|_{(u,v)=(u_0,v_0)}(v - v_0) + x(u_0, v_0)$$

$$y = \frac{\partial y}{\partial u}|_{(u,v)=(u_0,v_0)}(u - u_0) + \frac{\partial y}{\partial v}|_{(u,v)=(u_0,v_0)}(v - v_0) + y(u_0, v_0).$$

For the transformation $T(u, v) = (\frac{u^2}{v}, u^2v)$ defined on $\{(u, v) : u > 0, v > 0\}$:

- (a) Find the transformation dT relative to the point $(1, 1)$.
 (b) Find and draw the image of the box $[1, 2] \times [1, 2]$ under T and dT .

a) $\frac{\partial x}{\partial u} = \frac{2u}{v} \quad \frac{\partial x}{\partial v} = -\frac{u^2}{v^2}$

$\frac{\partial y}{\partial u} = 2uv \quad \frac{\partial y}{\partial v} = u^2$

so $dT_{(1,1)} = (2(u-1) - (v-1) + 1, 2(u-1) + (v-1) + 1)$
 $= (2u - v, 2u + v - 2)$


or $x = 2u - v$
 $y = 2u + v - 2$

b) Solve for u, v in T :

$x = \frac{u^2}{v} \quad y = u^2v \quad \Rightarrow \quad xy = u^4 \quad \Rightarrow \quad u = \sqrt[4]{xy}$
 $\frac{y}{x} = v^2 \quad \Rightarrow \quad v = \sqrt{\frac{y}{x}}$

so $u = 1 \Rightarrow y = \frac{1}{x} \quad v = 1 \Rightarrow y = x$

$u = 2 \Rightarrow y = \frac{16}{x} \quad v = 2 \Rightarrow y = 4x$



Solve for u, v in $dT_{(1,1)}$

$x = 2u - v \quad y = 2u + v - 2 \quad \Rightarrow \quad 4u = x + y + 2$
 $2v = y - x + 2$
 $\Rightarrow \quad u = \frac{1}{4}(x + y) + \frac{1}{2}$
 $v = \frac{1}{2}(y - x) + 1$

so $u = 1 \Rightarrow x + y = 2$
 $u = 2 \Rightarrow x + y = 6$
 $v = 1 \Rightarrow y = x$
 $v = 2 \Rightarrow y = x + 2$

Drawn on the xy plane in red.

4. A fly flies in a room along the curve

$$c(t) = (2 \sin(t), \cos(t), 2t).$$

The temperature at the point (x, y, z) of the room is given by the function

$$T(x, y, z) = x^2 + 2e^z y,$$

in °C.

(a) Find the gradient of T .

(b) As the fly moves, it experiences a different temperature at each point. Find the instantaneous rate of change of the temperature that the fly experiences at time π seconds (include units).

$$a) \nabla T(x, y, z) = \langle 2x, 2e^z, 2e^z y \rangle$$

$$b) \text{ Need: } \frac{d}{dt}(T \circ c)(t)$$

Use Chain Rule: write $c(t) = (x(t), y(t), z(t))$

Then, using that $c(\pi) = (0, -1, 2\pi)$

$$\frac{d}{dt}(T \circ c)(t) = \frac{\partial T}{\partial x} \frac{dx}{dt} + \frac{\partial T}{\partial y} \frac{dy}{dt} + \frac{\partial T}{\partial z} \frac{dz}{dt}$$

$$= \nabla T(c(\pi)) \cdot \left\langle \frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right\rangle_{t=\pi}$$

$$= \langle 0, 2e^{2\pi}, 2e^{2\pi}(-1) \rangle \cdot \langle 2\cos t, -\sin t, 2 \rangle_{t=\pi}$$


$$= \langle 0, 2e^{2\pi}, -2e^{2\pi} \rangle \cdot \langle -2, 0, 2 \rangle$$

$$= -4e^{2\pi} \text{ } ^\circ\text{C/s}$$

1. Make a change of variables to evaluate the integral

$$\iint_R e^{\frac{y-x}{y+x}} dA,$$

where R is the trapezoidal region with vertices $(1,0)$, $(2,0)$, $(0,2)$ and $(0,1)$.




$$\text{set } \begin{cases} u = y+x \\ v = y-x \end{cases} \Rightarrow \begin{cases} x = \frac{u-v}{2} \\ y = \frac{u+v}{2} \end{cases}$$

$$y = 2 - x \Rightarrow y + x = 2 \Rightarrow u = 2$$

$$y = 1 - x \Rightarrow y + x = 1 \Rightarrow u = 1$$

$$x = 0 \Rightarrow u = v$$

$$y = 0 \Rightarrow u = -v$$



Jacobian:

$$\frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{vmatrix} = \frac{1}{4} + \frac{1}{4} = \frac{1}{2}$$

So

$$\iint_R e^{\frac{y-x}{y+x}} dA = \int_1^2 \int_{-u}^u \frac{1}{2} e^{\frac{v}{u}} dv du$$

$$= \int_1^2 \frac{1}{2} u e^{\frac{v}{u}} \Big|_{-u}^u du$$

$$= \int_1^2 \frac{1}{2} u (e - e^{-1}) du$$

$$= \frac{1}{2} \frac{u^2}{2} (e - e^{-1}) \Big|_1^2 = \frac{3}{4} (e - e^{-1})$$

6. *(Chain Rule, an interesting conceptual question) What's wrong with the following argument? Suppose we are given a function $w = f(x, y, z)$, where $z = g(x, y)$. Then, wishing to compute $\frac{\partial w}{\partial x}$, we draw a tree diagram and find that at each point (x, y) we have

$$\frac{\partial w}{\partial x} = \frac{\partial w}{\partial x} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial x} \implies \frac{\partial w}{\partial z} \frac{\partial z}{\partial x} = 0, \quad (1)$$

so at each point either $\frac{\partial w}{\partial z} = 0$ or $\frac{\partial z}{\partial x} = 0$, which can't be true: there were no assumptions on w or z !

Hint: it might help you to work out a specific example.

The expressions $\frac{\partial w}{\partial x}$ in the left and right hand side are not the same! It may help to think of the function w as $w = f(u, v, z)$, where $u = x, v = y, z = g(x, y)$. Then, the chain rule becomes

$$\frac{\partial w}{\partial x} = \frac{\partial w}{\partial u} \frac{du}{dx} + \frac{\partial w}{\partial z} \frac{dz}{dx}$$

$$\implies \frac{\partial w}{\partial x} = \frac{\partial w}{\partial u} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial x}$$

which avoids misconceptions. In other words, $\frac{\partial w}{\partial x}$ on the right hand side means that we're fixing y and z and taking derivative wrt x , whereas on the left hand side the dependency of z on x is taking effect.