

• Last time

- For any Lie group G , endowed $T_1 G$ with a Lie algebra structure (for $G = GL_n(\mathbb{K})$, recovered usual commutator $[A, B] = AB - BA$ on $gl_n(\mathbb{K})$) getting a functor $\{ \text{category of Lie gps} \} \rightarrow \{ \text{category of Lie algs} \}$
- Stated three fundamental theorems of Lie theory, implying an equivalence $\{ \text{category of connected simply-connected Lie gps} \} \simeq \{ \text{finite-dimensional Lie algebras (over } \mathbb{R} \text{ or } \mathbb{C}) \}$
- Henceforth, we shall mostly focus on the Lie algebras alone and also their representations.

Def 1: a) A representation of a Lie algebra $\mathfrak{g}$ over a field $\mathbb{K}$ is a vector space V over $\mathbb{K}$ together with a homomorphism of Lie algebras $\rho: \mathfrak{g} \rightarrow gl(V)$, i.e. ρ is $\mathbb{K}$ -linear and

$$\boxed{\rho([x, y]) = [\rho(x), \rho(y)] = \rho(x)\rho(y) - \rho(y)\rho(x) \quad \forall x, y \in \mathfrak{g}}$$

b) Given two representations (V, ρ_V) & (W, ρ_W) of $\mathfrak{g}$, a homomorphism from the former to the latter is a linear map commuting with $\mathfrak{g}$ -action, i.e.

$$d: V \rightarrow W \text{ linear s.t. } \boxed{d \circ \rho_V(x) = \rho_W(x) \circ d \quad \forall x \in \mathfrak{g}}$$

↑
equality in $\text{Hom}_{\mathbb{K}}(V, W)$

Terminology: • representations of $\mathfrak{g}$ are also called $\mathfrak{g}$ -modules
• homom. of repr-s are also called intertwining operators

Conventions: for $\mathbb{K} = \mathbb{R}$ it's also common to consider complex V (not just real)
this shall actually simplify the theory

As a Corollary of the 2nd Fund. Thm of Lie theory, get:

Proposition 1: Let G be a real/cpx Lie group and $\mathfrak{g} = \text{Lie}(G)$.

- Every fin. dim. rep. $\rho: G \rightarrow GL(V)$ defines a Lie algebra repr-n $\rho_*: \mathfrak{g} \rightarrow gl(V)$ and any morphism of G -representations is also a morphism of $\mathfrak{g}$ -modules
- If G is connected, simply-connected, then $\rho \mapsto \rho_*$ from a) gives an equivalence of categories b/w the corresponding categories of finite dimensional repr-s. In particular, every f. dim. $\mathfrak{g}$ -module can be uniquely exponentiated to a representation of G

Lecture #9

Example: Besides for the trivial repr-n, every Lie algebra $\mathfrak{g}$ admits an adjoint representation ρ_{ad} of $\mathfrak{g}$ given by $\text{ad}(x): y \mapsto [x, y]$.

Indeed, as we saw before, ad-representation $\Leftrightarrow$ Jacobi for $[\cdot, \cdot]$.

As always, it's important to recall basic procedures to construct $\mathfrak{g}$ -modules.

Def d: A subspace W of a $\mathfrak{g}$ -module (V, ρ_V) is called a subrepresentation (a.k.a. submodule) if it is $\mathfrak{g}$ -invariant, i.e. $\rho_V(x)W \subseteq W \quad \forall x \in \mathfrak{g}$.

• Similar definition applies to G -modules. In either case, the quotient space V/W carries a natural structure of $\mathfrak{g}$ -module (G -module), called the quotient representation.

• Given a $\mathfrak{g}$ -module V , the dual space V^* is endowed with a $\mathfrak{g}$ -module structure:

$$\boxed{\rho_{V^*}(x) := -\rho_V(x)^*} \quad \leftarrow \text{dual representation}$$

• Given two modules V, W of $\mathfrak{g}$, we can equip $V \oplus W$ with a $\mathfrak{g}$ -module structure:

$$\boxed{\rho_{V \oplus W}(x) := \rho_V(x) \oplus \rho_W(x)} \quad \leftarrow \text{direct sum of representations}$$

• ———, we can equip $V \otimes W$ with a $\mathfrak{g}$ -module structure:

$$\boxed{\rho_{V \otimes W}(x) := \rho_V(x) \otimes \text{id}_W + \text{id}_V \otimes \rho_W(x)} \quad \leftarrow \text{tensor product of representations}$$

Exercise 1: a) Verify the above $\mathfrak{g}$ -actions define $\mathfrak{g}$ -actions

b) For $\mathfrak{g} = \text{Lie}(G)$, derive these formulas from the corresponding constructions for G -modules

c) Identifying $V^* \otimes W \cong \text{Hom}(V, W)$ as vector spaces, construct a natural action of $\mathfrak{g}$ on $\text{Hom}(V, W)$, $\forall \mathfrak{g}$ -modules V, W .

d) Verify $\text{Hom}(V, W)^{\mathfrak{g}} = \text{Hom}_{\mathfrak{g}}(V, W)$ [see notation below for LHS].

In particular, above constructions endow $S^n V$, $\wedge^n V$, $V^{\otimes n} \otimes (V^*)^m$ with canonical $\mathfrak{g}$ -module structures.

Notation: For any $\mathfrak{g}$ -module V , the $\mathfrak{g}$ -invariants are $V^{\mathfrak{g}} := \{v \in V \mid \rho_V(x)v = 0 \quad \forall x \in \mathfrak{g}\}$

Lecture #9

13

Major problem in representation theory is to classify all representations of a Lie group or a Lie algebra. To this end, we start with building blocks:

Def 3: a) A representation $V \neq 0$ of G or $\mathfrak{g}$ is irreducible if it has no subrepresentation other than V or 0 .

b) A representation $V \neq 0$ of G or $\mathfrak{g}$ is indecomposable if it does not admit decomposition into direct sum of nonzero modules, i.e.

$$V \underset{\mathfrak{g}\text{-mod}}{\simeq} W_1 \oplus W_2 \Rightarrow W_1 \text{ or } W_2 \text{ are zero.}$$

Clearly, every finite dimensional repr-n is isomorphic to $\oplus$ indecomposable, but not necessarily into $\oplus$ irreducibles (give a counter example)

Def 4: A repr-n V is called completely reducible (a.k.a. semisimple) if it is $\simeq \oplus$ irreducibles, i.e. $V \simeq \oplus n_i V_i$ (n_i are called multiplicities)
 V_i -pairwise nonisomorphic irreducible modules

Thus, the major questions are:

(1) classify all irreducible repr-s of G or $\mathfrak{g}$

(2) for a given semisimple repr-n V of G or $\mathfrak{g}$, find all the multiplicities

(3) for which G or $\mathfrak{g}$, all fin. dim. repr-s are completely reducible?

The following result is almost obvious (easy exercise to do at home):

Lemma 1: Let V be a fin. dimensional representation of G or $\mathfrak{g}$, and let $A: V \rightarrow V$ be a homomorphism of $\mathfrak{g}$ -repr-s. Then

$V = \bigoplus_{\lambda} \underline{V(\lambda)}$ is a decomposition of representations
 $\uparrow$ generalized eigenspace of A with eigenvalue λ .

Exercise 2: Verify that $V, S^n V, \wedge^n V$ are irreducible representations of the group $GL(V)$ or a Lie algebra $\mathfrak{gl}(V)$. Deduce that $V \otimes V$ is completely reducible.

As for finite groups, we shall start with the basic Schur's lemma.

Lecture #9

4

Lemma 2 (Schur Lemma): Let V, W be irreducible fin. dim. $\mathbb{C}$ -representations of the group G or a Lie algebra $\mathfrak{g}$. Then:

- a) if $V \neq W$, we have $\text{Hom}_{G \text{ or } \mathfrak{g}}(V, W) = 0$, i.e. no nonzero homomorphisms
- b) if $V \cong W$, then $\text{Hom}_{G \text{ or } \mathfrak{g}}(V, V) = \mathbb{C} \cdot \text{Id}$

Let $A: V \rightarrow W$ be a nonzero homomorphism of repr-s.

Then $\text{Ker } A$ is a proper submodule of $V \xRightarrow{V \text{ irred.}} \text{Ker } A = 0$
 $\text{Im } A$ is a nonzero submodule of $W \xRightarrow{W \text{ irred.}} \text{Im } A = W$ } $\Rightarrow A: V \xrightarrow{\sim} W$.

This proves part a). For part b), let $\lambda \in \mathbb{C}$ be an eigenvalue of A , then $A - \lambda \cdot \text{Id}$ is still a homomorphism, but has nontrivial kernel $\Rightarrow A - \lambda \cdot \text{Id} = 0 \Rightarrow A = \lambda \text{Id}_V$. This proves part b).

Note: As part b) may fail over $\mathbb{R}$, we will mostly consider $\mathbb{C}$ -representations (give counterexample!)

As the classical corollaries of Schur's lemma, we get:

Corollary 1: The center of G or $\mathfrak{g}$ acts on any irreducible repr-n via scalars.
In particular, every irreducible repr-n of abelian G or $\mathfrak{g}$ is 1-dim.

Corollary 2: Let $\{V_i\}$ be irreducible pairwise non-isomorphic complex representations of G or $\mathfrak{g}$, and $V = \bigoplus n_i V_i$, $W = \bigoplus m_i V_i$. Then:

- a) there is a vector space isomorphism $\text{Hom}_{G \text{ or } \mathfrak{g}}(V, W) \cong \bigoplus_i \text{Mat}_{m_i \times n_i}(\mathbb{C})$
- b) for $W = V$, the isomorphism in a) is an algebra isomorphism

Remark: a) For Corollary 1, the center of $\mathfrak{g}$ is $\{x \in \mathfrak{g} \mid [x, y] = 0 \ \forall y \in \mathfrak{g}\}$.
In particular, $\rho(x)\rho(y) - \rho(y)\rho(x) = \rho([x, y]) = 0 \ \forall y \in \mathfrak{g} \Rightarrow$
 $\Rightarrow \rho(x)$ defines intertwiner $V \rightarrow V \xRightarrow{\text{Lemma 2b}} \rho(x) \text{ acts as } \lambda \cdot \text{Id}.$

b) For Corollary 2, it boils down to $\text{Hom}_{\mathfrak{g}}(n_i V_i, m_j V_j) = 0 \ \forall i \neq j$
 $\text{Hom}_{\mathfrak{g}}(n_i V_i, m_i V_i) \cong \text{Mat}_{m_i \times n_i}$

Lecture #9

We conclude today's lecture with the discussion of $\mathfrak{sl}_2(\mathbb{C})$ repr. theory.
Down-to-earth, $\mathfrak{sl}_2(\mathbb{C})$ has a basis

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

with a commutator

$$[e, f] = h, \quad [h, e] = 2e, \quad [h, f] = -2f.$$

[Exercise 3] Prove these are defining relations for $\mathfrak{sl}_2$.

Note that $\text{Mat}_{2 \times 2}(\mathbb{C}) \curvearrowright \mathbb{C}^2$ actually gives rise to the action (check!)

$$\mathfrak{sl}_2 \curvearrowright \mathbb{C}[x, y] =: V \text{ via } e \mapsto x \partial_y, \quad h \mapsto x \partial_x - y \partial_y, \quad f = y \partial_x$$

each operator preserves degrees and so $V \cong \bigoplus_{n \geq 0} V_n$ is an $\mathfrak{sl}_2$ -decomposition, where $V_n = \{\text{degree } n \text{ polynomials in } x, y\}$, $\dim V_n = n+1$.

Down-to-earth, V_n has a basis $\{x^k y^{n-k} \mid 0 \leq k \leq n\}$ and action is $\stackrel{=}{=} \mathcal{V}_{k, n-k}$

$$\begin{cases} e(\mathcal{V}_{k, n-k}) = (n-k) \mathcal{V}_{k+1, n-(k+1)} \\ h(\mathcal{V}_{k, n-k}) = (2k-n) \mathcal{V}_{k, n-k} \\ f(\mathcal{V}_{k, n-k}) = k \mathcal{V}_{k-1, n-(k-1)} \end{cases}$$

The following is the key result on $\mathfrak{sl}_2$ -theory (which will be later used to treat most general simple Lie algebras):

Theorem 1: a) V_n are irreducible $\mathfrak{sl}_2$ -modules

b) If $V \neq 0$ is a f.in. dim. $\mathfrak{sl}_2$ -module, then e & f are nilpotent on V .
Therefore, $U := \text{Ker}(e) \neq 0$. Finally, h preserves U , acting diagonally with $\mathbb{Z}_{\geq 0}$ coefficients

c) Any irred. f.in. dim. $\mathfrak{sl}_2$ -module is $\cong V_n$ for some n

d) Any f.in. dim. $\mathfrak{sl}_2$ -module is completely reducible