

MA 174: Multivariable Calculus

EXAM II (practice)

NAME _____ INSTRUCTOR _____

NO CALCULATORS, BOOKS, OR PAPERS ARE ALLOWED. Use the back of the test pages for scrap paper.

Points awarded

- | | |
|------------------|------------------|
| 1. (5 pts) _____ | 9. (5 pts) _____ |
| 2. (5 pts) _____ | 9. (5 pts) _____ |
| 3. (5 pts) _____ | 9. (5 pts) _____ |
| 4. (5 pts) _____ | 9. (5 pts) _____ |
| 5. (5 pts) _____ | 9. (5 pts) _____ |
| 6. (5 pts) _____ | 9. (5 pts) _____ |

Total Points: _____

1. Suppose $z = f(x, y)$, where $x = e^t$ and $y = t^2 + 3t + 2$. Given that $\frac{\partial z}{\partial x} = 2xy^2 - y$ and $\frac{\partial z}{\partial y} = 2x^2y - x$, find $\frac{dz}{dt}$ when $t = 0$.

A. 3
B. 6
C.
D. 9
E. -1

2. Find the directional derivative of the function $f(x, y, z) = x^2y^2z^6$ at the point $(1, 1, 1)$ in the direction of the vector $\langle 2, 1, -2 \rangle$.

A. -6
B.
C. 0
D. 2
E. 6

3. Find the direction in which the function $z = x^2 + 3xy - \frac{1}{2}y^2$ is increasing most rapidly at $(-1, -1)$.

A. $3i$
B. $5\vec{i} + 2\vec{j} - \vec{k}$
C.
D. $2\vec{i} - 5\vec{j}$
E. $\sqrt{29}$

4. If $xz^3 - xyz = 4$, find $\frac{\partial z}{\partial x}$.

A. $\frac{\partial z}{\partial x} = \frac{xz}{z^3 - y^2}$

B. $\frac{\partial z}{\partial x} = \frac{3xz^2 - xy}{z^3 - yz}$

C. $\frac{\partial z}{\partial x} = 2x + xy$

D. $\frac{\partial z}{\partial x} = \frac{yz - z^3}{3xz^2 - xy}$

E. $\frac{\partial z}{\partial x} = z^3 - yz$

5. The directional derivative of $f(x, y) = x^3e^{-2y}$ in the direction of greatest increase of f at the point $(1, 0)$ is

A. 6

B. 5

C. $\sqrt{5}$

D. 13

E. $\sqrt{13}$

6. By using a linear approximation of $f(x, y) = \sqrt{x^2 + y}$ at $(4, 9)$, compute the approximate value of $f(5, 8)$.

A. 5.2

B. 5.3

C. 5.5

D. $\sqrt{5.7}$

E. 5.9

7. The max and min values of $f(x, y, z) = xyz$ on the surface $2x^2 + 2y^2 + z^2 = 2$ are

- A. $\pm \frac{\sqrt{2}}{9}$
- B. $\pm \frac{\sqrt{3}}{9}$
- C. $\pm \frac{\sqrt{6}}{9}$
- D. $\pm \frac{2\sqrt{2}}{9}$
- E. $\pm \frac{2\sqrt{3}}{9}$

8. Find the maximum value of $x^2 + y^2$ subject to the constraint $x^2 - 2x + y^2 - 4y = 0$.

- A. 0
- B. 2
- C. 4
- D. 16
- E. 20

9. If we use the method of Lagrange multipliers to find the maximum of $f(x, y) = 2x^2 - y^2 - y$ subject to the constraint $x^2 + y^2 = 1$, the Lagrange multipliers λ that we find are:

- A. $\lambda = 2$
- B. $\lambda = 0$
- C. $\lambda = -1$
- D. $\lambda = 2$ and $\lambda = -1$
- E. $\lambda = 0$ and $\lambda = -1$

10. For the function $f(x, y) = x^3 + 2y^2 + xy - 2x + 5y$, the point $(-1, -1)$ yields

A. a local minimum

B. a local maximum

C. a saddle point

D. $\nabla f(-1, -1) \neq 0$

E. The Second Derivative Test gives no information at $(-1, -1)$

11. Use the method of reversing the order of integration to compute

$$\int_0^1 \int_{2x}^2 e^{y^2} dy dx.$$

A. $\frac{1}{4}(e^4 - 1)$

B. $\frac{1}{2}(e^2 - 1)$

C. $\frac{1}{6}(e^3 - 1)$

D. $\frac{1}{2}(e^2 - e)$

E. $\frac{1}{4}(e^2 - e)$

12. A flat plate of constant density occupies the region in the xy -plane bounded by the curves $x = 0$ and $x = \sqrt{1 - y^2}$. If $(\bar{x}, \bar{y})$ is the center of mass, then $\bar{x}$ equals

A. $\frac{2}{3\pi}$

B. $\frac{1}{2}$

C. $\frac{2}{\pi}$

D. $\frac{3}{2\pi}$

E. $\frac{4}{3\pi}$

13. Find the volume of the solid whose base is the region in the xy -plane that is bounded by the parabola $y = 4 - x^2$ and the line $y = 3x$, while the top of the solid is bounded by the plane $z = x + 4$.

A. $\boxed{\frac{625}{12}}$

B. $\frac{625}{11}$

C. $\frac{542}{13}$

D. $\sqrt{15} \pi$

E. $\frac{\sqrt{8} \pi}{3}$

14. Which of the following integrals equals the volume of the solid bounded by $x = 0$, $y = 0$, $z = 0$ and $2x + y + z = 4$.

A. $\int_0^4 \int_0^4 \int_0^2 1 dx dy dz$

B. $\int_0^2 \int_0^{4-2x} \int_0^{4-y} 1 dz dy dx$

C. $\int_0^4 \int_0^{2x} \int_0^{4-y} 1 dz dy dx$

D. $\boxed{\int_0^2 \int_0^{4-2x} \int_0^{4-2x-y} 1 dz dy dx}$

E. $\int_0^2 \int_0^1 \int_0^1 1 dz dx dy$

15. Evaluate $\iint_R (x + 2y) dA$ where R is the region of the plane bounded by $x + y = 2$, $x = y$, $y = 0$.

A. $1/3$

B. $\boxed{5/3}$

C. $7/3$

D. $11/3$

E. $14/3$

16. Let

$$S: x = u - v, \ y = uv, \ z = u + v^2$$

If $(0, b, 5)$ is a point on the tangent plane to S at $(0, 1, 2)$ on S , then $b =$

- A. $\boxed{3}$
- B. 1
- C. -2
- D. 0
- E. 2

17. Find $\left(\frac{\partial w}{\partial y}\right)_x$ at $(w, x, y, z) = (4, 2, 1, -1)$ if

$$w = x^2y^2 + yz - z^3, \quad x^2 + y^2 + z^2 = 6$$

- A. -1
- B. 1
- C. 3
- D. $\boxed{5}$
- E. 7

18. Consider the function $f(x, y) = 2x^2 - 3xy + y^2$. Find two unit vectors such that the directional derivative of f at the point $(1, 1)$ in these two directions is 1.

Answer: $(1, 0)$ and $(0, -1)$

19. Find cubic approximation of $f(x, y) = \frac{1}{1 - x - y + xy}$ near the origin.

Answer: $1 + x + y + x^2 + xy + y^2 + x^3 + x^2y + xy^2 + y^3$

20. Find a equation for the tangent plane of

$$\cos(\pi x) - x^2y + e^{xz} + yz = 4 \quad \text{at} \quad (0, 1, 2)$$

Answer: $2x + 2y + z - 4 = 0$

21. Find absolute maximum and minimum values of

$$f(x, y) = x^2 + 2y^2 - x$$

on the disc $x^2 + y^2 \leq 1$.

Answer: $\max = \frac{9}{4}$, $\min = -\frac{1}{4}$