

# MA 261 QUIZ 6

## FEBRUARY 27, 2019

If you do not know how to do any one of these problems, circle “(E) I don’t know” as your answer choice. You will receive **two points** for doing that. **Each problem** is worth **five points**. You get **two points** for writing your **full name** and **three points** for writing your **section number**.

**Problem 6.1.** Find the maximum and minimum of  $f(x, y, z) = xyz$  subject to the constraint  $x + y + z = 1$  for  $x, y, z \geq 0$ .

(Hint: Use Lagrange multipliers.)

- (A) maximum  $1/27$ , minimum  $0$
- (B) maximum  $0$ , minimum  $0$
- (C) maximum  $1/27$ , minimum  $-1/27$
- (D) maximum  $1/3$ , minimum  $0$
- (E) I don’t know how to do this

*Solution.* Let  $g(x, y, z) = x + y + z$ . By the method of Lagrange multipliers,

$$\nabla f(x, y, z) = \langle yz, xz, xy \rangle = \langle \lambda, \lambda, \lambda \rangle = \lambda \nabla g(x, y, z).$$

Therefore, we have the following system of equations

$$\begin{aligned}yz &= \lambda, \\xz &= \lambda, \\xy &= \lambda.\end{aligned}$$

From here, it is not difficult to see that we must have the following sequence of equalities:  $x = y = z$ .

Feeding this back into the constraint,  $g(x, y, z) = 3x = 1$  we see that  $x = 1/3$  so  $y = z = 1/3$  and the maximum value must be  $1/27$ , subject to these constraints.

The minimum is trickier to come up with. Note that  $x, y, z \geq 0$  so, in particular, they can equal 0. Let  $x = 0$ . Then, since  $x + y + z = 1$ ,  $y + z = 1$  so let  $y = z = 1/2$  (or whatever else you can think of that adds up to 1). At any rate, the product  $xyz = 0$  for any choice of  $y$  and  $z$  since  $x = 0$ . Therefore, the minimum of  $f$  subject to the constraint  $g$  is 0.

**Answer:** (A) ◇

**Problem 6.2.** Find the absolute maximum of  $f(x, y) = 2x^2 - y^2 + 6y$  on  $x^2 + y^2 \leq 16$ .

- (A) 8

- (B) 9
- (C) 35
- (D) 40
- (E) I don't know how to do this

*Solution.* To find the absolute maximum of this function, subject to the constraints, we systematically apply the conclusion to the Extreme Value Theorem (EVT). Recall that the EVT says that a continuous function takes its extremal values at critical points within its domain or on the boundary of said domain.

In this case,  $f(x, y) = 2x^2 - y^2 + 6y$  has  $x^2 + y^2 \leq 16$  as its domain so we need to check the points at which  $\nabla f(x, y) = \langle 0, 0 \rangle$  or at points in the boundary, i.e.,  $(x, y)$  with  $x^2 + y^2 = 16$ .

Let's parametrize in terms of  $y$  to avoid leaving a square root dangling. Then

$$\tilde{f}(y) = f\left(\pm\sqrt{16 - y^2}, y\right) = 2(16 - y^2) - y^2 + 6y = 32 - 3y^2 + 6y.$$

Thus,

$$\tilde{f}'(y) = -6y + 6 = 0$$

for  $y = 1$ . It is easy to check, by the second derivative test (or simply observing that  $\tilde{f}'(y)$  is positive to the left of 1 and negative to the right) It follows that  $\tilde{f}(1) = f(\pm\sqrt{15}, 1) = 35$  is a max. Moreover, it is the largest value we have encountered, so it must be the global max in this domain.

**Answer:** (C). ◇

**Problem 6.3.** Find the value of the iterated integral

$$\iint_R 2 - x \, dA, \quad R = \{(x, y) : 0 \leq x \leq 2, 0 \leq y \leq 3\}.$$

- (A) 6
- (B) 8
- (C) 10
- (D) 12
- (E) I don't know how to do this

*Solution.* A straightforward calculation yields the answer:

$$\begin{aligned}\iint_R 2 - x \, dA &= \int_0^2 \int_0^3 2 - x \, dydx \\ &= 3 \int_0^2 2 - x \, dx \\ &= 3(2x - x^2/2)\Big|_0^2 \\ &= 3(4 - 2^2/2) \\ &= 6.\end{aligned}$$

**Answer:** (A)

◇