

# Axioms of Probability

Samy Tindel

Purdue University

Introduction to Probability Theory - MA 519

Mostly taken from *A first course in probability*  
by S. Ross

# Outline

- 1 Introduction
- 2 Sample space and events
- 3 Axioms of probability
- 4 Some simple propositions
- 5 Sample spaces having equally likely outcomes
- 6 Probability as a continuous set function

# Outline

- 1 Introduction
- 2 Sample space and events
- 3 Axioms of probability
- 4 Some simple propositions
- 5 Sample spaces having equally likely outcomes
- 6 Probability as a continuous set function

# Global objective

Aim: Introduce

- Sample space
- Events of an experiment
- Probability of an event
- Show how probabilities can be computed in certain situations

# Outline

- 1 Introduction
- 2 Sample space and events**
- 3 Axioms of probability
- 4 Some simple propositions
- 5 Sample spaces having equally likely outcomes
- 6 Probability as a continuous set function

# Sample space

**Situation:** We run an experiment for which

- Specific outcome is unknown
- Set  $S$  of possible outcomes is known

**Terminology:**

In the context above  $S$  is called **sample space**

# Examples of sample spaces

Tossing two dice: We have

$$\begin{aligned} S &= \{1, 2, 3, 4, 5, 6\}^2 \\ &= \{(i, j); i, j = 1, 2, 3, 4, 5, 6\} \end{aligned}$$

Lifetime of a transistor: We have

$$S = \mathbb{R}_+ = \{x \in \mathbb{R}; 0 \leq x < \infty\}$$

# Events

## Definition 1.

Consider

- Experiment with sample space  $S$
- A subset  $E$  of  $S$

Then

$E$  is called event



# Example of event (1)

Tossing two dice: We have

$$S = \{1, 2, 3, 4, 5, 6\}^2$$

Event: We define

$$E = (\text{Sum of dice is equal to 7})$$

## Example of event (2)

Description of  $E$  as a subset:

$$E = \{(1, 6); (2, 5); (3, 4); (4, 3); (5, 2); (6, 1)\} \subset S$$

## Second example of event (1)

Lifetime of a transistor: We have

$$S = \mathbb{R}_+ = \{x \in \mathbb{R}; 0 \leq x < \infty\}$$

Event: We define

$$E = (\text{Transistor does not last longer than 5 hours})$$

## Second example of event (2)

Description of  $E$  as a subset:

$$E = [0, 5] \subset S$$

# Operations on events

Complement:  $A^c$  is the set of elements of  $E$  not in  $A$

Two dice example:

$$E^c = \text{"Sum of two dice different from 7"}$$

Union, Intersection: For the two dice example, if

$$B = \text{"Sum of two dice is divisible by 3"}$$

$$C = \text{"Sum of two dice is divisible by 4"}$$


Then

$$B \cup C = \text{"Sum of two dice is divisible by 3 or 4"}$$


$$B \cap C = BC = \text{"Sum of two dice is divisible by 3 and 4"}$$

# Illustration (1)

Union and intersection:




(a) Shaded region:  $E \cup F$ .



(b) Shaded region:  $EF$ .

## Illustration (2)

Complement:



(c) Shaded region:  $E^c$ .

# Illustration (3)

Subset:




Figure:  $E \subset F$



# Laws for elementary operations

Commutative law:

$$E \cup F = F \cup E, \quad EF = FE$$

Associative law:


$$(E \cup F) \cup G = E \cup (F \cup G), \quad E(FG) = (EF)G$$

Distributive laws:


$$\begin{aligned}(E \cup F)G &= EG \cup EF \\ (EF) \cup G &= (E \cup G)(F \cup G)\end{aligned}$$

# Illustration


## Distributive law:



(a) Shaded region:  $EG$ .



(b) Shaded region:  $FG$ .



(c) Shaded region:  $(E \cup F)G$ .

Figure:  $(E \cup F)G = EG \cup FG$

# De Morgan's laws

## Proposition 2.

Let

- $S$  sample space
- $E_1, \dots, E_n$  events

Then

$$\left( \bigcup_{i=1}^n E_i \right)^c = \bigcap_{i=1}^n E_i^c$$

$$\left( \bigcap_{i=1}^n E_i \right)^c = \bigcup_{i=1}^n E_i^c$$

# Proof (1)

Proof of  $(\cup_{i=1}^n E_i)^c \subset \cap_{i=1}^n E_i^c$ :

Assume  $x \in (\cup_{i=1}^n E_i)^c$  Then

$$\begin{aligned}x \notin \cup_{i=1}^n E_i &\implies \text{for all } i \leq n, x \notin E_i \\ &\implies \text{for all } i \leq n, x \in E_i^c \\ &\implies x \in \cap_{i=1}^n E_i^c\end{aligned}$$

## Proof (2)

Proof of  $\bigcap_{i=1}^n E_i^c \subset (\bigcup_{i=1}^n E_i)^c$ :

Assume  $x \in \bigcap_{i=1}^n E_i^c$  Then

$$\begin{aligned} \text{for all } i \leq n, x \in E_i^c &\implies \text{for all } i \leq n, x \notin E_i \\ &\implies x \notin \bigcup_{i=1}^n E_i \\ &\implies x \in (\bigcup_{i=1}^n E_i)^c \end{aligned}$$

# Outline

- 1 Introduction
- 2 Sample space and events
- 3 Axioms of probability**
- 4 Some simple propositions
- 5 Sample spaces having equally likely outcomes
- 6 Probability as a continuous set function

# Definition of probability

## Definition 3.

A probability is an application which assigns a number (chances to occur) to any event  $E$ . It must satisfy 3 axioms

①

$$0 \leq \mathbf{P}(S) \leq 1$$

②

$$\mathbf{P}(S) = 1$$

③ If  $E_i E_j = \emptyset$  for  $i, j \geq 1$  such that  $i \neq j$ , then

$$\mathbf{P}\left(\bigcup_{i=1}^{\infty} E_i\right) = \sum_{i=1}^{\infty} \mathbf{P}(E_i)$$

# Easy consequence of the axioms

## Proposition 4.

Let  $\mathbf{P}$  be a probability on  $S$ . Then

①

$$\mathbf{P}(\emptyset) = 0$$

②

For  $n \geq 1$ ,  
if  $E_i E_j = \emptyset$  for  $1 \leq i, j \leq n$  such that  $i \neq j$  then

$$\mathbf{P}\left(\bigcup_{i=1}^n E_i\right) = \sum_{i=1}^n \mathbf{P}(E_i)$$



# Example: dice tossing

**Experiment:** tossing one dice

**Model:**  $S = \{1, \dots, 6\}$  and

$$\mathbf{P}(\{s\}) = \frac{1}{6}, \quad \text{for all } s \in S$$

**Probability of an event:** If  $E =$  "even number obtained", then

$$\begin{aligned} \mathbf{P}(E) &= \mathbf{P}(\{2, 4, 6\}) = \mathbf{P}(\{2\} \cup \{4\} \cup \{6\}) \\ &= \mathbf{P}(\{2\}) + \mathbf{P}(\{4\}) + \mathbf{P}(\{6\}) = \frac{3}{6} = \frac{1}{2} \end{aligned}$$

# Outline

- 1 Introduction
- 2 Sample space and events
- 3 Axioms of probability
- 4 Some simple propositions**
- 5 Sample spaces having equally likely outcomes
- 6 Probability as a continuous set function

# Probability of a complement

## Proposition 5.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- $E$  an event

Then

$$\mathbf{P}(E^c) = 1 - \mathbf{P}(E)$$

# Proof

Use Axioms 2 and 3:

$$1 = \mathbf{P}(S) = \mathbf{P}(E \cup E^c) = \mathbf{P}(E) + \mathbf{P}(E^c)$$

# Probability of a subset

## Proposition 6.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- $E, F$  two events, such that  $E \subset F$

Then

$$\mathbf{P}(E) \leq \mathbf{P}(F)$$

# Proof

Decomposition of  $F$ : Write

$$F = E \cup E^c F$$

Use Axioms 1 and 3: Since  $E$  and  $E^c F$  are disjoint,

$$\mathbf{P}(F) = \mathbf{P}(E \cup E^c F) = \mathbf{P}(E) + \mathbf{P}(E^c F) \geq \mathbf{P}(E)$$

# Probability of a non disjoint union

## Proposition 7.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- $E, F$  two events


Then

$$\mathbf{P}(E \cup F) = \mathbf{P}(E) + \mathbf{P}(F) - \mathbf{P}(E \cap F)$$

# Proof

Decomposition of  $E \cup F$ :

$$E \cup F = I \cup II \cup III$$





## Proof (2)

Decomposition for probabilities: We have

$$\mathbf{P}(E \cup F) = \mathbf{P}(I) + \mathbf{P}(II) + \mathbf{P}(III)$$

$$\mathbf{P}(E) = \mathbf{P}(I) + \mathbf{P}(II)$$

$$\mathbf{P}(F) = \mathbf{P}(II) + \mathbf{P}(III)$$

Conclusion: Since  $II = E \cap F$ , we get

$$\mathbf{P}(E \cup F) = \mathbf{P}(E) + \mathbf{P}(F) - \mathbf{P}(II) = \mathbf{P}(E) + \mathbf{P}(F) - \mathbf{P}(E \cap F)$$

# Application of Propositions 5 and 7

Experiment: dice tossing

$\hookrightarrow S = \{1, \dots, 6\}$  and  $\mathbf{P}(\{s\}) = \frac{1}{6}$  for all  $s \in S$

Events:

We consider  $A =$  "even outcome" and  $B =$  "outcome multiple of 3"

$\Rightarrow A = \{2, 4, 6\}$  and  $B = \{3, 6\}$

$\Rightarrow \mathbf{P}(A) = 1/2$  and  $\mathbf{P}(B) = 1/3$

Applying Propositions 5 and 7:

$\mathbf{P}(A^c) = 1 - \mathbf{P}(A) = 1/2$

$\mathbf{P}(A \cup B) = \mathbf{P}(A) + \mathbf{P}(B) - \mathbf{P}(A \cap B) = 1/2 + 1/3 - \mathbf{P}(\{6\}) = 2/3$

Verification:

$A^c = \{1, 3, 5\} \Rightarrow \mathbf{P}(A^c) = 1/2$

$A \cup B = \{2, 3, 4, 6\} \Rightarrow \mathbf{P}(A \cup B) = 4/6 = 2/3$

# Inclusion-exclusion identity

## Proposition 8.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- $n$  events  $E_1, \dots, E_n$

Then

$$\mathbf{P} \left( \bigcup_{i=1}^n E_i \right) = \sum_{r=1}^n (-1)^{r+1} \sum_{1 \leq i_1 < \dots < i_r \leq n} \mathbf{P} (E_{i_1} \cdots E_{i_r})$$

# Proof for $n = 3$

Apply Proposition 7:

$$\begin{aligned}\mathbf{P}(E_1 \cup E_2 \cup E_3) &= \mathbf{P}(E_1 \cup E_2) + \mathbf{P}(E_3) - \mathbf{P}((E_1 \cup E_2)E_3) \\ &= \mathbf{P}(E_1 \cup E_2) + \mathbf{P}(E_3) - \mathbf{P}(E_1E_3 \cup E_2E_3)\end{aligned}$$

Apply Proposition 7 to  $E_1 \cup E_2$  and  $E_1E_3 \cup E_2E_3$ :

$$\mathbf{P}(E_1 \cup E_2 \cup E_3) = \sum_{1 \leq i_1 \leq 3} \mathbf{P}(E_{i_1}) - \sum_{1 \leq i_1 < i_2 \leq 3} \mathbf{P}(E_{i_1}E_{i_2}) + \mathbf{P}(E_1E_2E_3)$$

Case of general  $n$ : By induction

# Bounds for $\mathbf{P}(\cup_{i=1}^n E_i)$

## Proposition 9.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- $n$  events  $E_1, \dots, E_n$

Then

$$\mathbf{P}\left(\bigcup_{i=1}^n E_i\right) \leq \sum_{1 \leq i \leq n} \mathbf{P}(E_i)$$

$$\mathbf{P}\left(\bigcup_{i=1}^n E_i\right) \geq \sum_{1 \leq i \leq n} \mathbf{P}(E_i) - \sum_{1 \leq i_1 < i_2 \leq n} \mathbf{P}(E_{i_1} E_{i_2})$$

# Bounds for $\mathbf{P}(\cup_{i=1}^n E_i)$ – Ctd

## Proposition 10.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- $n$  events  $E_1, \dots, E_n$

Then

$$\begin{aligned} & \mathbf{P}\left(\bigcup_{i=1}^n E_i\right) \\ & \leq \sum_{1 \leq i \leq n} \mathbf{P}(E_i) - \sum_{1 \leq i_1 < i_2 \leq n} \mathbf{P}(E_{i_1} E_{i_2}) + \sum_{1 \leq i_1 < i_2 < i_3 \leq n} \mathbf{P}(E_{i_1} E_{i_2} E_{i_3}) \end{aligned}$$

# Proof

Notation: Set

$$B_i = E_1^c \cdots E_{i-1}^c$$

Identity:

$$\mathbf{P}(\cup_{i=1}^n E_i) = \mathbf{P}(E_1) + \sum_{i=2}^n \mathbf{P}(B_i E_i)$$

Second identity: Since  $B_i = (\cup_{j<i} E_j)^c$ ,

$$\mathbf{P}(B_i E_i) = \mathbf{P}(E_i) - \mathbf{P}(\cup_{j<i} E_j E_i)$$

Partial conclusion:

$$\mathbf{P}(\cup_{i=1}^n E_i) = \sum_{1 \leq i \leq n} \mathbf{P}(E_i) - \sum_{1 \leq i \leq n} \mathbf{P}(\cup_{j<i} E_j E_i)$$

## Proof (2)

Recall:

$$\mathbf{P}(\cup_{i=1}^n E_i) = \sum_{1 \leq i \leq n} \mathbf{P}(E_i) - \sum_{1 \leq i < j \leq n} \mathbf{P}(E_j E_i) \quad (1)$$

Direct consequence of (1):

$$\mathbf{P}(\cup_{i=1}^n E_i) \leq \sum_{1 \leq i \leq n} \mathbf{P}(E_i) \quad (2)$$

Application of (2) to  $\mathbf{P}(\cup_{j < i} E_j E_i)$ :

$$\mathbf{P}(\cup_{j < i} E_j E_i) \leq \sum_{j < i} \mathbf{P}(E_j E_i)$$

Plugging into (1) we get

$$\mathbf{P}(\cup_{i=1}^n E_i) \geq \sum_{1 \leq i \leq n} \mathbf{P}(E_i) - \sum_{j < i} \mathbf{P}(E_j E_i)$$



# Outline

- 1 Introduction
- 2 Sample space and events
- 3 Axioms of probability
- 4 Some simple propositions
- 5 Sample spaces having equally likely outcomes**
- 6 Probability as a continuous set function

# Model

**Hypothesis:** We assume

- $S = \{s_1, \dots, s_N\}$  **finite**.
- $\mathbf{P}(\{s_i\}) = \frac{1}{N}$  for all  $1 \leq i \leq N$

**Alert:**

This is an important but **very particular** case of probability space

**Example:** tossing 4 dice

$\hookrightarrow S = \{1, \dots, 6\}^4$  and

$$\begin{aligned}\mathbf{P}(\{(1, 1, 1, 1)\}) &= \mathbf{P}(\{(1, 1, 1, 2)\}) = \dots = \mathbf{P}(\{(6, 6, 6, 6)\}) \\ &= \frac{1}{6^4} = \frac{1}{1296}\end{aligned}$$

# Computing probabilities

## Proposition 11.

**Hypothesis:** We assume

- $S = \{s_1, \dots, s_N\}$  finite.
- $\mathbf{P}(\{s_i\}) = \frac{1}{N}$  for all  $1 \leq i \leq N$

In this situation, let  $E \subset S$  be an event. Then

$$\mathbf{P}(E) = \frac{\text{Card}(E)}{N} = \frac{|E|}{N} = \frac{\# \text{ outcomes in } E}{\# \text{ outcomes in } S}$$

# Example: tossing one dice

**Model:** tossing one dice, that is

$$S = \{1, \dots, 6\}, \quad \mathbf{P}(\{s_i\}) = \frac{1}{6}$$

**Computing a simple probability:** Let  $E = \text{"even outcome"}$ . Then

$$\mathbf{P}(E) = \frac{|E|}{N} = \frac{3}{6} = \frac{1}{2}$$

**Main problem:** compute  $|E|$  in more complex situations

↔ Counting

# Example: drawing balls (1)

**Situation:** We have

- A bowl with 6 White and 5 Black balls
- We draw 3 balls

**Problem:** Compute

$$\mathbf{P}(E), \quad \text{with } E = \text{"Draw 1 W and 2 B"}$$

## Example: drawing balls (2)

Model 1: We take

- $S = \{\text{Ordered triples of balls, tagged from 1 to 11}\}$
- $\mathbf{P} = \text{Uniform probability on } S$

Computing  $|S|$ : We have

$$|S| = 11 \cdot 10 \cdot 9 = 990$$

Decomposition of  $E$ : We have

$$E = WBB \cup BWB \cup BBW$$

## Example: drawing balls (3)

Counting  $E$ :

$$|E| = |WBB| + |BWB| + |BBW| = 3 \times (6 \times 5 \times 4) = 360$$

Probability of  $E$ : We get

$$P(E) = \frac{|E|}{|S|} = \frac{360}{990} = \frac{4}{11} = 36.4\%$$

## Example: drawing balls (4)

Model 2: We take

- $S = \{\text{Non ordered triples of balls, tagged from 1 to 11}\}$
- $\mathbf{P} = \text{Uniform probability on } S$

Computing  $|S|$ : We have

$$|S| = \binom{11}{3} = 165$$

Decomposition of  $E$ : We have

$$E = \{\text{Triples with 2 B and 1 W}\}$$



## Example: drawing balls (5)

Counting  $E$ :

$$|E| = \binom{5}{2} \times \binom{6}{1} = 60$$

Probability of  $E$ : We get

$$\mathbf{P}(E) = \frac{|E|}{|S|} = \frac{60}{165} = \frac{4}{11} = 36.4\%$$

Remark:

When experiment  $\equiv$  draw  $k$  objects from  $n$  objects, two choices:

- 1 Considered the ordered set of possible draws
- 2 Consider the draws as unordered

# Example: poker game (1)

**Situation:** Deck of 52 cards and

- Hand: 5 cards
- Straight: distinct consecutive values, not of the same suit

**Problem:** Compute

$P(E)$ , with  $E = \text{"Straight is drawn"}$



## Example: poker game (2)

Model: We take

- $S = \{\text{Non ordered hands of cards}\}$
- $\mathbf{P} = \text{Uniform probability on } S$

Computing  $|S|$ : We have

$$|S| = \binom{52}{5} = 2,598,960$$

Decomposition of  $E$ : We have

$$E = \{\text{Straight hands}\}$$

## Example: poker game (3)

Counting  $E$ : We have

- # possible 1,2,3,4,5:  $4^5$
- # possible 1,2,3,4,5 not of the same suit:  $4^5 - 4$
- # possible values of straights: 10

Thus

$$|E| = 10(4^5 - 4) = 10,200$$

Probability of  $E$ : We get

$$P(E) = \frac{|E|}{|S|} = \frac{10(4^5 - 4)}{\binom{52}{5}} = 0.39\%$$

# Example: roommate pairing (1)

**Situation:** We have

- A football team with 20 Offensive and 20 Defensive players
- Players are paired by 2 for roommates
- Pairing made at random

**Problem:** Find probability of

- ① No offensive-defensive roommate pairs
- ②  $2i$  offensive-defensive roommate pairs

## Example: roommate pairing (2)

**Model:** We take

- $S = \{\text{Non ordered pairings of 40 players}\}$
- $\mathbf{P} = \text{Uniform probability on } S$

**Computing  $|S|$ :** We have

$$|S| = \frac{1}{20!} \binom{40}{2, 2, \dots, 2} = \frac{40!}{2^{20} 20!} \simeq 3.20 \cdot 10^{23}$$

**First event  $E_0$ :** We set

$$E_0 = \{\text{No Offensive-Defensive pairing}\}$$

## Example: roommate pairing (3)

Counting  $E_0$ : We have

$$\begin{aligned} |E_0| &= (\# \text{ O-O pairings}) \times (\# \text{ D-D pairings}) \\ &= \left( \frac{20!}{2^{10} 10!} \right)^2 \end{aligned}$$

Computing  $\mathbf{P}(E_0)$ :

$$\mathbf{P}(E_0) = \frac{|E_0|}{|S|} = \frac{(20!)^3}{(10!)^2 40!} \approx 1.34 \cdot 10^{-6}$$

## Example: roommate pairing (4)

Events  $E_{2i}$ : We set

$$E_{2i} = \{2i \text{ Offensive-Defensive pairings}\}$$

Counting  $E_{2i}$ : We have

- # selections of  $2i$  O &  $2i$  D:  $\binom{20}{2i}^2$
- #  $2i$  O-D pairings:  $(2i)!$
- #  $(20 - i)$  O & D intra-pairings:  $\left(\frac{(20-2i)!}{2^{10-i} (10-i)!}\right)^2$

Thus we get

$$|E_{2i}| = \binom{20}{2i}^2 (2i)! \left(\frac{(20-2i)!}{2^{10-i} (10-i)!}\right)^2$$



## Example: roommate pairing (5)

Computing  $\mathbf{P}(E_{2i})$ :

$$\mathbf{P}(E_{2i}) = \frac{|E_{2i}|}{|S|} = \frac{\binom{20}{2i}^2 (2i)! \left( \frac{(20-2i)!}{2^{10-i} (10-i)!} \right)^2}{\frac{40!}{2^{20} 20!}}$$

Some values of  $\mathbf{P}(E_{2i})$ :

$$\mathbf{P}(E_0) \simeq 1.34 \cdot 10^{-6}$$

$$\mathbf{P}(E_{10}) \simeq 0.35$$

$$\mathbf{P}(E_{20}) \simeq 7.6 \cdot 10^{-6}$$

# Example: husband-wife placement (1)

**Situation:** We have

- A round table
- 10 married couples
- Placement at random

**Problem:** Find probability that

- 1  $n$  couples sit next to each other
- 2 No husband sits next to his wife

## Example: husband-wife placement (2)

**Model:** We take

- $S = \{\text{Permutations of 20 persons}\} / \{\text{Cyclic transformations}\}$
- $\mathbf{P} = \text{Uniform probability on } S$

**Computing  $|S|$ :** We have

$$|S| = \frac{20!}{20} = 19!$$

**Events  $E_i$ :** We set

$$E_i = \{\textit{i}th \text{ husband sits next to his wife}\}$$

## Example: husband-wife placement (3)

**Basic idea:** Let  $i_1 < \dots < i_n$ . Then on  $E_{i_1} \cdots E_{i_n}$

- The  $n$  couples  $i_1, \dots, i_n$  are considered as one entity
- We are left with the placement of  $20 - n$  entities

**Counting  $E_{i_1} \cdots E_{i_n}$ :** We have

- # placements of  $(20 - n)$  entities:  $(20 - n - 1)!$
- # wife-husband placements next to each other:  $2^n$

Thus

$$|E_{i_1} \cdots E_{i_n}| = 2^n (19 - n)!$$

## Example: husband-wife placement (4)

Second event,  $n$  couples sit together: For  $1 \leq n \leq 10$ , define

$$\begin{aligned} A_n &= \{n \text{ couples sitting next to each other}\} \\ &= \bigcup_{1 \leq i_1 < \dots < i_n \leq 10} (E_{i_1} \cdots E_{i_n}) \end{aligned}$$

Then

$$\begin{aligned} \mathbf{P}(A_n) &= \sum_{1 \leq i_1 < \dots < i_n \leq 10} \mathbf{P}(E_{i_1} \cdots E_{i_n}) \\ \mathbf{P}(A_n) &= \binom{10}{n} \frac{2^n (19-n)!}{19!} \end{aligned}$$

## Example: husband-wife placement (5)

Third event, no couple sits together: Define

$$A_0 = \{\text{no couple sitting next to each other}\}$$

Then

$$\begin{aligned} A_0^c &= \{\text{at least one couple sitting next to each other}\} \\ &= \bigcup_{i=1}^{10} E_i \end{aligned}$$

## Example: husband-wife placement (6)

Computing  $\mathbf{P}(A_0^c)$ : Thanks to Proposition 8

$$\begin{aligned}\mathbf{P}(A_0^c) &= \mathbf{P}\left(\bigcup_{i=1}^{10} E_i\right) \\ &= \sum_{n=1}^{10} (-1)^{n+1} \sum_{1 \leq i_1 < \dots < i_n \leq 10} \mathbf{P}(E_{i_1} \cdots E_{i_n}) \\ &= \sum_{n=1}^{10} (-1)^{n+1} \binom{10}{n} \frac{2^n (19-n)!}{19!}\end{aligned}$$

Computing  $\mathbf{P}(A_0)$ : We get

$$\mathbf{P}(A_0) = 1 + \sum_{n=1}^{10} (-1)^n \binom{10}{n} \frac{2^n (19-n)!}{19!}$$

# Outline

- 1 Introduction
- 2 Sample space and events
- 3 Axioms of probability
- 4 Some simple propositions
- 5 Sample spaces having equally likely outcomes
- 6 Probability as a continuous set function**



# Probabilities for increasing sequences

## Proposition 12.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- An **increasing** family of events  $\{E_i; i \geq 1\}$
- Set  $\lim_{n \rightarrow \infty} E_n = \bigcup_{i=1}^{\infty} E_i$

Then

$$\mathbf{P} \left( \lim_{n \rightarrow \infty} E_n \right) = \lim_{n \rightarrow \infty} \mathbf{P} (E_n)$$

# Proof (1)

Decomposition with exclusive sets: Define

$$F_n = E_n E_{n-1}^c$$

Then the  $F_i$  are mutually exclusive and we have

$$\begin{aligned}\bigcup_{i=1}^{\infty} E_i &= \bigcup_{i=1}^{\infty} F_i \\ \bigcup_{i=1}^n E_i &= \bigcup_{i=1}^n F_i\end{aligned}$$

## Proof (2)

Computation for  $\mathbf{P}(\lim_{n \rightarrow \infty} E_n)$ :

$$\begin{aligned}\mathbf{P}\left(\lim_{n \rightarrow \infty} E_n\right) &= \mathbf{P}\left(\bigcup_{i=1}^{\infty} E_i\right) \\ &= \mathbf{P}\left(\bigcup_{i=1}^{\infty} F_i\right) \\ &= \sum_{i=1}^{\infty} \mathbf{P}(F_i) \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \mathbf{P}(F_i) \\ &= \lim_{n \rightarrow \infty} \mathbf{P}\left(\bigcup_{i=1}^n F_i\right) \\ &= \lim_{n \rightarrow \infty} \mathbf{P}\left(\bigcup_{i=1}^n E_i\right) \\ &= \lim_{n \rightarrow \infty} \mathbf{P}(E_n)\end{aligned}$$

# Probabilities for decreasing sequences

## Proposition 13.

Let

- $\mathbf{P}$  a probability on a sample space  $S$
- An **decreasing** family of events  $\{E_i; i \geq 1\}$
- Set  $\lim_{n \rightarrow \infty} E_n = \bigcap_{i=1}^{\infty} E_i$

Then

$$\mathbf{P} \left( \lim_{n \rightarrow \infty} E_n \right) = \lim_{n \rightarrow \infty} \mathbf{P} (E_n)$$