

# Math 353, Practice Midterm 1

Name: \_\_\_\_\_

This exam consists of 8 pages including this front page.

## Ground Rules

1. You may take 3 by 5 cards two sides as note.
2. No calculator is allowed.
3. Show your work for every problem unless otherwise stated.

<i>Score</i>		
1	10	
2	15	
3	15	
4	20	
5	20	
6	20	
<i>Total</i>	100	

1. The following are true/false questions. You don't have to justify your answers. Just write down either T or F in the table below.  $A, B, C, X, b$  are always matrices here.

- (a) Let  $W_1, W_2$  be subspaces of a vector space  $V$ , then  $W_1 \cap W_2$  is a subspace of  $V$ .
- (b) Let  $V$  be a vector space of dimension  $n$ . Then any set of  $m$  vectors with  $m < n$  is linearly independent.
- (c) Let  $T : V \rightarrow W$  be a linear transformation of vector spaces  $V$  and  $W$ . If  $S \subset V$  is a basis of  $V$ , then  $T(S)$  spans  $R(T)$ .
- (d) Let  $\beta$  be a basis of  $V$  and  $T : V \rightarrow V$  a linear transformation. Write  $A = [T]_\beta$ . Then  $T$  is an isomorphism if and only if  $N(A) = \{0\}$ .
- (e) Let  $A \in M_{n \times n}(F)$  be a square matrix. Then the linear transformation  $L_A : F^n \rightarrow F^n$  is an isomorphism if and only if  $A$  is invertible.

	(a)	(b)	(c)	(d)	(e)
Answer	T	F	T	T	T

2. Multiple Choice,  $A$ ,  $B$ ,  $C$ ,  $X$ ,  $b$  are always matrices here:

- (i) Which of the following statement is correct.
- (a) Any set of 4 vectors in  $F^4$  is a basis of  $F^4$
  - (b) Any set of 4 vectors in  $F^3$  is linearly dependent.
  - (c) Any set of 2 vectors in  $F^3$  is linearly independent.
  - (d) Any set of 5 vectors in  $F^4$  must span  $F^4$
  - (e) Any linearly independent subset of  $F^3$  is a basis of  $F^3$

The correct answer is (b).

- (ii) Which of the given subsets of  $\mathbb{R}_3$  is a subspace?
- (a) The set of all vectors of the form  $(a, b, c)$  such that  $2a + b = c$
  - (b) The set of all vectors of the form  $(a, b, c)$  such that  $a + b < c$
  - (c) The set of all vectors of the form  $(a, b, c)$  such that  $2a + b = 1$
  - (d) The set of unit sphere.
  - (e) The set of all vectors of the form  $(a, a^2, a^3)$ .

The correct answer is (a).

(iii) Which of the following map is a linear transformation

- (a)  $L : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  by  $L\left(\begin{pmatrix} x \\ y \\ z \end{pmatrix}\right) = \begin{pmatrix} x + y \\ y + 1 \end{pmatrix}$ .
- (b)  $T : P_3(\mathbb{R}) \rightarrow \mathbb{R}$  by  $T(f(x)) = \int_0^1 f(x)^2 dx$
- (c)  $S : P_3(\mathbb{R}) \rightarrow \mathbb{R}$  by  $S(f(x)) = f'(3) + 2$ .
- (d)  $L : M_{2 \times 2}(F) \rightarrow M_{2 \times 2}(F)$  by  $L(X) = AXA$  where  $A = \begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix}$ .
- (e)  $L : \mathbb{R} \rightarrow \mathbb{R}$  by  $L(x) = e^x$ .

The correct answer is (d).

(iv) Let  $T : V \rightarrow W$  be a linear transformation. Assume that  $\dim V = \dim W = n$  and let  $\alpha \subset V$  and  $\beta \subset W$  be bases. Then which of the following statement is NOT equivalent to that  $T$  is an isomorphism

- (a)  $T$  is one-to-one.

- (b)  $T$  is onto.
- (c) The matrix  $[T]_{\alpha}^{\beta}$  is invertible.
- (d)  $T$  sends *any* linearly independent set of  $V$  to linearly independent set in  $W$ .
- (e)  $T$  sends *any* linearly dependent set of  $V$  to linearly dependent set in  $W$ .

The correct answer is (e).

- (v) Let  $A$  be a  $3 \times 5$ -matrix of real numbers and  $A \neq 0$ . Which of the following statement is FALSE?
  - (a)  $L_A$  define by  $L_A(x) = Ax$  is a linear transformation from  $\mathbb{R}^3$  to  $\mathbb{R}^5$ .
  - (b)  $\text{rank}(A) \leq 3$
  - (c) If  $\alpha$  and  $\beta$  are the standard bases for  $\mathbb{R}^5$  and  $\mathbb{R}^3$  respectively then  $[L_A]_{\alpha}^{\beta} = A$ .
  - (d)  $\text{Nullity}(A) \geq 2$ .
  - (e)  $L_A$  can not be one-to-one.

The correct answer is (a).

3. Let  $p_1 = 1+x$ ,  $p_2 = x+2x^2-x^3$ ,  $p_3 = 1+2x+2x^2-x^3$ ,  $p_4 = 1+2x+3x^2+x^3$  be polynomials in  $P_3(\mathbb{R})$ . Find a basis for  $\text{Span}\{p_1, p_2, p_3, p_4\} \subset P_3(\mathbb{R})$ .

*Solutions:* Consider equation of vectors  $a_1p_1 + a_2p_2 + a_3p_3 + a_4p_4 = 0$  with  $a_i$  being unknowns. compare the coefficients of each degree, we arrive the following system of linear equations

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 1 & 1 & 2 & 2 \\ 0 & 2 & 2 & 3 \\ 0 & -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

Use elementary operation to simplify the equation, we arrive

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

Then the equation has a solution  $a_4 = 0$ ,  $a_1 = 1$ ,  $a_2 = 1$ ,  $a_3 = -1$  So it is clear that  $p_1, p_2, p_3$  are linear dependent. If we remove  $p_3$  or equivalently set  $a_3 = 0$ . We see that this forces  $a_1 = a_2 = a_4 = 0$ . So  $p_1, p_2, p_4$  are linearly independent and hence a basis for  $\text{Span}\{p_1, p_2, p_3, p_4\}$ .

4. Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation such that  $T \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$  and

$$T \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}.$$

(a) Find  $T \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ .

(b) Find the standard matrix representing  $T$ .

(c) Find nullity of  $T$  and the rank of  $T$ .

(d) Given a basis  $\beta = \left\{ \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right\}$ . Find matrix  $[T]_\beta$ .

*Solutions:*

(a) Solve equation  $\begin{pmatrix} 2 \\ 1 \end{pmatrix} = x \begin{pmatrix} 1 \\ 1 \end{pmatrix} + y \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ . We see  $x = 3, y = -1$ . Then

$$T \begin{pmatrix} 2 \\ 1 \end{pmatrix} = 3T \begin{pmatrix} 1 \\ 1 \end{pmatrix} - T \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 4 \\ 2 \end{pmatrix}.$$

(b) We find  $\begin{pmatrix} 1 \\ 0 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ 2 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \end{pmatrix} = - \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 2 \end{pmatrix}$ . So similarly as the above, we have

$$T \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \quad T \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

So the standard matrix representing  $T$  is just  $A = \begin{pmatrix} 2 & 0 \\ 1 & 0 \end{pmatrix}$ .

(c) Since rows in  $A$  are just multiple to each other, row space of  $A$  only has 1 linearly independent vector. So  $\text{rank}(A) = \text{rank}(T) = 1$  and the nullity of  $T$  is  $2 - \text{rank}(A) = 1$ .

(d) Let  $Q = \begin{pmatrix} 2 & 1 \\ 1 & 0 \end{pmatrix}$ . Then  $[T]_\beta = Q^{-1}AQ = \begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix}$ .

5. Let  $v_1, \dots, v_n \in F^n$  and  $A = (v_1, \dots, v_n)$  be the  $n \times n$ -matrix. Show that  $\{v_1, \dots, v_n\}$  is a basis of  $F^n$  if and only if  $A$  is an invertible matrix via the following steps:
- (a) Show that there exists a linear transformation  $T : F^n \rightarrow F^n$  so that  $T(e_i) = v_i$  for all  $i$ . Here  $\alpha := \{e_1, \dots, e_n\}$  is the standard basis of  $F^n$ .
  - (b) Show that  $[T]_\alpha = A$ .
  - (c) Show that if  $\{v_1, \dots, v_n\}$  is a basis of  $F^n$  then there exists a linear transformation  $U : F^n \rightarrow F^n$  so that  $U(v_i) = e_i$  for all  $i$ .
  - (d) Show that  $\{v_1, \dots, v_n\}$  is a basis of  $F^n$  if and only if  $A$  is an invertible matrix.

*Proof:*

- (a) Since  $\{e_1, \dots, e_n\}$  is the standard basis of  $F^n$ . By Theorem 2.6, there exists unique linear transformation  $T : F^n \rightarrow F^n$  such that  $T(e_i) = v_i$  for all  $i = 1, \dots, n$ .
- (b) Since  $\alpha$  is the standard basis of  $F^n$ , we always have  $v = [v]_\alpha$  for any  $v \in F^n$ . So

$$[T]_\alpha = ([T(e_1)]_\alpha, \dots, [T(e_n)]_\alpha) = ([v_1]_\alpha, \dots, [v_n]_\alpha) = (v_1, \dots, v_n) = A.$$

- (c) If  $v_1, \dots, v_n$  is a basis of  $F^n$  then we can apply Theorem 2.6 to  $\{v_i\}$  and  $\{e_i\}$  again, which implies that there exists a unique linear transformation  $U : F^n \rightarrow F^n$  so that  $U(v_i) = e_i$  for all  $i = 1, \dots, n$ .
- (d) Applies Theorem 2.18 to  $T$ , we see  $T$  is an isomorphism if and only if  $A = [T]_\alpha$  is invertible. Now if  $v_1, \dots, v_n$  is a basis, then (c) constructed a linear transformation  $U : F^n \rightarrow F^n$ . Note that  $UT(e_i) = I_{F^n}(e_i)$  for all  $i$ . Using Theorem 2.6 again,  $UT$  is necessarily  $I_{F^n}$ . Similarly, we see that  $TU = I_{F^n}$ . So  $T$  is invertible and hence  $A = [T]_\alpha$  is invertible. Conversely, if  $A$  is invertible then  $T$  is invertible, which means it is one-to-one and onto. In particular, by Theorem 2.2,  $\{v_i = T(e_i), i = 1, \dots, n\}$  spans  $F^n$ . Since  $\{v_1, \dots, v_n\}$  has  $n$  vectors, this implies that  $\{v_1, \dots, v_n\}$  is a basis of  $F^n$ .

6. Let  $V, W$  be vector spaces over field  $F$ . Let  $v_1, \dots, v_n \in V$  be *linearly independent* vectors.
- Show that given  $w_1, \dots, w_n \in W$  then there exists a linear transformation  $T : V \rightarrow W$  such that  $T(v_i) = w_i$  for all  $i = 1, \dots, n$ .
  - Could we drop the assumption of linear independence for so that the above statement is still true? why or why not?
  - Is  $T$  necessarily unique, why or why not?

*Solutions:*

- Proof:* Since  $v_i$  are linearly independent, then one can always extend  $v_i$ , say  $\{v_1, \dots, v_n, v_{n+1}, \dots, v_m\}$ , to be a basis of  $V$ . Now extend  $w_i$  to  $m$ -vectors  $w_1, \dots, w_n, w_{n+1}, \dots, w_m$ . For example,  $w_{n+1} = \dots = w_m = 0$ . Now by Theorem 2.6, there exists a unique linear transformation  $T : V \rightarrow W$  so that  $T(v_i) = w_i$  for all  $i = 1, \dots, m$ .
- No. We can not drop the assumption that  $v_i$  are linearly independent. For example, let  $V = W = \mathbb{R}$ ,  $v_1 = v_2 = 1$ , but  $w_1 = 0$  and  $w_2 = 2$ . It is not possible to define a linear transformation  $T : \mathbb{R} \rightarrow \mathbb{R}$  so that  $T(v_i) = w_i$ .
- Contrary to Theorem 2.6,  $T$  here in general is not unique as  $v_i$  may not form a basis. For example, let  $V = W = \mathbb{R}^2$ ,  $v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $w_1 = v_1$ . We see  $I_V(v_1) = w_1$ . On the other hand, consider linear transformation  $T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \\ 0 \end{pmatrix}$ . We still have  $T(v_1) = w_1$ .