

THE INTEGRAL OF THE NORMAL AND FLUXES OVER SETS OF FINITE PERIMETER

IDO BRIGHT AND MONICA TORRES

ABSTRACT. Given two intersecting sets of finite perimeter, E_1 and E_2 , with unit normals ν_1 and ν_2 respectively, we obtain a bound on the integral of ν_1 over the reduced boundary of E_1 inside E_2 . This bound depends only on the perimeter of E_2 . For any vector field $\mathbf{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ with the property that $\mathbf{F} \in L^\infty$ and $\operatorname{div} \mathbf{F}$ is a (signed) Radon measure, we obtain bounds on the flux of \mathbf{F} over the portion of the reduced boundary of E_1 inside E_2 . These results are then applied to study the limit of surfaces with perimeter growing to infinity.

1. INTRODUCTION

An application of the Gauss-Green Theorem shows that the integral over the reduced boundary of any set of finite perimeter $E \subset \mathbb{R}^n$ of the normal vector field is the zero vector (see Lemma 3.3); namely,

$$(1.1) \quad \int_{\partial^* E} \nu(x) d\mathcal{H}^{n-1}(x) = \mathbf{0}.$$

With the aid of (1.1) we prove in this paper an estimate of the integral of the normal over a surface that lies on the boundary of a set of finite perimeter (see Theorem 3.2). More precisely, given two intersecting sets of finite perimeter E_1 and E_2 with unit normals ν_1 and ν_2 respectively, we show in Theorem 3.2 that

$$(1.2) \quad \left| \int_{\partial^* E_1 \cap E_2^1} \nu_1(x) d\mathcal{H}^{n-1} \right| \leq \frac{1}{2} \mathcal{H}^{n-1}(\partial^* E_2),$$

where $\partial^* E_1$ is the reduced boundary of E_1 and E_2^1 is the measure-theoretic interior of E_2 . The significance of this result is that the bound depends only on the perimeter of E_2 . We then use estimate (1.2) to study the limit of sets with perimeter growing to infinity.

Given a vector field \mathbf{F} , we ask the question whether we can estimate the integral of $\mathbf{F} \cdot \nu$ over a surface S that lies on the boundary of a set of finite perimeter E . That is, we would like to estimate the flux of the vector field \mathbf{F} across S . This is a delicate question and the answer depends on the regularity of \mathbf{F} . Indeed, if the vector field \mathbf{F} is continuous then $\mathbf{F} \cdot \nu$ is defined on any surface $S \subset \mathbb{R}^n$. However, a vector field $\mathbf{F} \in L^p(\mathbb{R}^n; \mathbb{R}^n)$ is defined only up to a set of Lebesgue measure zero and therefore, given any surface $S \subset \mathbb{R}^n$, such \mathbf{F} might not even be defined on S . However, it has been proved in Chen-Torres-Ziemer [7, 6] and Šilhavý [14] that if $\mathbf{F} \in L^\infty(\mathbb{R}^n; \mathbb{R}^n)$ satisfies that $\operatorname{div} \mathbf{F}$ is a Radon measure (we will refer to these vector fields as *divergence-measure fields*), then \mathbf{F} has an interior normal trace on the reduced boundary of any set of finite perimeter E . This normal trace is a function, denoted as $\mathcal{F} \cdot \nu$, that satisfies $\mathcal{F} \cdot \nu \in L^\infty(\partial^* E)$. Therefore, the flux of a divergence-measure vector field $\mathbf{F} \in L^\infty$ through S ,

$$\int_S \mathcal{F} \cdot \nu(x) d\mathcal{H}^{n-1}(x)$$

Key words and phrases. Gauss-Green theorem, divergence-measure fields, sets of finite perimeter, normal traces, shape optimization, occupational measures.

is defined on any surface S that is the boundary of a set of finite perimeter. In Theorems 4.2 and 4.3 we obtain estimates on the flux of bounded divergence-measure vector fields using the Gauss-Green formula proved in [7, 6, 14] (see also Frid [11] and the references therein). We refer to the book of Dafermos [8, Chapter 1] and [6] for the connections between fluxes of divergence-measure fields and the theory of hyperbolic conservation laws.

We then apply our main Theorem 3.2 to the analysis of sets with perimeter growing to infinity (see Theorem 5.2). Our results rely on the analysis of occupational measures, which are probability measures essentially supported on the boundaries of sets of finite perimeter. Occupational measures appear in the study of stochastic processes, and also in the context of optimization in the study of infinite horizon optimal control (see Finlay-Gaitsgory-Lebedev [10], Artstein-Bright [2] and Gaitsgory-Quincampoix [12]).

The aforementioned bounds extend previous bounds by Bright-Lee [3] from the smooth to non-smooth settings. Some of the results on sets of finite perimeter presented in this paper, and in particular Theorem 5.2, will be used by the authors to study shape optimization problems in a forthcoming paper.

The organization of this paper is as follows. In Section 2 we present some facts about sets of finite perimeter and we state the Gauss-Green formula that we will use in the paper, as well as other results. In Section 3 we obtain our estimate on the integral of the normal. In Section 4 we show that the same technique used in Section 3 can be applied to estimate the flux of bounded divergence-measure fields. Finally, in Section 5 we introduce occupational measures and use them to study the limit of sets with perimeter growing to infinity.

2. SETS OF FINITE PERIMETER AND THE GAUSS-GREEN FORMULA FOR DIVERGENCE-MEASURE FIELDS

In this section we first recall some properties of Radon measures, sets of finite perimeter, and related BV functions ([1, 9]). For the sake of completeness, we start with some basic notions and definitions. First, denote by \mathcal{H}^{n-1} the $(n-1)$ -dimensional Hausdorff measure in \mathbb{R}^n , and by \mathcal{L}^n the Lebesgue measure in \mathbb{R}^n (recall that $\mathcal{L}^n = \mathcal{H}^n$). The space $\mathcal{M}(\Omega)$ consists of all finite Radon measures μ in the open set Ω ; that is, the total variation of μ , denoted as $\|\mu\|$, satisfies $\|\mu\|(\Omega) < \infty$. For any set $E \subset \mathbb{R}^n$, we denote by \bar{E} and ∂E its topological closure and boundary, respectively. We denote by $E \Subset \mathbb{R}^n$ that the closure of E is compact and contained in \mathbb{R}^n . The complement of the set E is denoted by $E^c = \mathbb{R}^n \setminus E$. Also, we denote $B(x, r)$ as the open ball of radius r and center at x . The symmetric difference of sets is denoted by

$$A \Delta B := (A \setminus B) \cup (B \setminus A).$$

Definition 2.1. For every $\alpha \in [0, 1]$ and every \mathcal{L}^n -measurable set $E \subset \mathbb{R}^n$, define

$$(2.1) \quad E^\alpha := \{y \in \mathbb{R}^n : D(E, y) = \alpha\},$$

where

$$(2.2) \quad D(E, y) := \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap B(y, r))}{\mathcal{L}^n(B(y, r))}.$$

Then E^α is the set of all *points with density* α . We define the *measure-theoretic boundary* of E , $\partial^m E$, as

$$(2.3) \quad \partial^m E := \mathbb{R}^n \setminus (E^0 \cup E^1).$$

Definition 2.2. Let $E \subset \mathbb{R}^n$. We say that E is a *set of finite perimeter* in Ω if

$$\text{Per}(E, \Omega) := \sup \left\{ \int_E \text{div} \varphi dx : \varphi \in C_c^1(\Omega), \|\varphi\|_\infty \leq 1 \right\} < \infty.$$

We say that E is a set of *locally finite perimeter* if it is a set of *finite perimeter* in every $\Omega \Subset \mathbb{R}^n$.

Definition 2.3. A function $f \in L^1(\Omega)$ is called a function of bounded variation if its distributional derivative Df is a finite Radon vector measure; that is, $\|Df\|(\Omega) < \infty$, where $\|Df\|$ is the total variation of Df . Notationally, we write $f \in BV(\Omega)$.

Remark 2.4. If E is a set of finite perimeter in \mathbb{R}^n and E has finite Lebesgue measure in \mathbb{R}^n , namely, $\mathcal{L}^n(E) < \infty$ then its characteristic function, denoted as χ_E , is in $BV(\mathbb{R}^n)$. Moreover,

$$(2.4) \quad \|D\chi_E\| = \mathcal{H}^{n-1} \llcorner \partial^* E,$$

where $\partial^* E$ is the reduced boundary of the set E (see the definition below).

Definition 2.5. Let E be a set of finite perimeter in \mathbb{R}^n . The *reduced boundary* of E , denoted as $\partial^* E$, is the set of all points $y \in \mathbb{R}^n$ such that

- (1) $\|D\chi_E\|(B(y, r)) > 0$ for all $r > 0$;
- (2) The limit $\nu_E(y) := \lim_{r \rightarrow 0} \frac{D\chi_E(B(y, r))}{\|D\chi_E\|(B(y, r))}$ exists and $|\nu_E(y)| = 1$.

Remark 2.6. Throughout the paper we use indistinctly the notation

$$Per(E) = Per(E, \mathbb{R}^n) = \mathcal{H}^{n-1}(\partial^* E)$$

to denote the perimeter of the set E .

The unit vector, $\nu_E(y)$, is called the *measure-theoretic interior unit normal* to E at y (we sometimes write ν instead of ν_E for notational simplicity). In view of the following, we see that ν is aptly named because $\nu(y)$ satisfies

$$D(\{x : (x - y) \cdot \nu(y) > 0, x \notin E\} \cup \{x : (x - y) \cdot \nu(y) < 0, x \in E\}, y) = 0,$$

which means that $\nu_E(y)$ is the interior unit normal to E in the sense that E (in the limit and in measure) lies in the appropriate half-space determined by the hyperplane orthogonal to ν .

The following result is due to Federer (see also [15] Lemma 5.9.5. and [1], Theorem 3.61):

Theorem 2.7. *If E is a set of finite perimeter in \mathbb{R}^n , then*

$$(2.5) \quad \partial^* E \subset E^{\frac{1}{2}} \subset \partial^m E, \quad \mathcal{H}^{n-1}(\mathbb{R}^n \setminus (E^0 \cup \partial^* E \cup E^1)) = 0.$$

In particular, E has density either 0 or 1/2 or 1 at \mathcal{H}^{n-1} -a.e. $x \in \mathbb{R}^n$ and \mathcal{H}^{n-1} -a.e. $x \in \partial^m E$ belongs to $\partial^ E$.*

We will refer to the sets E^0 and E^1 as the *measure-theoretic exterior and interior* of E . We note that, in general, the sets E^0 and E^1 do not coincide with the topological exterior and interior of the set E . The sets E^0 and E^1 also motivate the definition of measure-theoretic boundary. We note that (2.5) implies, for any set $E \subseteq \mathbb{R}^n$ of finite perimeter,

$$\mathbb{R}^n = E^1 \cup \partial^* E \cup E^0 \cup \mathcal{N}$$

where $\mathcal{H}^{n-1}(\mathcal{N}) = 0$.

We recall that the BV space, the space of functions of bounded variation, in fact represents equivalence classes of functions so that, when a function in a class is changed on a set of \mathcal{L}^n -measure zero, it remains in this class. The same is true for sets of finite perimeter because, by definition, the characteristic function χ_E of a set of finite perimeter, E , is a function of bounded variation. Thus, it follows that E may be altered by a set of \mathcal{L}^n -measure zero and still determine the same essential boundary $\partial^m E$.

Remark 2.8. Throughout this paper, we will choose a preferred representative for E and thereby adopt the following convention:

$$E = E^1 \cup \partial^m E$$

The following result by Šilhavý [13, Proposition 3.8.5, page 81], which is easily verified, will be needed in the sequel.

Proposition 2.9. *If E_1, E_2 are sets of finite perimeter in \mathbb{R}^n , with corresponding measure theoretic normal vectors ν_1 and ν_2 , then:*

- (1) *Up to a set of zero \mathcal{H}^{n-1} -measure*

$$\partial^m(E_1 \cap E_2) = (\partial^m E_1 \cap E_2^1) \cup (E_1^1 \cap \partial^m E_2) \cup \mathcal{S},$$

where $\mathcal{S} = \{x \in \partial^* E_1 \cap \partial^* E_2 \mid \nu_1(x) = \nu_2(x)\}$.

- (2) *Let $E_3 = E_1 \cap E_2$ then its corresponding measure theoretic normal vector ν_3 satisfies, for \mathcal{H}^{n-1} -a.e point $x \in \partial^* E_3$,*

$$\nu_3(x) = \begin{cases} \nu_1(x) & x \in \partial^* E_1 \cap E_2^1 \\ \nu_2(x) & x \in \partial^* E_2 \cap E_1^1 \\ \nu_1(x) = \nu_2(x) & x \in \mathcal{S} \end{cases}.$$

We will use the following

Lemma 2.10. *If $E \subset \mathbb{R}^n$ is a set of finite perimeter then, for every $x \in \mathbb{R}^n$ and almost every $r > 0$, $\mathcal{H}^{n-1}(\partial^* E \cap \partial B(x, r)) = 0$.*

Proof. By the coarea formula we have

$$(2.6) \quad 0 = \mathcal{L}^n(\partial^* E) = \int_0^\infty \mathcal{H}^{n-1}(\partial B(x, r) \cap \partial^* E) d\mathcal{L}(r)$$

which clearly implies the desired result. \square

Definition 2.11. The vector field $\mathbf{F} \in L^\infty(\Omega; \mathbb{R}^n)$ is called a *divergence measure field* in Ω if $\operatorname{div} \mathbf{F}$, the distributional divergence of \mathbf{F} , is a finite Radon measure in Ω . That is $\|\operatorname{div} \mathbf{F}\|(\Omega) < \infty$.

In this paper we use the Gauss-Green Formula (see (2.9) below) for bounded divergence-measure fields \mathbf{F} over sets of finite perimeter E , which was proved in Šilhavý [14], Chen-Torres [6] and Chen-Torres-Ziemer [7]. In [6], the interior normal trace $\mathcal{F} \cdot \nu \in L^\infty(\partial^* E)$ was obtained using a product rule for divergence-measure fields. In this paper we prefer to use Šilhavý's formula for the trace since it is expressed in terms of the normal vector $\nu(x)$ (see 2.7 below). In Chen-Torres-Ziemer [7], the normal trace $\mathcal{F} \cdot \nu$ was constructed with a different approach; namely, $\mathcal{F} \cdot \nu$ was obtained as the limit of the classical normal traces $\mathbf{F} \cdot \nu$, which are defined on almost every surface that approximate the reduced boundary of E . This requires to show that a set of finite perimeter E can be approximated with smooth surfaces from "one-side" of the reduced boundary with respect to the measure $\operatorname{div} \mathbf{F}$.

Theorem 2.12. *Suppose $\mathbf{F}: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a vector field with the properties $\mathbf{F} \in L^\infty(\mathbb{R}^n; \mathbb{R}^n)$ and $\operatorname{div} \mathbf{F} \in \mathcal{M}(\mathbb{R}^n)$; that is, \mathbf{F} is a divergence-measure field in \mathbb{R}^n . Let $E \subset \mathbb{R}^n$ be a set of finite perimeter. Then the interior normal trace of \mathbf{F} on $\partial^* E$, denoted as $\mathcal{F} \cdot \nu$, satisfies $\mathcal{F} \cdot \nu \in L^\infty(\partial^* E)$ and is given, for \mathcal{H}^{n-1} -almost every $x \in \partial^* E$, by the formula $\mathcal{F} \cdot \nu(x) = T_{\mathbf{F}}(x, \nu(x))$, where*

$$(2.7) \quad T_{\mathbf{F}}(x, \nu(x)) = \lim_{r \rightarrow 0} \frac{n}{\omega_{n-1} r^n} \int_{B(x, r, \nu(x))} \mathbf{F}(y) \cdot \frac{y-x}{|y-x|} d\mathcal{L}^n(y),$$

where $B(x, r, \nu(x)) = B(x, r) \cap \{y \mid (y-x) \cdot \nu(x) > 0\}$ and ω_{n-1} is the volume of the $(n-1)$ -dimensional unit ball in \mathbb{R}^{n-1} . For every Lipschitz continuous test function with compact support ϕ ,

$$(2.8) \quad - \int_{\partial^m E} \phi(x) \mathcal{F} \cdot \nu(x) d\mathcal{H}^{n-1}(x) = \int_{E^1} \phi(x) \operatorname{div} \mathbf{F} + \int_{E^1} \nabla \phi(x) \cdot \mathbf{F}(x) d\mathcal{L}^n(x).$$

In particular, if $E \in \mathbb{R}^n$ then

$$(2.9) \quad - \int_{\partial^m E} \mathcal{F} \cdot \nu(x) d\mathcal{H}^{n-1}(x) = \int_{E^1} \operatorname{div} \mathbf{F}.$$

Moreover,

$$(2.10) \quad \|\mathcal{F} \cdot \nu\|_{\infty; \partial^* E} \leq \|\mathbf{F}\|_{\infty; \mathbb{R}^n}.$$

It was proven in Chen-Torres-Ziemer [7] (and it is also implied in Šilhavý [14]) that if \mathbf{F} is continuous then $\mathcal{F} \cdot \boldsymbol{\nu} = \mathbf{F} \cdot \boldsymbol{\nu}$, the classical dot product, which is defined on any surface. Hence, we have:

Theorem 2.13. *If \mathbf{F} is continuous, the above Gauss-Green formula reduces to*

$$-\int_{\partial^m E} \phi(x) \mathbf{F}(x) \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) = \int_{E^1} \phi(x) \operatorname{div} \mathbf{F} + \int_{E^1} \nabla \phi(x) \cdot \mathbf{F}(x) d\mathcal{L}^n(x),$$

for every Lipschitz continuous test function with compact support ϕ . In particular, if $E \in \mathbb{R}^n$ we have

$$(2.11) \quad -\int_{\partial^m E} \mathbf{F}(x) \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) = \int_{E^1} \operatorname{div} \mathbf{F}.$$

3. ESTIMATES ON THE NORMAL VECTOR

In this section we verify bounds on the integrals of the normal vector over the boundary of a set of finite perimeter. We recall that throughout the paper, whenever E is a set of finite perimeter, we imply that E is the precise representative $E := E^1 \cup \partial^m E$ (see Remark 2.8).

We first prove the following:

Lemma 3.1. *Let $E \subset \mathbb{R}^n$ be a set of finite perimeter and $O \subset \mathbb{R}^n$ an open set. Then*

$$E^1 \cap O = (E \cap O)^1 \text{ and } E^0 \cap O = (E \cap O)^0.$$

Proof. Let $x \in E^1 \cap O$. Since $x \in E^1$;

$$\lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap B(x, r))}{\mathcal{L}^n(B(x, r))} = 1.$$

Since x is in the open set O , $B(x, r) \subset O$ for r small enough. Thus:

$$\begin{aligned} \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap B(x, r))}{\mathcal{L}^n(B(x, r))} &= \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap B(x, r) \cap O)}{\mathcal{L}^n(B(x, r))} \\ &= \lim_{r \rightarrow 0} \frac{\mathcal{L}^n((E \cap O) \cap B(x, r))}{\mathcal{L}^n(B(x, r))} \\ &= 1. \end{aligned}$$

This implies that $x \in (E \cap O)^1$. Conversely, let $x \in (E \cap O)^1$. Thus,

$$\lim_{r \rightarrow 0} \frac{\mathcal{L}^n((E \cap O) \cap B(x, r))}{\mathcal{L}^n(B(x, r))} = 1;$$

that is,

$$(3.1) \quad \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap (O \cap B(x, r)))}{\mathcal{L}^n(B(x, r))} = 1.$$

Notice that if a point y satisfies $y \notin O$, then

$$(3.2) \quad \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap (O \cap B(y, r)))}{\mathcal{L}^n(B(y, r))} \leq \frac{1}{2},$$

and this together with (3.1) implies that $x \in O$. Hence, since O is open it follows that $B(x, r) \subset O$ for r small enough. Thus,

$$1 = \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap (O \cap B(x, r)))}{\mathcal{L}^n(B(x, r))} = \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E \cap B(x, r))}{\mathcal{L}^n(B(x, r))},$$

and hence $x \in E^1$. We conclude that $x \in E^1 \cap O$.

The second equality follows from the fact that

$$x \in E^0 \iff x \in (E^0 \cup \partial^m E)^1.$$

□

We now state our main result:

Theorem 3.2. *Let $E_1, E_2 \subset \mathbb{R}^n$ be sets of finite perimeter, then for $F = E_2, E_2^1, E_2^0$ or $E_2^0 \cup \partial^m E_2$*

$$\left| \int_{\partial^* E_1 \cap F} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \leq \frac{\mathcal{H}^{n-1}(\partial^* E_2)}{2}.$$

In order to prove the main result we use the following Lemma.

Lemma 3.3. *Let $E \subset \mathbb{R}^n$ be a set of finite perimeter. Then,*

$$\int_{\partial^* E} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) = \mathbf{0}.$$

Proof. By replacing E by $E^0 \cup \partial^m E$, if necessary, we can assume that $\mathcal{L}^n(E) < \infty$. Let $\mathbf{F}_i \equiv (0, \dots, 1, \dots, 0)$ (the i -th position is 1) and note that \mathbf{F}_i is a continuous vector field. We consider first the case $E \Subset \mathbb{R}^n$ and use the Gauss-Green formula (2.11) to obtain,

$$\int_{E^1} \operatorname{div} \mathbf{F}_i(x) d\mathcal{L}^n(x) = 0 = - \int_{\partial^* E} \nu_i(y) d\mathcal{H}^{n-1}(y),$$

where $\boldsymbol{\nu}(y) = (\nu_1(y), \dots, \nu_n(y))$. Hence,

$$\int_{\partial^* E} \nu_i(x) d\mathcal{H}^{n-1}(x) = 0, \quad i = 1, 2, \dots, n.$$

We conclude

$$(3.3) \quad \int_{\partial^* E} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) = \mathbf{0}.$$

Otherwise, when E is not bounded, there exists $\epsilon > 0$ and an open ball B satisfying

$$(3.4) \quad \mathcal{H}^{n-1}(\partial B \cap \partial^* E) = 0, \quad \mathcal{H}^{n-1}(\partial^* E \cap B^c) < \epsilon \text{ and } \mathcal{H}^{n-1}(\partial B \cap E) < \epsilon.$$

Indeed, using Lemma 2.10, for almost every r , $\mathcal{H}^{n-1}(\partial B(\mathbf{0}, r) \cap \partial^* E) = 0$, and, clearly, $\mathcal{H}^{n-1}(\partial^* E \setminus B(\mathbf{0}, r)) < \epsilon$ for every r large enough. Also, notice that the coarea formula yields:

$$\int_0^\infty \mathcal{H}^{n-1}(\partial B(\mathbf{0}, r) \cap E) d\mathcal{L}(r) = \mathcal{L}^n(E) < \infty,$$

and hence the set $S = \{r | \mathcal{H}^{n-1}(\partial B(\mathbf{0}, r) \cap E) \geq \epsilon\}$ has finite measure. Therefore, the complement of this set, $\{r | \mathcal{H}^{n-1}(\partial B(\mathbf{0}, r) \cap E) < \epsilon\}$, has infinite measure. Hence, there exists an r such that $\mathcal{H}^{n-1}(\partial B(\mathbf{0}, r) \cap E) < \epsilon$.

For every i we obtain, using the Gauss-Green formula (2.11) and Proposition 2.9 (1),

$$(3.5) \quad \int_{(E \cap B)^1} \operatorname{div} \mathbf{F}_i(x) d\mathcal{L}^n(x) = \mathbf{0} = - \int_{\partial^* E \cap B} \nu_i(x) d\mathcal{H}^{n-1}(x) - \int_{\partial B \cap E^1} \nu_i(x) d\mathcal{H}^{n-1}(x).$$

This implies that

$$\begin{aligned} \left| \int_{\partial^* E} \nu_i(x) d\mathcal{H}^{n-1}(x) \right| &\leq \left| \int_{\partial^* E \cap B} \nu_i(x) d\mathcal{H}^{n-1}(x) \right| + \left| \int_{\partial^* E \setminus B} \nu_i(x) d\mathcal{H}^{n-1}(x) \right| \\ &\leq \left| \int_{\partial B \cap E^1} \nu_i(x) d\mathcal{H}^{n-1}(x) \right| + \epsilon, \text{ from (3.4) and (3.5)} \\ &\leq 2\epsilon, \text{ from (3.4)}. \end{aligned}$$

Since ϵ is arbitrary the proof is complete. □

Corollary 3.4. *Let $E \subset \mathbb{R}^n$ be a set of finite perimeter, $C \subset \partial^* E$. Then*

$$\left| \int_C \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| = \left| \int_{\partial^* E \setminus C} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \leq \frac{1}{2} \mathcal{H}^{n-1}(\partial^* E).$$

Proof. For any $C \subset \partial^* E$,

$$\mathbf{0} = \int_{\partial^* E} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) = \int_C \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) + \int_{\partial^* E \setminus C} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x).$$

Therefore,

$$\left| \int_C \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| = \left| \int_{\partial^* E \setminus C} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| = \alpha(C).$$

Since

$$\alpha(C) \leq \mathcal{H}^{n-1}(C) \text{ and } \alpha(C) \leq \mathcal{H}^{n-1}(\partial^* E \setminus C)$$

we obtain:

$$2\alpha(C) \leq \mathcal{H}^{n-1}(C) + \mathcal{H}^{n-1}(\partial^* E \setminus C) = \mathcal{H}^{n-1}(\partial^* E).$$

We conclude:

$$\alpha(C) \leq \frac{1}{2} \mathcal{H}^{n-1}(\partial^* E).$$

□

Proof. [Proof of Theorem 3.2] Let $E_3 = E_1 \cap E_2$ and $\boldsymbol{\nu}_1, \boldsymbol{\nu}_2$ and $\boldsymbol{\nu}_3$ be the measure theoretic normal vectors corresponding to the sets E_1, E_2 and E_3 , respectively. We first prove the theorem for $F = E_2^1$. By Lemma 3.3 and Proposition 2.9 we have that

$$\begin{aligned} (3.6) \quad \mathbf{0} &= \int_{\partial^*(E_1 \cap E_2)} \boldsymbol{\nu}_3(x) d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial^* E_1 \cap E_2^1} \boldsymbol{\nu}_1(x) d\mathcal{H}^{n-1}(x) + \int_{\partial^* E_2 \cap E_1^1} \boldsymbol{\nu}_2(x) d\mathcal{H}^{n-1}(x) + \int_S \boldsymbol{\nu}_2(x) d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial^* E_1 \cap E_2^1} \boldsymbol{\nu}_1(x) d\mathcal{H}^{n-1}(x) + \int_C \boldsymbol{\nu}_2(x) d\mathcal{H}^{n-1}(x), \end{aligned}$$

where $S = \{x \in \partial^* E_1 \cap \partial^* E_2 \mid \boldsymbol{\nu}_1(x) = \boldsymbol{\nu}_2(x)\}$, and C is the disjoint union $S \cup (\partial^* E_2 \cap E_1^1) \subset \partial^* E_2$. From Corollary 3.4 we obtain

$$\left| \int_{\partial^* E_1 \cap E_2^1} \boldsymbol{\nu}_1(x) d\mathcal{H}^{n-1}(x) \right| = \left| \int_C \boldsymbol{\nu}_2(x) d\mathcal{H}^{n-1}(x) \right| \leq \frac{\mathcal{H}^{n-1}(\partial^* E_2)}{2},$$

which proves the bound for $F = E_2^1$.

For $F = E_2^0$, the inequality follows by applying the latter result to $F = (E_2^0 \cup \partial^m E_2)^1 = E_2^0$. The proof for $F = E_2^0 \cup \partial^m E_2$ follows from Lemma 3.3 since

$$0 = \int_{\partial^* E_1} \boldsymbol{\nu}_1(x) d\mathcal{H}^{n-1}(x) = \int_{\partial^* E_1 \cap E_2^1} \boldsymbol{\nu}_1(x) d\mathcal{H}^{n-1}(x) + \int_{\partial^* E_1 \cap (E_2^0 \cup \partial^m E_2)} \boldsymbol{\nu}_1(x) d\mathcal{H}^{n-1}(x),$$

and a similar argument holds for $F = E_2 = E_2^1 \cup \partial^m E_2$.

□

A weaker result follows when E_1 has only locally finite perimeter, and E_2 has finite measure. For this we need the following well known result.

Lemma 3.5. *Let $E \subset \mathbb{R}^n$ be a set of finite perimeter with finite measure and $K \subset \mathbb{R}^n$ a convex set. Then*

$$\text{Per}(E \cap K) \leq \text{Per}(E).$$

Proof. If $K = \mathbb{R}^n$ the result is clear. Otherwise, we first assume that $K = H = \{y \in \mathbb{R}^n \mid y \cdot v \leq r\}$ is the half-space defined by $v \in \mathbb{R}^n$ and $r \in \mathbb{R}$. Let $\tilde{E} = E \setminus H$. Proceeding as in Lemma 2.10 and using Proposition 2.9 (1) it follows that, up to a set of \mathcal{H}^{n-1} -measure zero and for almost every r ,

$$\partial^* \tilde{E} = (\partial^* E \cap H^c) \cup (E \cap \partial H)$$

Therefore, by Lemma 3.3, for almost every r ,

$$(3.7) \quad \begin{aligned} \mathbf{0} &= \int_{\partial^* \tilde{E}} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) = \int_{\partial^* \tilde{E} \setminus H} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) + \int_{\partial^* \tilde{E} \cap H} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) \\ &= \int_{\partial^* E \setminus H} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) + \int_{\partial H \cap E} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y). \end{aligned}$$

(Note that since E has finite measure then, by the coarea formula, the last term in (3.7) is finite for almost every r).

Since H is a half-space, then its normal vector is in the direction $\boldsymbol{\nu}_{\partial H} \equiv -v$, and by the triangle inequality the latter expression implies

$$(3.8) \quad \mathcal{H}^{n-1}(\partial H \cap E) = \left| \int_{\partial^* E \setminus H} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) \right| \leq \int_{\partial^* E \setminus H} |\boldsymbol{\nu}(y)| d\mathcal{H}^{n-1}(y) = \mathcal{H}^{n-1}(\partial^* E \setminus H).$$

From (3.8), we conclude that, for almost every r ,

$$(3.9) \quad \begin{aligned} Per(E) &= \mathcal{H}^{n-1}(\partial^* E) = \mathcal{H}^{n-1}(\partial^* E \cap H) + \mathcal{H}^{n-1}(\partial^* E \setminus H) \\ &\geq \mathcal{H}^{n-1}(\partial^* E \cap H) + \mathcal{H}^{n-1}(\partial H \cap E) = Per(H \cap E). \end{aligned}$$

For the general case, since K is a convex set, the closure \overline{K} is an intersection of countably many half-spaces. Thus, \overline{K} is an intersection of countably many half-spaces satisfying (3.9). Writing $E \cap \overline{K} = \lim_{N \rightarrow \infty} E \cap \bigcap_{i=1}^N H_i$ (in L^1), the lower semi-continuity of the perimeter implies

$$(3.10) \quad Per(E \cap \overline{K}) \leq Per(E).$$

Moreover, since K is convex we have that K and \overline{K} are Lebesgue equivalent and thus $\partial^* K = \partial^* \overline{K}$. Moreover, $\partial K = \partial \overline{K}$. Therefore, the reduced boundary and the topological boundary of K are \mathcal{H}^{n-1} equivalent, which implies that

$$Per(E \cap K) = Per(E \cap \overline{K}) \leq Per(E).$$

□

Corollary 3.6. *Let $E_1 \subset \mathbb{R}^n$ be of locally finite perimeter and $E_2 \subset \mathbb{R}^n$ of finite perimeter. If E_2 has finite measure then, for every open ball $B = B(x, r)$,*

$$(3.11) \quad \left| \int_{\partial^* E_1 \cap E_2^1 \cap B} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \leq \frac{\mathcal{H}^{n-1}(\partial^* E_2)}{2}.$$

Furthermore, when $\mathcal{H}^{n-1}(\partial^* E_1 \cap E_2) < \infty$, and, in particular, when $E_2 \Subset \mathbb{R}^n$,

$$(3.12) \quad \left| \int_{\partial^* E_1 \cap E_2^1} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right|, \left| \int_{\partial^* E_1 \cap E_2} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \leq \frac{\mathcal{H}^{n-1}(\partial^* E_2)}{2}.$$

Proof. Let \hat{B} be an open ball so that $\overline{B} \subset \hat{B}$. Consider the sets of finite perimeter $E := (E_1 \cap \hat{B})^1 \cup \partial^m(E_1 \cap \hat{B})$ and $F = (E_2 \cap B)^1 \cup \partial^m(E_2 \cap B)$. Thus, since $\partial \hat{B} \cap (E_2^1 \cap B) = \emptyset$ and $B \subset \hat{B}$,

Proposition 2.9 (1) yields

$$\begin{aligned}
\left| \int_{\partial^* E_1 \cap E_2^1 \cap B} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| &= \left| \int_{\partial^*(E_1 \cap \hat{B}) \cap (E_2^1 \cap B)} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \\
&= \left| \int_{\partial^*(E_1 \cap \hat{B}) \cap (E_2 \cap B)^1} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right|, \text{ by Lemma 3.1} \\
&= \left| \int_{\partial^* E \cap F^1} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \\
&\leq \frac{\mathcal{H}^{n-1}(\partial^* F)}{2}, \text{ by Theorem 3.2} \\
&= \frac{\mathcal{H}^{n-1}(\partial^*(E_2 \cap B))}{2} \leq \frac{\mathcal{H}^{n-1}(\partial^* E_2)}{2}, \text{ by Lemma 3.5,}
\end{aligned}$$

which proves inequality (3.11).

When $\mathcal{H}^{n-1}(\partial^* E_1 \cap E_2) < \infty$, we fix $\epsilon > 0$ and choose an open ball B , such that

$$(3.13) \quad \mathcal{H}^{n-1}(\partial^* E_1 \cap E_2 \cap B^c) < \epsilon.$$

By the previous inequality,

$$\begin{aligned}
\left| \int_{\partial^* E_1 \cap E_2^1} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| &= \left| \int_{\partial^* E_1 \cap E_2^1 \cap B} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| + \left| \int_{\partial^* E_1 \cap E_2^1 \cap B^c} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \\
&\leq \left| \int_{\partial^* E_1 \cap E_2^1 \cap B} \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| + \epsilon, \text{ by (3.13)} \\
&\leq \frac{\mathcal{H}^{n-1}(\partial^* E_2)}{2} + \epsilon.
\end{aligned}$$

Taking ϵ to zero completes the proof of the first inequality in (3.12). For the second inequality, we proceed in the same way, integrating over $\partial^* E_1 \cap E_2$ and then replacing $E_2 \cap B$ by F . This introduces an extra term that can be controlled by (3.13). Finally, we apply Theorem 3.2 as before with F in place of E_2 . \square

4. ESTIMATES ON THE FLUX OF A VECTOR FIELD.

We recall that a vector field $\mathbf{F} \in L^p(\Omega; \mathbb{R}^n)$ is defined up to a set of \mathcal{L}^n -measure zero. In particular, given $\mathbf{F} \in L^\infty(\Omega; \mathbb{R}^n)$ and a surface $S \subset \mathbb{R}^n$, the vector field \mathbf{F} might not be defined on S . However, it is proven in [6, 7, 14] that if $\mathbf{F} \in L^\infty$ is a divergence-measure field and S is a surface that is the boundary of a set of finite perimeter E , then \mathbf{F} has an interior normal trace on $\partial^* E$, denoted as $\mathcal{F} \cdot \boldsymbol{\nu}$. Therefore, the flux of a divergence-measure vector field $\mathbf{F} \in L^\infty$,

$$(4.1) \quad \int_S \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x)$$

is defined on any surface S that is the boundary of a set of finite perimeter. The same arguments used in Section 3 and the Gauss-Green formula (2.9) yield the following result that estimate the flux of a bounded divergence-measure vector field.

Lemma 4.1. *Let $\mathbf{F} \in L^\infty(\mathbb{R}^n; \mathbb{R}^n)$ be a divergence-measure vector field satisfying $\operatorname{div} \mathbf{F} = 0$. Then for any set of finite perimeter $E \Subset \mathbb{R}^n$ and any $C \subset \partial^* E$,*

$$\int_C \mathcal{F} \cdot \boldsymbol{\nu} d\mathcal{H}^{n-1} \leq \frac{1}{2} \|\mathbf{F}\|_\infty \mathcal{H}^{n-1}(\partial^* E).$$

Proof. We proceed as in the proof of Corollary 3.4. For any $C \subset \partial^* E$, from the Gauss-Green formula (2.9) we have

$$\mathbf{0} = \int_{\partial^* E} \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) = \int_C \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) + \int_{\partial^* E \setminus C} \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x).$$

Therefore,

$$\left| \int_C \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| = \left| \int_{\partial^* E \setminus C} \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| = \alpha(C).$$

Since $\mathcal{F} \cdot \boldsymbol{\nu} \in L^\infty(\partial^* E)$ and $\|\mathcal{F} \cdot \boldsymbol{\nu}\|_{\infty; \partial^* E} \leq \|\mathbf{F}\|_{\infty; \mathbb{R}^n}$ we obtain

$$\alpha(C) \leq \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(C) \text{ and } \alpha(C) \leq \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E \setminus C),$$

which implies

$$\alpha(C) \leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E).$$

□

Theorem 4.2. *Let $\mathbf{F} \in L^\infty(\mathbb{R}^n; \mathbb{R}^n)$ be a divergence-measure vector field satisfying $\operatorname{div} \mathbf{F} = 0$. Then, for any E_1, E_2 sets of finite perimeter with $\mathcal{L}^n(E_2) < \infty$ we have*

$$(4.2) \quad \left| \int_{\partial^* E_1 \cap E_2^1} \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E_2),$$

where $\mathcal{F} \cdot \boldsymbol{\nu}$ is the interior normal trace of \mathbf{F} on $\partial^* E_1$.

Proof. The Gauss-Green formula (2.9) we use applies to bounded sets. To overcome this problem we, essentially, approximate E_2 by a bounded set \hat{E}_2 and then apply a limiting argument. Let $E_3 = E_1 \cap \hat{E}_2$ with $\hat{E}_2 := (E_2 \cap B)^1 \cup \partial^m(E_2 \cap B)$, B an open ball, and $\boldsymbol{\nu}_1, \boldsymbol{\nu}_2$ and $\boldsymbol{\nu}_3$ be the measure theoretic normal vectors corresponding to the sets E_1, \hat{E}_2 and E_3 , respectively. We emphasize here the Šilhavý's pointwise formula for the normal trace $T_{\mathbf{F}}(x, \boldsymbol{\nu}(x))$ (see (2.7)), which depends only on the point x and the normal to the set at that point, $\boldsymbol{\nu}(x)$. With the aid of this formula we can now use Proposition 2.9 to compute

$$\begin{aligned} 0 &= \int_{\partial^* E_3} T_{\mathbf{F}}(x, \boldsymbol{\nu}_3(x)) d\mathcal{H}^{n-1}(x) = \int_{\partial^*(E_1 \cap \hat{E}_2)} T_{\mathbf{F}}(x, \boldsymbol{\nu}_3(x)) d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial^* E_1 \cap \hat{E}_2^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) + \int_{\partial^* \hat{E}_2 \cap E_1^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) + \int_S T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial^* E_1 \cap \hat{E}_2^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) + \int_C T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \end{aligned}$$

where $C = (\partial^* \hat{E}_2 \cap E_1^1) \cup S$, $(\partial^* \hat{E}_2 \cap E_1^1) \cap S = \emptyset$ and $S \subset \partial^* E_1 \cap \partial^* \hat{E}_2$ is the set where $\boldsymbol{\nu}_1(x) = \boldsymbol{\nu}_2(x)$. Therefore,

$$(4.3) \quad \left| \int_{\partial^* E_1 \cap \hat{E}_2^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) \right| = \left| \int_C T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \right| \leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* \hat{E}_2),$$

from Lemma 4.1 applied to the set of finite perimeter \hat{E}_2 . Since $\hat{E}_2^1 = E_2^1 \cap B$ and $\partial^m \hat{E}_2 = \partial^m(E_2 \cap B)$, (4.3) reduces to

$$(4.4) \quad \begin{aligned} \left| \int_{\partial^* E_1 \cap E_2^1 \cap B} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) \right| &\leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^*(E_2 \cap B)) \\ &\leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E_2), \text{ by Lemma 3.5,} \end{aligned}$$

for any open ball B . We note that the condition $\mathcal{L}^n(E_2) < \infty$ is used in the previous inequality since Lemma 3.5 holds for sets of finite Lebesgue measure. We now choose concentric open balls

B_i , $\mathbb{R}^n = \bigcup_{i=1}^{\infty} B_i$. Then, with our notation $\mathcal{F} \cdot \boldsymbol{\nu}(x) := T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x))$, we have from (2.10) and the Dominated Convergence Theorem

$$\begin{aligned} \left| \int_{\partial^* E_1 \cap E_2^1} \mathcal{F} \cdot \boldsymbol{\nu} d\mathcal{H}^{n-1}(x) \right| &= \lim_{i \rightarrow \infty} \left| \int_{\partial^* E_1 \cap E_2^1 \cap B_i} \mathcal{F} \cdot \boldsymbol{\nu} d\mathcal{H}^{n-1}(x) \right| \\ &\leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E_2) \text{ by (4.4)} \end{aligned}$$

and hence we conclude that

$$\left| \int_{\partial^* E_1 \cap E_2^1} \mathcal{F} \cdot \boldsymbol{\nu} d\mathcal{H}^{n-1}(x) \right| \leq \frac{1}{2} \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E_2).$$

□

Theorem 4.3. *Let $\mathbf{F} \in L^{\infty}(\mathbb{R}^n; \mathbb{R}^n)$ be a divergence-measure vector field (i.e., $\operatorname{div} \mathbf{F} = \mu \in \mathcal{M}(\mathbb{R}^n)$). Then, for every E_1, E_2 sets of finite perimeter with $\mathcal{L}^n(E_2) < \infty$ we have*

$$(4.5) \quad \left| \int_{\partial^* E_1 \cap E_2^1} \mathcal{F} \cdot \boldsymbol{\nu}(x) d\mathcal{H}^{n-1}(x) \right| \leq \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E_2) + \|\operatorname{div} \mathbf{F}\|(E_2^1),$$

where $\mathcal{F} \cdot \boldsymbol{\nu}$ is the interior normal trace of \mathbf{F} on $\partial^* E_1$.

Proof. We proceed as in the previous proof and let $E_3 = E_1 \cap \hat{E}_2$ with $\hat{E}_2 := (E_2 \cap \hat{B})^1 \cup \partial^m(E_2 \cap \hat{B})$, where \hat{B} is an open ball. Let $\boldsymbol{\nu}_1, \boldsymbol{\nu}_2$ and $\boldsymbol{\nu}_3$ be the measure theoretic normal vectors corresponding to the sets E_1, \hat{E}_2 and E_3 , respectively. The Gauss-Green formula (2.9) yields

$$\begin{aligned} - \int_{E_3^1} \operatorname{div} \mathbf{F} &= \int_{\partial^* E_3} T_{\mathbf{F}}(x, \boldsymbol{\nu}_3(x)) d\mathcal{H}^{n-1}(x) = \int_{\partial^*(E_1 \cap \hat{E}_2)} T_{\mathbf{F}}(x, \boldsymbol{\nu}_3(x)) d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial^* E_1 \cap \hat{E}_2^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) + \int_{\partial^* \hat{E}_2 \cap E_1^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \\ &\quad + \int_S T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \\ &= \int_{\partial^* E_1 \cap \hat{E}_2^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) + \int_C T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \end{aligned}$$

where $C = (\partial^* \hat{E}_2 \cap E_1^1) \cup S$, $(\partial^* \hat{E}_2 \cap E_1^1) \cap S = \emptyset$ and $S \subset \partial^* E_1 \cap \partial^* \hat{E}_2$ is the set where $\boldsymbol{\nu}_1(x) = \boldsymbol{\nu}_2(x)$. In this case we can not apply Lemma 4.1 since \mathbf{F} is not divergence-free. However, we still have the following estimate

$$(4.6) \quad \begin{aligned} \left| \int_{\partial^* E_1 \cap \hat{E}_2^1} T_{\mathbf{F}}(x, \boldsymbol{\nu}_1(x)) d\mathcal{H}^{n-1}(x) \right| &\leq \left| \int_{E_3^1} \operatorname{div} \mathbf{F} \right| + \left| \int_C T_{\mathbf{F}}(x, \boldsymbol{\nu}_2(x)) d\mathcal{H}^{n-1}(x) \right| \\ &\leq \|\operatorname{div} \mathbf{F}\|(E_3^1) + \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(C) \\ &\leq \|\operatorname{div} \mathbf{F}\|(E_3^1) + \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* \hat{E}_2). \end{aligned}$$

We now proceed to show that

$$E_3^1 \subset E_2^1.$$

Indeed, let $x \in E_3^1$. This means that

$$\lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E_3 \cap B(x, r))}{\mathcal{L}^n(B(x, r))} = 1,$$

and since $\partial^m(E_2 \cap \hat{B})$ has \mathcal{L}^n -measure zero, this is equivalent to

$$(4.7) \quad \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E_1 \cap E_2^1 \cap \hat{B} \cap B(x, r))}{\mathcal{L}^n(B(x, r))} = 1.$$

Notice that (4.7) implies that $x \in \hat{B}$. Thus, for r small enough we have $B(x, r) \subset \hat{B}$ and (4.7) becomes

$$(4.8) \quad \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E_1 \cap E_2^1 \cap B(x, r))}{\mathcal{L}^n(B(x, r))} = 1,$$

and since

$$\frac{\mathcal{L}^n(E_1 \cap E_2^1 \cap B(x, r))}{\mathcal{L}^n(B(x, r))} \leq \frac{\mathcal{L}^n(E_2 \cap B(x, r))}{\mathcal{L}^n(B(x, r))} \leq 1,$$

it follows from (4.8) that

$$(4.9) \quad \lim_{r \rightarrow 0} \frac{\mathcal{L}^n(E_2 \cap B(x, r))}{\mathcal{L}^n(B(x, r))} = 1,$$

which means that $x \in E_2^1$. Furthermore, since $\hat{E}_2^1 = E_2^1 \cap \hat{B}$ and $\partial^m \hat{E}_2 = \partial^m(E_2 \cap \hat{B})$, (4.6) becomes

$$(4.10) \quad \left| \int_{\partial^* E_1 \cap E_2^1 \cap \hat{B}} T_{\mathbf{F}}(x, \nu_1(x)) d\mathcal{H}^{n-1}(x) \right| \leq \|\operatorname{div} \mathbf{F}\|(E_2^1) + \|\mathbf{F}\|_{\infty} \mathcal{H}^{n-1}(\partial^* E_2)$$

Since (4.10) is true for every open ball \hat{B} , we proceed as in Theorem 4.2 to conclude the desired result. \square

5. LIMITS OF SETS WITH PERIMETER GROWING TO ∞

It is well known that when a sequence of sets E_i , with $\chi_{E_i} \in BV(\mathbb{R}^n)$, converges in L^1 to a set E_0 , then E_0 satisfies

$$\mathcal{H}^{n-1}(\partial^* E_0) \leq \liminf_{i \rightarrow \infty} \mathcal{H}^{n-1}(\partial^* E_i).$$

In this section, we consider two degenerate cases. The first when the perimeter grows indefinitely, and the second when the Lebesgue measure shrinks to zero. Specifically, the limit is studied by means of occupational measures. The results from this section will be applied to shape optimization in a forthcoming paper.

For every set of finite perimeter we define a corresponding probability measure in the following manner.

Definition 5.1. We define the *occupational measure* $\mu \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ corresponding to a set of finite perimeter E by

$$\mu(U \times V) = \frac{1}{\mathcal{H}^{n-1}(\partial^* E)} \mathcal{H}^{n-1}(x \in \partial^* E \mid (x, \nu(x)) \in U \times V),$$

for every Borel sets $U \subset \mathbb{R}^n$ and $V \subset \mathbb{S}^{n-1}$.

A useful property of occupational measures is that, for every continuous function $g \in C(\mathbb{R}^n \times \mathbb{S}^{n-1})$ we have that

$$\frac{1}{\mathcal{H}^{n-1}(\partial^* E)} \int_{\partial^* E} g(x, \nu(x)) d\mathcal{H}^{n-1}(x) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} g(x, v) d\mu(x, v).$$

We endow the set of probability measures $P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ with the weak topology, namely, a sequence of measures $\mu_1, \mu_2, \dots \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ converges weakly to a measure $\mu_0 \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ if for every bounded continuous function $g \in C(\mathbb{R}^n \times \mathbb{S}^{n-1})$,

$$\lim_{i \rightarrow \infty} \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} g(x, v) d\mu_i(x, v) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} g(x, v) d\mu_0(x, v).$$

Furthermore, if the sequence $\mu_1, \mu_2, \dots \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ converges weakly to μ_0 , and $f(x, v)$ is a bounded Borel function with compact support such that the set of its discontinuity points is μ_0 -negligible, then (see Ambrosio-Fusco-Pallara [1, Proposition 1.62, pages 27-28].)

$$\lim_{i \rightarrow \infty} \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f(x, v) d\mu_i(x, v) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f(x, v) d\mu_0(x, v).$$

The space $P(K \times \mathbb{S}^{d-1})$ is compact in the weak topology, whenever $K \subset \mathbb{R}^n$ is compact (see, Billingsley [4, page 72]).

Another tool we need for the next theorem is disintegration of measures. Given a probability measure $\mu \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$, we denote its disintegration by $\mu = p \otimes \mu^x$; the marginal measure is $p \in P(\mathbb{R}^n)$, which is the push forward of the projection map $\pi : \mathbb{R}^n \times \mathbb{S}^{n-1} \rightarrow \mathbb{R}^n$; that is $p = \pi_{\#}\mu$, and $p(A) = \mu(A \times \mathbb{S}^{n-1})$ for every Borel set $A \subset \mathbb{R}^n$. The measure-valued function $\mu^x \in P(\mathbb{S}^{n-1})$ is the disintegration with respect to p , for p -almost every x . With this notation, for every Borel sets $C \subset \mathbb{R}^n$ and $D \subset \mathbb{S}^{n-1}$, we have that $\mu(C \times D) = \int_C \mu^x(D) dp(x)$.

Note that when μ is the occupational measure of a set of finite perimeter, then the disintegration is a Dirac measure p -almost everywhere.

Theorem 5.2. *Let $E_1, E_2, \dots \subset \mathbb{R}^n$ be sets of finite perimeter, with perimeter growing to infinity, namely, $\lim_{i \rightarrow \infty} \mathcal{H}^{n-1}(\partial^* E_i) = \infty$. If the corresponding sequence of occupational measures μ_1, μ_2, \dots converges weakly to $\mu_0 \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ then*

$$h(x) = \int_{\mathbb{S}^{n-1}} v d\mu_0^x(v) = 0,$$

for p_0 -almost every x , where $\mu_0 = p_0 \otimes \mu_0^x$ is the disintegration of μ_0 with respect to its projection, p_0 .

Proof. Let us fix x in the support of p_0 . By the definition of the occupational measure and Theorem 3.2 we obtain, for every i ,

$$\left| \int_{B(x,r) \times \mathbb{S}^{n-1}} v d\mu_i(y, v) \right| = \left| \frac{1}{\mathcal{H}^{n-1}(\partial^* E_i)} \int_{\partial^* E_i \cap B(x,r)} \nu(y) d\mathcal{H}^{n-1}(y) \right| \leq \frac{\mathcal{H}^{n-1}(\partial B(x,r))}{2\mathcal{H}^{n-1}(\partial^* E_i)}.$$

The weak convergence implies that

$$\left| \int_{B(x,r) \times \mathbb{S}^{n-1}} v d\mu_0(y, v) \right| = \lim_{i \rightarrow \infty} \left| \int_{B(x,r) \times \mathbb{S}^{n-1}} v d\mu_i(y, v) \right| \leq \lim_{i \rightarrow \infty} \frac{\mathcal{H}^{n-1}(\partial B(x,r))}{2\mathcal{H}^{n-1}(\partial^* E_i)} = 0,$$

for almost every $r > 0$, for which $\mu_0(\partial B(x,r) \times \mathbb{S}^{d-1}) = 0$. With the disintegration notation one obtains

$$\left| \int_{B(x,r) \times \mathbb{S}^{n-1}} v d\mu_0(y, v) \right| = \left| \int_{B(x,r)} \left(\int_{\mathbb{S}^{n-1}} v d\mu_0^y(v) \right) dp_0(y) \right| = \int_{B(x,r)} h(y) dp_0(y) = 0,$$

for almost every $r > 0$.

If the measure p_0 was the Lebesgue measure, by the Lebesgue differentiation theorem $h(x) = 0$ almost everywhere. The Lebesgue-Besicovitch differentiation theorem extends this result to Radon measures (see Evans-Gariepy [9, page 43]). \square

Theorem 5.3. *Let $E_1, E_2, \dots \subset \mathbb{R}^n$ be sets of finite perimeter. If $\lim_{i \rightarrow \infty} \mathcal{L}^n(E_i) = 0$ and the corresponding sequence of occupational measures μ_1, μ_2, \dots converges weakly to $\mu_0 \in P(\mathbb{R}^n \times \mathbb{S}^{n-1})$ then*

$$h(x) = \int_{\mathbb{S}^{n-1}} v d\mu_0^x(v) = 0$$

for p_0 -almost every x , where $\mu_0 = p_0 \otimes \mu_0^x$ is the disintegration of μ_0 with respect to its projection, p_0 .

Proof. Let us fix x in the support of p_0 . From Lemma 2.10 it follows that $\mathcal{H}^{n-1}(\partial^* E_i \cap \partial B(x, r)) = 0$, for almost every $r > 0$ and every $i = 1, 2, \dots$. Hence, Lemma 3.3 and Proposition 2.9 (1) applied to $E_i \cap B(x, r)$ imply, for almost every $r > 0$ and every $i = 1, 2, \dots$,

$$\mathbf{0} = \int_{\partial^*(E_i \cap B(x, r))} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) = \int_{\partial^* E_i \cap B(x, r)} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) + \int_{\partial B(x, r) \cap E_i} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y),$$

and therefore,

$$\begin{aligned} \left| \int_{B(x, r) \times \mathbb{S}^{n-1}} v d\mu_i(y, v) \right| &= \frac{1}{\mathcal{H}^{n-1}(\partial^* E_i)} \left| \int_{\partial^* E_i \cap B(x, r)} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) \right| \\ &= \frac{1}{\mathcal{H}^{n-1}(\partial^* E_i)} \left| \int_{E_i \cap \partial B(x, r)} \boldsymbol{\nu}(y) d\mathcal{H}^{n-1}(y) \right| \\ &\leq \frac{\mathcal{H}^{n-1}(\partial B(x, r) \cap E_i)}{\mathcal{H}^{n-1}(\partial^* E_i)}. \end{aligned}$$

We integrate both sides of the previous inequality with respect to $\mathcal{L}(r)$ and use the coarea formula to obtain

$$\begin{aligned} (5.1) \quad \int_0^1 \left| \int_{B(x, r) \times \mathbb{S}^{n-1}} v d\mu_i(y, v) \right| d\mathcal{L}(r) &\leq \frac{1}{\mathcal{H}^{n-1}(\partial^* E_i)} \int_0^1 \mathcal{H}^{n-1}(\partial B(x, r) \cap E_i) d\mathcal{L}(r) \\ &\leq \frac{1}{\mathcal{H}^{n-1}(\partial^* E_i)} \int_0^\infty \mathcal{H}^{n-1}(\partial B(x, r) \cap E_i) d\mathcal{L}(r) \\ &= \frac{\mathcal{L}^n(E_i)}{\mathcal{H}^{n-1}(\partial^* E_i)} \\ &= \frac{\mathcal{L}^n(E_i)^{\frac{1}{n}} \mathcal{L}^n(E_i)^{\frac{n-1}{n}}}{\mathcal{H}^{n-1}(\partial^* E_i)}. \end{aligned}$$

By the isoperimetric inequality there exists a constant C_1 , depending only on the dimension n , such that $\mathcal{L}^n(E)^{(n-1)/n} \leq C_1 \mathcal{H}^{n-1}(\partial^* E)$ for every set of finite perimeter E . Hence, (5.1) is bounded by $C_1 (\mathcal{L}^n(E_i))^{1/n}$. The weak convergence and Fatou's lemma imply that

$$\begin{aligned} \int_0^1 \left| \int_{B(x, r) \times \mathbb{S}^{n-1}} v d\mu_0(y, v) \right| d\mathcal{L}(r) &= \int_0^1 \lim_{i \rightarrow \infty} \left| \int_{B(x, r) \times \mathbb{S}^{n-1}} v d\mu_i(y, v) \right| d\mathcal{L}(r) \\ &\leq \liminf_{i \rightarrow \infty} \int_0^1 \left| \int_{B(x, r) \times \mathbb{S}^{n-1}} v d\mu_i(y, v) \right| d\mathcal{L}(r) \\ &\leq \liminf_{i \rightarrow \infty} C_1 (\mathcal{L}^n(E_i))^{1/n} = 0. \end{aligned}$$

Applying a disintegration argument similar to Theorem 5.2 we complete the proof. \square

Remark 5.4. Theorem 5.3 holds when $\lim_{i \rightarrow \infty} \min(\mathcal{L}^n(E_i), \mathcal{L}^n(\mathbb{R}^n - E_i^1)) = 0$, since replacing E_i by $\mathbb{R}^n - E_i^1$, when needed, only changes the sign of the measurable normal vector, and as we take the absolute value of all our integrals, it will not affect our final result.

REFERENCES

- [1] L. Ambrosio, N. Fusco, and D. Pallara. Functions of Bounded Variation and Free Discontinuity Problems. *Oxford Mathematical Monographs*. The Clarendon Press, Oxford University Press: New York, 2000.
- [2] Z. Artstein and I. Bright. Periodic optimization suffices for infinite horizon planar optimal control. *SIAM J. Control Optim.*, 48: 4963–4986, 2010.
- [3] I. Bright and J. M. Lee. A note on flux integrals over smooth regular domains. *Pacific Journal of Mathematics*, 272(2): 305–322, 2014.
- [4] P. Billingsley. Convergence of probability measures. *Wiley-Interscience*, 2009.

- [5] L. A. Caffarelli and R. de la Llave. Planelike minimizers in periodic media. *Comm. Pure Appl. Math.* 54(12): 1403–1441, 2001.
- [6] G.-Q. Chen and M. Torres. Divergence-measure fields, sets of finite perimeter, and conservation laws. *Arch. Rational Mech. Anal.*, 175(2):245-267, 2005.
- [7] G.-Q. Chen, M. Torres and W. P. Ziemer. Gauss-Green Theorem for Weakly Differentiable Vector Fields, Sets of Finite Perimeter, and Balance Laws. *Communications on Pure and Applied Mathematics*, 62(2): 242-304, 2009.
- [8] C. M. Dafermos. Hyperbolic Conservation Laws in Continuum Physics, 2nd Ed., *Springer-Verlag: Berlin*, 2005.
- [9] C. Evans and D. Gariepy. Measure Theory and Fine Properties of Functions. *CRC Press, Boca Raton, FL*, 1992.
- [10] L. Finlay, V. Gaitsgory, and I. Lebedev. Duality in linear programming problems related to deterministic long run average problems of optimal control. *SIAM J. Control Optim.*, 47: 1667–1700, 2008.
- [11] H. Frid. Remarks on the theory of the divergence-measure fields. *Quartely of Applied Mathematics*, 70(3):579-596, 2012.
- [12] V. Gaitsgory and M. Quincampoix. Linear programming approach to deterministic infinite horizon optimal control problems with discounting. *SIAM J. Control Optim.*, 48: 2480-2512, 2009.
- [13] M. Šilhavý. The mechanics and thermodynamics of continuous media. *Springer-Verlag*, 1997.
- [14] M. Šilhavý. Divergence measure fields and Cauchy’s stress theorem. *Rend. Sem. Mat. Univ. Padova*, 113:15-45, 2005.
- [15] W.P. Ziemer. Weakly Differentiable Functions, Graduate Texts in Mathematics. *Springer-Verlag: New York*, 120, 1989.

[I. BRIGHT] DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195, USA.

E-mail address: `ibright@uw.edu`

[M. TORRES] DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, 150 N. UNIVERSITY STREET, WEST LAYAYETTE, IN 47907-2067, USA. [HTTP://WWW.MATH.PURDUE.EDU/~TORRES](http://www.math.purdue.edu/~torres)

E-mail address: `torres@math.purdue.edu`