

DRAFT COMPLETE, COMMENTS WELCOME!

MODEL THEORY OF OPERATOR SYSTEMS AND C*-ALGEBRAS

THOMAS SINCLAIR

ABSTRACT. We survey the model theory of operator systems and C*-algebras.

1. INTRODUCTION

At the heart of the theory of operator algebras is the “noncommutative order” imposed on the algebra $\mathcal{B}(H)$ of bounded linear operators on some Hilbert space H by the cone of positive semidefinite operators, that is, operators T which admit a sum-of-squares decomposition $T = S_1^*S_1 + \dots + S_n^*S_n$ for some n and $S_1, \dots, S_n \in \mathcal{B}(H)$ (equivalently $n = 1$). This partial order structure, even in the finite-dimensional case, is incredibly rich and complex and its understanding is deeply connected with many important and outstanding open problems in fields as diverse as quantum information theory and quantum computing, numerical linear algebra and optimization, combinatorics, computer science, and random matrix theory. Beginning in the 1960s and 70s with work of Arveson, Choi, Effros, Lance, Kirchberg, and Stinespring it became apparent that the full power of the noncommutative order is captured not just by the “level one” order structure but the higher-level order structure imposed by taking matrix amplifications. In fact, the higher-order structure on a C*-algebra is powerful enough to capture the norm-structure as well as many properties which would be considered algebraic in nature. This conception of noncommutative order crystallized in the notion of an *operator system* first systematically studied in the groundbreaking work of Choi and Effros [9].

In any semisimple category of algebras the finite-dimensional, simple objects are of primary importance. In the case of C*-algebras, these are the (complex) matrix algebras. Thus, it was a natural theme from the very beginning of the subject to try to understand the structure of a C*-algebra by quantifying how well (or how poorly) it was algebraically approximated by direct sums of matrix algebras. With the theory of operator systems, it was realized that finite-dimensional *order* approximation through matrix algebras was an equally vital aspect of the theory, and such properties as nuclearity and exactness have become standard tools of the trade for working operator algebraists. The surprising relations discovered by celebrated work of Kirchberg between two other well-studied finite-dimensional order approximation properties, the local lifting and weak expectation properties, and a famous conjecture of Connes have helped catalyze an entirely new chapter in operator algebras through their connection with a deep problem of Tsirelson arising in quantum information theory: see Goldbring’s article in this volume.

The goal of this set of notes is to introduce the reader to some aspects of the continuous model theory of operator systems and C*-algebras mainly through the model theoretic

properties of the noncommutative order and their relation with finite-dimensional approximation properties.

This article is organized as follows.

- Section 2 gives a brief overview of the theory of operator systems. While not totally self-contained, the section aims at presenting enough material for the reader to gain a basic working familiarity with operator systems, including such topics as the Choi–Effros representation theorem, Arveson’s extension theorem, duality and the double dual, and quotients. The latter part of the section provides several examples of canonical ways of constructing operator systems. The reader may wish to consult the excellent books of Brown and Ozawa [6] and Paulsen [36] for a fuller account of the theory. The reader may also wish to read the article of Szabo in this volume for an introduction to C^* -algebras before proceeding beyond this section.
- Section 3 introduces the model theory of operator systems and C^* -algebras. The first part of the section constructs the language for operator systems, while the next few sections serve to introduce basic model-theoretic concepts such as theories, ultraproducts, and definability, illustrating these ideas in the operator systems context before concluding with a discussion of the model theory of C^* -algebras. This section is largely based on the accounts found in the paper [20] of Goldbring and the author and the monograph of Farah, et al. [15], though many of the proofs of results given here are new and the centering of the account from the perspective of matrix completion problems is also somewhat novel. A reader wishing to explore these topics further would do good to consult [5] and [15].
- Section 4 surveys some applications of the model theory developed in the Section 3 to the theory of finite-dimensional approximation properties for operator systems and C^* -algebras. The discussion of exactness and nuclearity is largely sourced from [15] and [21], while that of the local lifting property is from [20], [21], and [43]. The last part of this section on the weak expectation property is sourced from [21] and [35], though the treatment here is substantially new. We refer the reader to Goldbring’s article in this volume for the model-theoretic aspects of the local lifting and weak expectation properties in relation to the famous conjectures of Connes and Kirchberg, and the reader may also wish to consult the recent monograph of Pisier [40] on this subject.

2. GENERAL BACKGROUND ON OPERATOR SYSTEMS

Definition 2.1. An *operator system* E is a closed subspace of $\mathcal{B}(H)$ which is closed under taking adjoints and contains the unit.

Remark 2.2. Let E^h be the set of hermitian elements. From the property that E is closed under adjoints, it is immediate that E^h is a real, unital subspace with $E = \mathbb{C}E^h$. Since E contains the unit, we have that for each $x \in E^h$ we may decompose x as a difference of positive elements in E , namely as $x = (\|x\|1 + x) - \|x\|1$, which shows that $E^+ := E \cap \mathcal{B}(H)^+$ is a cone in E . In this way E^h is an ordered (real) vector space

under $x \preceq y \iff y - x \in E^+$. Moreover 1 is an *order unit*, that is, for each $x \in E^h$ $-r1 \preceq x \preceq r1$ for some $r > 0$ which is *archimedean* in the sense that $-\epsilon 1 \preceq x \preceq \epsilon 1$ for all $\epsilon > 0$ implies that $x = 0$.

So far we have just established that an operator system is a Banach space with a pleasant real, ordered structure. In the real case, these are referred to as *function systems* and were studied and classified by Kadison [24] (see [2, Chapter II] as well), while a study of the complex case was undertaken by Paulsen and Tomforde [38]. What distinguishes operator systems from function systems is the following enrichment of structure of the objects or, to put it in a glass-half-empty way, restriction of the category maps between the objects.

The enrichment comes from the following, seemingly modest, observation: if $E \subset \mathcal{B}(H)$ is an operator system, then so is $E_n := M_n(E) \subset M_n(\mathcal{B}(H)) \cong \mathcal{B}(H^{\oplus n})$, where $H^{\oplus n}$ is the direct sum of n copies of H . Thus, from each operator system there may be derived a sequence of order unit spaces $(E_n, E_n^+, 1_n)$. Moreover, there is a natural family of connecting maps defined as follows. For $a \in M_{n,k}$ define $\text{ad}(a) : E_n \rightarrow E_k$ by $\text{ad}(a) : x \mapsto a^*xa$. Notice that $\text{ad}(a)$ preserves the order structure. For $v \in M_{n,k}$ with $v^*v = 1_k$ we will often refer to $\text{ad}(v)$ as the *compression* induced by v . (Notice that vv^* is a rank k projection in M_n , so $n \geq k$.)

Given a $*$ -linear map $\varphi : E \rightarrow F$ between operator systems E and F , we can define a $*$ -linear map $\varphi_n : E_n \rightarrow F_n$ coordinate-wise by $\varphi_n([x_{ij}]) := [\varphi(x_{ij})]$. Thinking about $M_n(E) \cong M_n \otimes E$, φ_n is nothing other than $\text{id}_{M_n} \otimes \varphi$. Note that the φ_n 's commute with the structure maps (thus, compressions): $\text{ad}(a) \circ \varphi_n = \varphi_k \circ \text{ad}(a)$.

Definition 2.3. A map $\varphi : E \rightarrow F$ is said to be *n-positive* ($n=1,2,3,\dots$) if $\varphi_n(E_n^+) \subset F_n^+$ and *completely positive* if φ is n -positive for all n . We say that φ is a complete order embedding if it is unital and both φ and $\varphi^{-1} : \varphi(E) \rightarrow E$ are completely positive.

It is easy to see that n -positivity implies k -positivity for all $k \leq n$ and that positive maps are $*$ -linear.

For an operator system $E \subset \mathcal{B}(H)$, we denote by $\|\cdot\|_n$ the restriction of the operator norm on $\mathcal{B}(H^{\oplus n})$ to E_n .

Definition 2.4. A linear map $\varphi : E \rightarrow F$ is said to be *n-bounded* if $\varphi_n : (E_n, \|\cdot\|_n) \rightarrow (F_n, \|\cdot\|_n)$ is bounded, in which case we denote the norm of φ_n by $\|\varphi\|_n$. The map φ is said to be *completely bounded* if it is bounded for all n , in which case we write $\|\varphi\|_{\text{cb}} := \sup_n \|\varphi\|_n$. (Note that $\|\varphi\|_k \leq \|\varphi\|_n$ for all $k \leq n$.)

Perhaps the most important foundational fact in the theory of operator systems is that the higher-order norm structure is totally determined by the structure of the positive cones.

Proposition 2.5. *Let $E \subset \mathcal{B}(H)$ be an operator system. For $x \in E_n$ we have that*

$$\|x\|_n = \inf \left\{ t > 0 : \begin{bmatrix} t1 & x \\ x^* & t1 \end{bmatrix} \in E_{2n}^+ \right\}.$$

Proof. It clearly suffices to check the case $n = 1$. We have that $\begin{bmatrix} a & b \\ b^* & c \end{bmatrix} \in M_2(\mathcal{B}(H))^+$ if and only if $\langle a\xi, \xi \rangle + 2\operatorname{Re}\langle b\xi, \eta \rangle + \langle c\eta, \eta \rangle \geq 0$ for all $\xi, \eta \in H$ with $\|\xi\| = \|\eta\| = 1$. Since we have that

$$\|x\| = \sup\{|\langle x\xi, \eta \rangle| : \|\xi\| = \|\eta\| = 1\},$$

it follows that $\begin{bmatrix} t1 & x \\ x^* & t1 \end{bmatrix}$ is positive if and only if $2t + 2\operatorname{Re}\langle x\xi, \eta \rangle \geq 0$ for all unit-norm vectors $\xi, \eta \in H$. This, in turn, is seen to be equivalent to $t \geq |\langle x\xi, \eta \rangle|$ for all such ξ, η by making substitutions of the form $\eta \mapsto e^{is}\eta$ for $s \in \mathbb{R}$ chosen suitably. This completes the proof. \square

Exercise 2.6. A unital, 2-positive map $\varphi : E \rightarrow F$ is contractive.

Proposition 2.7. *If $\varphi : E \rightarrow F$ is a unital, $*$ -linear contractive map between operator systems, then φ is positive.*

Proof. Let $x \in E^+$ with $0 \preceq x \preceq 1$, and define $y = 2x - 1$ so that $-1 \preceq y \preceq 1$, equivalently $\|y\| \leq 1$. Since φ is unital and $*$ -linear $\varphi(y) = 2\varphi(x) - 1$ is hermitian, and $\|\varphi(y)\| \leq 1$ since φ is contractive. Thus, $-1 \preceq 2\varphi(x) - 1 \preceq 1$, which implies $\varphi(x) \succeq 0$. \square

Exercise 2.8. Any unital, contractive linear map between operator systems is $*$ -linear.

We derive the following easily as a consequence of the previous two results.

Corollary 2.9. *Every complete order embedding of operator systems is a completely isometric embedding. Every unital completely isometric embedding is a complete order embedding.*

Exercise 2.10 (Kadison–Cauchy–Schwarz). Let $E \subset \mathcal{B}(H)$ be an operator system. If $\varphi : E \rightarrow \mathcal{B}(K)$ is unital and 2-positive, then $\varphi(x)\varphi(x^*) \preceq \varphi(xx^*)$ for all $x \in E$. See [36, Chapter 3] for this and many other basic properties satisfied by completely positive maps.

Exercise 2.11. If $\varphi : E \rightarrow F$ is unital, completely positive, then $\|\varphi\|_{\text{cb}} = \|\varphi(1)\| = 1$. See [36, Proposition 3.6].

Proposition 2.12 (Choi's Theorem). *Every completely positive map $\varphi : M_n \rightarrow M_k$ is of the form $\varphi(x) = \sum_{i=1}^k a_i^* x a_i$ for some $a_1, \dots, a_k \in M_{n,k}$.*

The converse is also seen to be true as $x \mapsto a^* x a$ is completely positive. See [3, Theorem 2.21] for a proof.

Exercise 2.13. Use Choi's Theorem to deduce that if $\varphi : M_n \rightarrow M_k$ is completely positive, then the *adjoint* map $\varphi^\dagger : M_k \rightarrow M_n$ defined by the functional equation $\operatorname{tr}(\varphi(A)B) = \operatorname{tr}(A\varphi^\dagger(B))$ for all $A \in M_n, B \in M_k$ is again completely positive.

Exercise 2.14. For $E = M_2$ show that

$$\varphi : \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mapsto \begin{bmatrix} (a+d)/2 & b \\ c & (a+d)/2 \end{bmatrix}$$

is unital and 1-positive, but not 2-positive. Show that $|\lambda| = 1/2$ is the optimal constant so that

$$\varphi : \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mapsto \begin{bmatrix} (a+d)/2 & \lambda b \\ \lambda c & (a+d)/2 \end{bmatrix}$$

is completely positive.

Exercise 2.15. Let v_1, v_2 be the generators of the Cuntz algebra \mathcal{O}_2 , and consider $E := \text{span}\{1, v_1, v_2, v_1^*, v_2^*\}$. We have that $\varphi : E \rightarrow E$ given by $\varphi(a1 + bv_1 + cv_2 + dv_1^* + ev_2^*) = a1 + bv_1^* + cv_2^* + dv_1 + ev_2$ is unital and positive, but not 2-positive, as φ_2 is not contractive.

Exercise 2.16. We have that $\varphi : M_2 \rightarrow M_2$ given by transposition, $\varphi(x) := x^t$, is positive but not 2-positive.

Remarkably, the structure of higher ordered cones and connecting maps is totally sufficient to abstractly characterize operator systems, as discovered in the seminal work of Choi and Effros [9].

Definition 2.17. An *abstract operator system* $(E, E^+, 1)$ is a real ordered $*$ -vector space with archimedean order unit 1 along with a real ordered $*$ -structure $(E_n, E_n^+, 1_n)$ for each $n = 1, 2, \dots$, where $E_n \cong M_n \otimes E$, $(a \otimes x)^* = a^* \otimes x^*$, and $1_n = I_n \otimes 1$, so that $E_n^+ \subset E_n^h$ and $a^* E_n^+ a \subset E_k^+$ for all $a \in M_{n,k}$.

Remark 2.18. Omitting the requirement of an order unit in the previous definition leads to the definition of a *matrix ordered $*$ -vector space*

Theorem 2.19 (Choi and Effros). *Every abstract operator system E admits a complete order embedding into $\mathcal{B}(H)$ for some Hilbert space H .*

The representation obtained by the Choi–Effros theorem is canonical, but, like the Gelfand–Naimark theorem, it is impractical to work with. Nonetheless, the abstract characterization of operator provides us with a way to axiomatize the class of operator systems as metric structures. Additionally, the proof of this theorem introduces some important ideas, as we will see in the following sketch.

To begin, we introduce the notion of a *state* $\varphi : E \rightarrow \mathbb{C}$ to just mean that φ is a unital, positive map. It follows essentially by an application of the Hahn–Banach theorem for archimedean ordered $*$ -vector spaces that the states separate points in E and completely determined the positivity structure; to wit,

Lemma 2.20. *For an abstract operator system E , we have that $x \in E^+$ if and only if $\varphi(x) \geq 0$ for every state φ .*

See [38, Proposition 3.12] for a proof.

Remark 2.21. Even if (E, E^+, e) is an ordered $*$ -vector space with e being an not necessarily archimedean order unit, one may still consider the set of all states on E and define E' to be their closed, linear span. For each $x \in E$ we can define the *evaluation map* $\hat{x} : E' \rightarrow \mathbb{C}$ by $\hat{x}(\varphi) := \varphi(x)$. It can be seen that $(\widehat{E}, \widehat{E}^+, \hat{e})$ is now an archimedean ordered, $*$ -vector space which is known as the *archimedeanization* of E . We refer the reader to [38, Section 3.2] for details.

The crucial point is that e is an archimedean order unit *exactly when* the evaluation map $x \mapsto \hat{x}$ is faithful, that is, has trivial kernel. In many of the constructions going forward we will define some sort of higher-level positivity/order structure on a archimedean ordered $*$ -vector space, making it an operator system, by proscribing the set of higher-order positive maps. At a technical level the positive maps really induce an operator system structure on \widehat{E} which we can identify with E canonically as long as we know the evaluation map is faithful.

Lemma 2.22. *For an abstract operator system E , every positive map $\varphi : E \rightarrow \mathbb{C}$ is completely positive. In particular, every state is completely positive.*

Proof. Let $x \in E_n^+$. We want to show that $\varphi_n(x) \in M_n$ is positive semidefinite. To this end, choose $a \in M_{n,1}$, and observe that $a^* \varphi_n(x) a = \varphi(a^* x a) \geq 0$ since $a^* x a \in E^+$. We are therefore done since $a^* \varphi_n(x) a \geq 0$ for all $a \in M_{n,1}$ characterizes the matrix $\varphi_n(x)$ as being positive semidefinite. \square

Notation 2.23. For (abstract) operator systems E and F , we use $\text{CP}(E, F)$ and $\text{UCP}(E, F)$, respectively, to denote the set of all completely positive maps from E to F and the set of all unital, completely positive maps from E to F .

Lemma 2.24. *Let E and F be abstract operator systems. There is an affine bijection between $\text{CP}(E_n, F)$ and $\text{CP}(E, F_n)$, $\text{CP}(E_n, F) \ni \varphi \leftrightarrow \tilde{\varphi} \in \text{CP}(E, F_n)$, given by*

$$\tilde{\varphi}(x) := [\varphi(e_{ij} \otimes x)]_{ij}, \quad \varphi([x_{ij}]) = \sum_{i,j} \tilde{\varphi}(x_{ij})_{ij},$$

where e_{ij} are the standard matrix units for M_n .

Proof. Let's start with the assumption that $\varphi : E_n \rightarrow F$ is completely positive. It is clear that the map $\Delta_n : M_n \rightarrow M_n \otimes M_n$ given by

$$\Delta_n([x_{ij}]) := \sum_{i,j} e_{ij} \otimes x_{ij} e_{ij}$$

is a (non-unital) $*$ -homomorphism, hence is completely positive. Therefore by Proposition 2.12, we have that for all $x \in E_n^+$ that $\Delta_n \otimes \text{id}_E(x) \in E_n^+$. Since φ is completely positive, so is $\text{id}_{M_n} \otimes \varphi : M_n \otimes E_n \rightarrow F_n$. Let $1_n \in M_{n,1}$ be the matrix of all 1's. We have that the connecting map $\text{ad}(1_n^*) : E \rightarrow E_n$ as defined at the beginning of this section is completely positive. We now compute that

$$\tilde{\varphi}(x) = (\text{id}_{M_n} \otimes \varphi_n) \circ (\Delta_n \otimes \text{id}_E) \circ \text{ad}(1_n^*)(x),$$

so $\tilde{\varphi} : E \rightarrow F_n$ is completely positive.

We now turn to the case that $\tilde{\varphi} : E \rightarrow F_n$ is completely positive, so $\text{id}_{M_n} \otimes \tilde{\varphi} : E_n \rightarrow M_n \otimes F_n$ is as well. We have that

$$\varphi([x_{ij}]) = \text{ad}(1_n) \circ (\Delta_n^\dagger \otimes \text{id}_F) \circ (\text{id}_{M_n} \otimes \tilde{\varphi})([x_{ij}])$$

is completely positive by Exercise 2.13. \square

We now sketch two slightly different proofs of Theorem 2.19.

Sketch of a First Proof of Theorem 2.19. Consider all $\varphi \in \text{UCP}(E, M_n)$, over all $n = 1, 2, \dots$, where we will write $n(\varphi) := n$ for clarity. By Lemma 2.24 we have that $\text{CP}(E, M_n) \cong \text{CP}(E_n, \mathbb{C})$, which by Lemmas 2.22 and 2.20 shows that $\text{UCP}(E, M_n)$ completely determines the positive cone in E_n . For concision let's denote $I := \bigcup_{n=1}^{\infty} \text{UCP}(E, M_n)$.

It is now relatively straightforward to check that setting $H = \bigoplus_{\varphi \in I} \mathbb{C}^{n(\varphi)}$ we have that $\Phi : E \rightarrow \mathcal{B}(H)$ given by

$$\Phi(x) = \bigoplus_{\varphi \in I} \varphi(x) \in \bigoplus_{\varphi \in I} M_{n(\varphi)} \subset \mathcal{B}(H)$$

is a complete order embedding. \square

For the second proof we will take a slightly different tack, essentially following the proof of [36, Theorem 13.1]. This proof will be important later on when we attempt the axiomatization of operator systems as metric structures.

Sketch of a Second Proof of Theorem 2.19. Following the ideas from Proposition 2.5 we begin by defining a semi-norm $\|x\|'_n$ on each E_n by

$$\|x\|'_n := \inf \left\{ t > 0 : \begin{bmatrix} te_n & x \\ x^* & te_n \end{bmatrix} \in E_{2n}^+ \right\},$$

where $e_n = I_n \otimes e$ is the order unit in E_n . The fact that e is an archimedean order unit for E can then be used to show that $\|x\|'_n$ is actually a norm and that E_n^+ forms a closed cone. (We refer the reader to [36, Proposition 13.3] for details.) Once we know that each E_n^+ is closed, setting E'_n to be the norm-closed linear span of $\text{UCP}(E, M_n)$, the Krein–Milman theorem for cones can be used to show, with some work, that the evaluation map $E_n \ni x \mapsto \hat{x} : E' \rightarrow \mathbb{C}$ satisfies $\hat{x}(\varphi) \geq 0$ for all $\varphi \in \text{UCP}(E, M_n)$ if and only if $x \in E_n^+$. Borrowing the notation from the sketch of the first proof,

$$\Phi(x) = \bigoplus_{\varphi \in I} \varphi(x) \in \bigoplus_{\varphi \in I} M_{n(\varphi)} \subset \mathcal{B}(H)$$

is a unital complete order embedding. It follows from Corollary 2.9 that Φ is a complete isometric embedding. \square

One of the main upshots of the Choi–Effros representation theorem from our vantage is that it describes the following “dual” way of viewing an operator system structure via a coherent collection of admissible positive maps into each matrix algebra. Let $(E, E^+, 1)$ be a real ordered $*$ -vector space with archimedean order unit, and let $\mathcal{S}(E)$ denote the set of states of E .

Proposition 2.25. *Suppose that for each $n = 1, 2, \dots$ we define a closed set \mathcal{S}_n of unital, linear maps from E to M_n so that $\mathcal{S}_1 = \mathcal{S}(E)$ and for each $f \in \mathcal{S}_n$ and $\varphi \in \text{UCP}(M_n, M_k)$ we have that $\varphi \circ f \in \mathcal{S}_k$. Then the collection (\mathcal{S}_n) completely determines an operator system structure on E so that $\mathcal{S}_n = \text{UCP}(E, M_n)$ for all n .*

Sketch of Proof. For each $\varphi \in \mathcal{S}_n \subset \mathcal{L}(E, M_n)$, let $\varphi(x) = [\varphi_{ij}(x)]$. Define $E_n^+ \subset E_n$ to be all $x \in E_n$ so that $\sum_{i,j} \varphi_{ij}(x_{ij}) \geq 0$ for all $\varphi \in \mathcal{S}_n$. We leave it as an exercise that this defines a compatible family of real ordered structures on each E_n satisfying Definition 2.17. \square

Perhaps the most important basic result in the theory of operator systems is Arveson's extension theorem.

Theorem 2.26 (Arveson's Extension Theorem). *For every unital inclusion $E \subset F$ of operator systems and every unital, completely positive map $\varphi : E \rightarrow \mathcal{B}(H)$, there exists a unital, completely positive map $\tilde{\varphi} : F \rightarrow \mathcal{B}(H)$ extending φ .*

We will outline a proof here: for a detailed proof see [6, Theorem 1.6.1] or [36, Theorem 7.5]

Sketch of Proof. We first note that any positive map $\varphi : E \rightarrow \mathbb{C}$ extends to a positive (hence, completely positive) map $\tilde{\varphi} : F \rightarrow \mathbb{C}$ by the Hahn–Banach extension theorem for ordered vector spaces [13, Corollary 9.12]. We now consider the case when $\dim(H) = n < \infty$. We have that $\text{CP}(E, \mathcal{B}(H)) \cong \text{CP}(E, M_n) \cong \text{CP}(M_n(E), \mathbb{C})$. As before, we can now extend any $\varphi \in \text{CP}(M_n(E), \mathbb{C})$ to $\tilde{\varphi} \in \text{CP}(M_n(F), \mathbb{C}) \cong \text{CP}(F, M_n)$. Since $\tilde{\varphi}$ is unital and (2-)positive, it is contractive by Exercise 2.6.

Now let H be arbitrary and let $(H_i)_{i \in I}$ be the directed set of all finite-dimensional subspaces of H , ordered by inclusion. Let $p_i : H \rightarrow H_i$ be the orthogonal projection onto H_i and consider the unital completely positive maps $\varphi_i : E \rightarrow \mathcal{B}(H_i)$ defined by $\varphi_i(x) = p_i \varphi(x) p_i$. We may extend each to $\tilde{\varphi}_i : F \rightarrow \mathcal{B}(H_i)$ unittally and completely positively. Since all maps are contractive, we can now take a pointwise-ultraweak cluster point of the net $\tilde{\varphi}_i$ to obtain a unital, completely positive map $\tilde{\varphi} : F \rightarrow \mathcal{B}(H)$. It is straightforward to check that $\tilde{\varphi}$ extends φ . \square

We now turn our attention to describing the dual of an operator system. We begin with the following proposition.

Lemma 2.27. *Let E be an operator system. Every bounded linear functional $\varphi : E \rightarrow \mathbb{C}$ is a linear combination of four states. Every completely bounded map $\varphi : E \rightarrow M_n$ is a linear combination of four completely positive maps.*

Proof. Suppose $E \subset \mathcal{B}(H)$. By the Hahn-Banach theorem every bounded linear contraction $\varphi : E \rightarrow \mathbb{C}$ extends to a linear contraction $\tilde{\varphi} : \mathcal{B}(H) \rightarrow \mathbb{C}$. Now $\tilde{\varphi}$ is a linear combination of four states by the Jordan decomposition theorem for C^* -algebras [44, Proposition 2.1]. The rest of result follows from this by essentially the same proof as Lemma 2.24 by noting the correspondence described therein converts any bounded linear map $\varphi : E \rightarrow M_n$ to a bounded linear functional $\tilde{\varphi} : M_n(E) \rightarrow \mathbb{C}$. \square

In this we see that E^* is equipped with a matricial ordered $*$ -vector space structure, where the positive cone in $M_n(E^*)$ is identified with the completely positive maps $E \rightarrow M_n$. This does not give E^* the structure of an operator system, however, as there is no clear choice of an archimedean order unit. In the case that E a finite-dimensional operator system there is a way of placing an operator system structure on E^* which is non-canonical as it will depend on choosing an order unit. This is essentially a consequence of the following lemma: see [26, Lemma 2.5] for a proof.

Lemma 2.28. *For any finite-dimensional operator system E we can find a basis $\{x_0 = 1, x_1, \dots, x_n\}$ of self-adjoint elements so that the dual basis $\{x_0^*, x_1^*, \dots, x_n^*\}$, defined by $x_i^*(x_j) = \delta_{ij}$, is a hermitian basis for E^* with δ_0 an order unit.*

A related, useful lemma appears as [6, Lemma B.10].

Lemma 2.29. *For any finite-dimensional operator system E , we can find a spanning set $1 = x_1, \dots, x_n$ consisting of hermitian elements of unit norm so that the dual basis x_1^*, \dots, x_n^* defined by $x_i^*(x_j) = \delta_{ij}$ satisfies $\|x_i^*\| = 1$ as well for all $i = 1, \dots, n$. Moreover, x_1^* may be taken to be an order unit in the cone $\mathcal{S}(E)$.*

Even though E^* is not an operator system, E^{**} is, as it turns out that the evaluation map $\hat{1} : E^* \rightarrow \mathbb{C}$ is an archimedean order unit for the matricial ordered structure on E^{**} induced from E^* . In fact, the following is true.

Proposition 2.30. *The natural embedding $E \hookrightarrow E^{**}$ given by sending $x \mapsto \hat{x}$ is a complete order embedding of operator systems. If E is finite-dimensional, then it is a complete order isomorphism.*

Proof. We have that $E \subset \mathcal{B}(H)$ for some Hilbert space H . By [44, Section III.2] there is a unital $*$ -algebra embedding $\pi : \mathcal{B}(H) \rightarrow \mathcal{B}(K)$ so that there is a unique extension $\pi' : \mathcal{B}(H)^{**} \rightarrow \mathcal{B}(K)$ which is a homeomorphism from the weak*-topology on $\mathcal{B}(H)$ to the ultraweak topology on $\mathcal{B}(K)$. This implies that $x \in M_n(\mathcal{B}(H)^{**})$ is positive in the matricial order structure defined on $\mathcal{B}(H)^{**}$ if and only if $\pi'(x)$ is positive in $M_n(\mathcal{B}(K))$ as both are determined from taking the closures of $M_n(\mathcal{B}(H))^+$ in the respective weak topologies. Since $1 = \pi(1) \in \mathcal{B}(K)$ is archimedean, we have that $\hat{1} \in E^{**}$ is archimedean, thus E^{**} is an operator system. \square

Definition 2.31. For an operator system E , a subspace $J \subset E$ is said to be a *kernel* if $J = \bigcap_{i \in I} \ker(\varphi_i)$ where $\{\varphi_i : i \in I\}$ is a collection of states on E .

Given a kernel $J \subset E$, we may put a *quotient operator system* structure on $\widehat{E/J}$ for the ordered $*$ -vector space quotient E/J by identifying $\text{UCP}(E/J, M_n)$ with the weak*-closed, convex subset $\text{UCP}(E, M_n; J)$ of $\text{UCP}(E, M_n)$ consisting of all φ with $J \subset \ker(\varphi)$. Note that in general the quotient structure must be placed on $\widehat{E/J}$ and not E/J itself as the class of 1 in E/J is an order unit which is possibly not archimedean, so we must pass through the evaluation map to obtain the archimedeanization. Quotients of operator systems were first defined by Choi and Effros [9]. We refer the reader to [26, 29] for an in-depth treatment of quotients of operator systems.

2.1. Examples of Operator Systems.

Example 2.32. Every unital C^* -algebra A is canonically an operator system where A_n^+ is just the cone of positive elements in the C^* -algebra $M_n(A)$. In particular the matrix algebra M_k is naturally an operator system under $(M_k)_n^+ = (M_n \otimes M_k)^+$. Note that not every element of $(M_n \otimes M_k)^+$ can be written as a sum of simple tensors $x \otimes y$ with $x \in M_n^+$ and $y \in M_k^+$ as soon as $n, k \geq 2!$ In quantum information theory the elements in $(M_n \otimes M_k)^+$ which are not in the span of simple tensors of positive elements are said to be *entangled*.

Example 2.33. Let X be a finite set. For a vector space V , let $V[X]$ denote the vector space of all functions $f : X \rightarrow V$. We have that $\mathbb{C}[X]$ is a real, ordered $*$ -vector space under pointwise conjugation and the cone $\mathbb{R}_{\geq 0}[X]$ of all non-negatively-valued functions $f : X \rightarrow \mathbb{R}_{\geq 0}$, with order unit being the characteristic function 1_X .

We can define the following “free” canonical operator system structure on $\mathbb{C}[X]$ as follows. We have that $M_n(\mathbb{C}[X]) \cong M_n[X]$ canonically, so we can define $\mathbb{C}[X]_n^+$ as the cone of all functions $f : X \rightarrow M_n^+$. In the dual picture, consider a map $g : X \rightarrow M_k^+$, which extends to a positive linear map $\tilde{g} : \mathbb{C}[X] \rightarrow M_k$. We see that $\tilde{g}_n : M_n[X] \rightarrow M_n(M_k) \cong M_n \otimes M_k$ is given by $\tilde{g}_n(f) = \sum_{x \in X} f(x) \otimes g(x)$; hence, every positive map from $\mathbb{C}[X]$ to M_k is completely positive. Conversely, consider the a positive map $g : \mathbb{X} \rightarrow M_n^+$. For $f \in M_n[X]$ we have the pairing $\sum_{i,j} \tilde{g}_{ij}(f_{ij})$ as given in the proof of Proposition 2.25 can be computed as

$$\sum_{i,j} \tilde{g}_{ij}(f_{ij}) = \sum_{x \in X} \text{tr}(f(x)g(x)).$$

Since $A \in M_n$ is positive semidefinite if and only if $\text{tr}(AB) \geq 0$ for all B positive semidefinite, it follows that $\sum_{i,j} \tilde{g}_{ij}(f_{ij}) \geq 0$ for all $g : X \rightarrow M_n^+$ if and only if $f(x) \in M_n^+$ for all $x \in X$.

In summary, we see that this operator system structure on $\mathbb{C}[X]$ has the following two universal properties: for every operator system E every positive map $\varphi : E \rightarrow \mathbb{C}[X]$ is completely positive and every function $f : X \rightarrow E^+$ induces a completely positive map $\tilde{f} : \mathbb{C}[X] \rightarrow E$.

Example 2.34. We now describe the construction of universal operator systems from sets of linear relations. We equip $\mathbb{C}[X]$ with the usual inner product structure $\langle f, g \rangle := \sum_{x \in X} f(x)\bar{g}(x)$. Consider a set of elements $r_1, \dots, r_k \in \mathbb{R}[X]$ so that $\langle r_i, 1_X \rangle = \sum_x r_i(x) = 0$ for all $i = 1, \dots, k$. Let $R = \text{span}\{r_1, \dots, r_k\}$. We define an operator system structure on the quotient space $\mathbb{C}[X]/R$ by defining the completely positive maps from $\mathbb{C}[X]/R$ to M_n to be exactly the set of linear maps induced by functions $f : X \rightarrow M_n^+$ so that $\sum_{x \in X} r_i(x)f(x) = 0$ for all $i = 1, \dots, k$. Since $1_X \in R^\perp$ we have that the constant function $f(x) \equiv \frac{1}{|X|}I_n$ is a unital, completely positive map, so that $\text{UCP}(\mathbb{C}[X]/R, M_n)$ is nonempty for each n .

This is a special case of the quotienting construction detailed above, as it is easy to check that $R \subset \mathbb{C}[X]$ is a kernel. This construction seems rather simple, but this belies

a very rich structure. For example, let us take $|X| = 2n$ and label the elements of X as $s_1, \dots, s_n, t_1, \dots, t_n$. Consider the single element r defined by $r(s_i) = 1$ and $r(t_j) = -1$ for all $i, j = 1, \dots, n$. It is a result of Kavruk [27] that in this case that $\mathbb{C}[X]/R$ is canonically completely order isomorphic the operator subsystem of $C^*(\mathbb{Z}/n\mathbb{Z} * \mathbb{Z}/n\mathbb{Z})$ spanned by the generating set $(\mathbb{Z}/n\mathbb{Z} * 1) \cup (1 * \mathbb{Z}/n\mathbb{Z})$.

Example 2.35. Let $P \subset \mathbb{R}^k$ be a closed cone with archimedean order unit e . We can define an operator system structure on $V = (\mathbb{C}^n, P, e)$ by declaring $\text{UCP}(V : M_n)$ to be the set of all linear maps $f : \mathbb{C}^n \rightarrow M_n$ so that $f(\mathbb{R}^n) \subset M_n^h$, $f(e) = I_n$, and $f(p) \in M_n^+$ for all $p \in P$. This is the *maximal operator system structure* $\text{MAX}(V)$ over V as defined in [37]: the following is proved therein.

Proposition 2.36. *The operator system $\text{MAX}(V)$ is characterized by the universal property that $\varphi : \text{MAX}(V) \rightarrow \mathcal{B}(H)$ is completely positive if and only if $\varphi(P) \subset \mathcal{B}(H)^+$.*

We record this property by introducing new terminology.

Definition 2.37. We say that an operator system E is *k-maximal* if every k -positive map $\varphi : E \rightarrow \mathcal{B}(H)$ is completely positive. For ease of reference, we will say that E is *maximal* if it is 1-maximal.

Remark 2.38. If $K \subset \mathbb{R}^\ell$ is compact, convex set with 0 as an interior point, we can form the cone $P_K \subset \mathbb{R}^{\ell+1}$ to consist of all vectors of the form $te - tv$ where $e = (1, 0, \dots, 0)$, $v \in 0 \oplus K$, and $t \geq 0$. It can be checked that e is an archimedean order unit for P_K and $P_K - P_K = \mathbb{R}^\ell$.

Proposition 2.39. *For the maximal operator system structure on $V_K := (\mathbb{C}^{\ell+1}, P_K, e)$, we have that $\text{UCP}(V_K, M_n)$ is parametrized by the set of all ℓ -tuples X_1, \dots, X_ℓ of hermitian elements in M_n satisfying the linear matrix inequalities $I_n \succeq v_1 X_1 + \dots + v_\ell X_\ell$ for all $v \in K$.*

We leave the proof as an exercise to the reader. Conversely, for a closed cone P with archimedean order unit e , consider the “slice” $K = \{p \in P : \langle p, e \rangle = 1\}$. It can be shown that P_K is affinely isomorphic to P , so that all maximal operator systems essentially arise in this way. This is essentially a special case of the Webster–Winkler Duality Theorem [45].

Example 2.40. Again, let $P \subset \mathbb{R}^k$ be a closed cone with archimedean order unit e . We can define an operator system structure on $V = (\mathbb{C}^n, P, e)$ by embedding V via the evaluation map into the algebra $C(\mathcal{S}(V))$ of continuous functions on the set $\mathcal{S}(V)$ of all states, which is a closed and bounded, hence compact, subset of \mathbb{C}^n . This is the *minimal operator system structure* $\text{MIN}(V)$ defined on V as described in [37]. The following is proved therein.

Proposition 2.41. *The minimal operator system structure on V is characterized by the universal property that for every operator system E every positive map $\varphi : E \rightarrow V$ extends to a completely positive map $\varphi : E \rightarrow \text{MIN}(V)$.*

3. MODEL THEORY OF OPERATOR SYSTEMS

A linear matrix $*$ -polynomial in n -variables $x = (x_1, \dots, x_n)$ is an expression of the form

$$p(x) = a_1 \otimes x_1 + \dots + a_n \otimes x_n + b_1 \otimes x_1^* + \dots + b_n \otimes x_n^* + c \otimes 1$$

where $(a_1, \dots, a_n, b_1, \dots, b_n, c)$ are elements in M_k for some k . Equivalently, we may think of $p(x)$ as a $k \times k$ matrix where each entry is a linear $*$ -polynomial in $1, x_1, \dots, x_n$.

Definition 3.1. We say that a linear matrix $*$ -polynomial is *hermitian* if $p(x^*)^* = p(x)$ which is to say that c is hermitian and $a_i^* = b_i$ for all $i = 1, \dots, n$. A linear matrix $*$ -polynomial is said to be *homogeneous* if $p(0) = 0$, that is, if $c = 0$. We say that $p(x)$ has *degree* k if $a_1, \dots, a_n, b_1, \dots, b_n, c \in M_k$, and we write $\deg(p) = k$.

We define a *linear matrix inequality* to be an expression of the form $p(x) \succeq 0$ where $p(x)$ is a hermitian linear matrix $*$ -polynomial. Note that given a system $p_1(x) \succeq 0, \dots, p_m(x) \succeq 0$ of linear matrix inequalities we may without loss of generality combine them into a single linear linear matrix inequality via a block diagonal embedding of the matrix coefficients of each p_i into a larger matrix. Many problems in functional analysis can be essentially phrased as the following general problem:

Problem 3.2 (Matrix Completion Problem). Does there exist $X = (X_1, \dots, X_n) \in \mathcal{D}^n$ for some domain $\mathcal{D} \subset \mathcal{B}(H)$ so that $p(X) \succeq 0$? (Here 1 is interpreted as the unit in $\mathcal{B}(H)$.)

From the perspective of logic, the Matrix Completion Problem can be seen as whether some existential sentence is true over some (hopefully) suitable domain of quantification \mathcal{D} . Notice that by Proposition 2.5, this framework captures problems involving norm estimates as well: if we are interested in the statement

$$\exists x \in \mathcal{D}^n : \|p(x)\| \leq 1,$$

then setting $p^*(x) := p(x^*)^*$ we may write this equivalently as

$$\exists x \in \mathcal{D}^n : \begin{bmatrix} 1 & p(x) \\ p^*(x) & 1 \end{bmatrix} \succeq 0.$$

If H is a finite-dimensional Hilbert space, the Matrix Completion Problem frequently occurs in the context of *semidefinite programming* in terms of whether a given semidefinite program is feasible. (We will not go any further into this here, but we refer the reader to [19, 34] for an introduction to modern developments in the theory of semidefinite programming.)

3.1. Building the Language. One could say that the broad goal of building a model theory for matrix operator systems is to devise a general framework where problems such as the Matrix Completion Problem can be systematically studied. Following [20, Appendix B] we give a description of how to axiomatize operator systems in the context of first order continuous logic for metric structures. (We refer the reader to [15, Section 2.1] for the basics of first order continuous logic for metric structures.) In terms of building the language, this immediately brings to attention the following considerations.

- (1) We will need a collection of sorts E_1, E_2, E_3, \dots intended to capture the operator system $E = E_1$ as well as its matrix amplifications so that E_n should be interpreted as $M_n(E)$ when our work is done. Each of these sorts will need domains of quantification $\mathcal{D}_r(E_n)$ corresponding to the norm r -balls about the origin. Since we are about to give each E_n a real vector space structure, it will suffice to consider $\mathcal{D}_1(E_n)$ alone.
- (2) We will need another collection of sorts C_1, C_2, C_3, \dots for which each C_n will need to be interpreted as the cone of positive elements E_n^+ in E_n with domains of quantification to be interpreted as the restriction of the r -balls in E_n to C_n .
- (3) Sorts $M_{n,k}$ for the complex $n \times k$ matrices with each domains of quantification the operator norm r -balls.
- (4) Sorts for \mathbb{C} , \mathbb{R} , and the non-negative reals $\mathbb{R}_{\geq 0}$ with the domains of quantification being the standard r -balls.

For each of these sorts we will need the corresponding relational symbols.

- (1) We will need constant symbols $0, 1$ in $\mathcal{D}_1(E_1)$ for 0 and the order unit. We require function symbols

$$\cdot_n : \mathbb{C} \times E_n \rightarrow E_n, \quad +_n : E_n \times E_n \rightarrow E_n, \quad \text{and} \quad *_n : E_n \rightarrow E_n$$

to be used for scalar multiplication, addition, and involution, respectively. Additionally one further set of function symbols

$$f_{n,k} : M_{n,k} \times E_n \rightarrow E_k$$

is needed for $(a, x) \mapsto a^*xa$ and another

$$p_n^{ij} : E_n \rightarrow E_1$$

for projections giving the matrix coordinates. There are predicate symbols

$$\|\cdot\|_n : E_n \rightarrow \mathbb{R}_{\geq 0}$$

which we will want to interpret as the norms induced from the matrix ordering as in Proposition 2.5.

- (2) We define constant symbols $0', 1'$ in $\mathcal{D}_1(C_1)$ and function symbols

$$\cdot'_n : \mathbb{R}_{\geq 0} \times C_n \rightarrow C_n, \quad +'_n : C_n \times C_n \rightarrow C_n, \quad \text{and} \quad g_{n,k} : M_{n,k} \times C_n \rightarrow C_k$$

to be used for scalar multiplication, addition, and $(a, x) \mapsto a^*xa$, respectively.

- (3) Similarly to the first two items, there will need to constant symbols for 0 and function symbols for scalar multiplication, addition, and taking adjoints $*$: $M_{n,k} \rightarrow M_{k,n}$ for each $M_{n,k}$.
- (4) Again, we must define symbols for $0, 1$ in $\mathbb{R}_{\geq 0}$, \mathbb{R} , and \mathbb{C} with an additional symbol i in \mathbb{C} , all in the unit domain of quantification. Function symbols are needed for addition, scalar multiplication, and, in the case of \mathbb{C} , conjugation.
- (5) Besides these we need a couple of other sets of function symbols

$$i_n : C_n \rightarrow E_n,$$

obviously for the inclusion of the positive cone in each E_n and

$$h_n : E_n \rightarrow E_{2n}$$

which we want to interpret as

$$h_n : x \mapsto \begin{bmatrix} 1_n & x \\ x^* & 1_n \end{bmatrix}$$

to be used to guarantee our predicate symbols are interpreted as intended.

Pertaining to the use of the h_n 's we require the following result.

Lemma 3.3. *For an operator system E and $x \in E_n$ we have that*

$$\|x\|_n \dot{\div} 1 = \text{dist} \left(\begin{bmatrix} 1 & x \\ x^* & 1 \end{bmatrix}, E_{2n}^+ \right).$$

Here $x \dot{\div} y := \max\{x - y, 0\}$.

Proof. We may assume without loss of generality that $n = 1$ and $r := \|x\| > 1$. By Proposition 2.5 we have that

$$\begin{bmatrix} 1 & \frac{1}{r}x \\ \frac{1}{r}x^* & 1 \end{bmatrix} \in E_2^+;$$

hence,

$$h(x) := \text{dist} \left(\begin{bmatrix} 1 & x \\ x^* & 1 \end{bmatrix}, E_2^+ \right) = \text{dist} \left(\begin{bmatrix} 0 & (1 - 1/r)x \\ (1 - 1/r)x^* & 0 \end{bmatrix}, E_2^+ \right).$$

Noting that

$$\left\| \begin{bmatrix} a & b \\ b^* & c \end{bmatrix} \right\| = \left\| \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a & b \\ b^* & c \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \right\| = \left\| \begin{bmatrix} a & -b \\ -b^* & c \end{bmatrix} \right\|,$$

we see by simple arithmetic that

$$\left\| \begin{bmatrix} 0 & y \\ y^* & 0 \end{bmatrix} - \begin{bmatrix} a & b \\ b^* & c \end{bmatrix} \right\| = \left\| \begin{bmatrix} 0 & y \\ y^* & 0 \end{bmatrix} - \begin{bmatrix} a & -b \\ -b^* & c \end{bmatrix} \right\|.$$

Hence, by averaging the two expressions and using the triangle inequality we see that $h(x)$ must be approached by elements in E_2^+ of the form $\begin{bmatrix} a & 0 \\ 0 & c \end{bmatrix}$. However, by the same argument as before

$$\left\| \begin{bmatrix} 0 & y \\ y^* & 0 \end{bmatrix} - \begin{bmatrix} a & 0 \\ 0 & c \end{bmatrix} \right\| = \left\| \begin{bmatrix} 0 & -y \\ -y^* & 0 \end{bmatrix} - \begin{bmatrix} a & 0 \\ 0 & c \end{bmatrix} \right\|$$

which shows by averaging that $a = c = 0$ is optimal. Therefore

$$h(x) = \left\| \begin{bmatrix} 0 & (1 - 1/r)x \\ (1 - 1/r)x^* & 0 \end{bmatrix} \right\|_2 = \left(1 - \frac{1}{\|x\|}\right) \|x\|$$

since

$$\begin{bmatrix} x & 0 \\ 0 & x^* \end{bmatrix} = \begin{bmatrix} 0 & x \\ x^* & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

□

We require one further technical result: see [20, Lemma B.1].

Exercise 3.4. For an operator system E and $x \in E_n^+$ we have that $\text{dist}(-x, E_n^+) = \|x\|_n$.

To complete the language we need axioms that tell us that the symbols enumerated above are interpreted as they should be. These would include, for instance, axioms stating that each E_n has the structure of a $*$ -vector space, E_n isomorphic to $M_n(E_1)$, and the $*$ -vector space structure on E_n is the natural one induced from the $*$ -vector space structure on E_1 via matrix amplification. In addition to these, we need the following, more specialized, axioms.

- (1) There is an axiom that each $\|\cdot\|_n$ is seminorm on E_n , and that this predicate is the metric predicate, that is,

$$\sup_{x,y \in \mathcal{D}_1(E_n)} |\text{dist}(x,y) - \|x-y\|_n| = 0.$$

This implies that each $\|\cdot\|_n$ is, in fact, a norm.

- (2) We need an axiom stating that $i_n : C_n \rightarrow E_n$ is an isometric inclusion, an axiom for ensuring that the range of i_n lies in the hermitian part of E_n ,

$$\sup_{x \in \mathcal{D}_1(C_n)} \text{dist}(i_n(x), i_n(x)^*) = 0,$$

and an axiom (by way of Exercise 3.4) ensuring that $-i_n(C_n) \cap i_n(C_n) = \{0\}$,

$$\sup_{x,y \in \mathcal{D}_1(C_n)} |\|x\|_n - \text{dist}(i_n(x), -i_n(y))| = 0.$$

- (3) Finally, there is an axiom stating the content of Lemma 3.3, that is,

$$\sup_{x \in \mathcal{D}_r(E_n)} |(\|x\|_n - 1) - \text{dist}(h_n(x), C_{2n})| = 0.$$

Remark 3.5. The last axiom is important as it guarantees that 1 is interpreted as a (complete) order unit. To see this, let $x \in E_n$ with $x = x^*$ and set $r := \|x\|_n$. We have that $\|\frac{1}{r}x\|_n = 1$, thus

$$h_n\left(\frac{1}{r}x\right) = \begin{bmatrix} 1_n & \frac{1}{r}x \\ \frac{1}{r}x & 1_n \end{bmatrix} \in C_{2n}.$$

We then have that $v^*h_n(x)v = 1_n + \frac{1}{r}x \in C_n$ where $v = [\frac{1}{\sqrt{2}}I_n, \frac{1}{\sqrt{2}}I_n]^t$.

Remark 3.6. As a consequence of previous remark we have that $i_n(C_n)$ spans E_n for all n . Indeed, for all $x \in E_n$ with $\|x\|_n \leq 1$ we have that

$$x = ((1_n + (x + x^*)/2) - 1_n) + \sqrt{-1}((1_n - \sqrt{-1}(x - x^*)/2) - 1_n)$$

which expresses x as a linear combination of four elements in the positive cone, each with norm at most 2.

So far the axiomatization says that any model can be interpreted as a matrix ordered $*$ -vector space with order unit (see Remark 2.18), the only difference from being an abstract operator system then is the issue of whether 1 is an *archimedean* order unit.

To bridge this last gap, we will rely crucially on the fact that the metric structure gives rise to a norm $\|\cdot\|_n$ on each E_n .

Lemma 3.7. *Let E be an matrix ordered $*$ -vector space with order unit 1. Setting*

$$\|x\|'_n := \inf \left\{ t > 0 : \begin{bmatrix} t1_n & x \\ x^* & t1_n \end{bmatrix} \in E_{2n}^+ \right\},$$

we have that that E is an abstract operator system if and only if one of the following holds:

- (1) 1 is an archimedean order unit for E ;
- (2) 1_n is an archimedean order unit for E_n for all n ;
- (3) $\|\cdot\|'_n$ is a norm for all n .
- (4) $\|\cdot\|'_1$ is norm.

Proof. (1 \Leftrightarrow 2) Suppose that 1_n is not archimedean for some n , that is, there exists a non-zero $x \in E_n^h$ so that $r1_n \pm x \in E_n^+$ for all $r > 0$. We must have that $y = v^*xv \neq 0$ for some unit vector $v \in \mathbb{R}^n$. Seeing that $r1 \pm y = v^*(r1_n \pm x)v \in E_1^+$, we verify that 1 is not archimedean. The converse is trivial.

(2 \Rightarrow 3) We leave it as an exercise to check that $\|\cdot\|'_n$ is always a seminorm, that is, $\|x\|'_n = \|-x\|'_n$, $\|\lambda x\|'_n = |\lambda| \|x\|'_n$, and $\|x + y\|'_n \leq \|x\|'_n + \|y\|'_n$ for all $\lambda \in \mathbb{C}$ and $x, y \in E_n$. For a nonzero $x \in E_n$, we have that

$$y := \begin{bmatrix} 0 & x \\ x^* & 0 \end{bmatrix} \in E_{2n}^h;$$

thus, there is a $t > 0$ so that for all $s < t$ it is not the case that $s1_{2n} \pm y \in E_{2n}^+$. It follows that $\|x\|'_n = t$.

(3 \Rightarrow 2) We can assume without loss of generality that 1 is not archimedean as witnessed by $x \in E^h$. Now since $x + r1 \in E^+$,

$$\begin{bmatrix} x + r1 & x + r1 \\ x + r1 & x + r1 \end{bmatrix} = [1, 1]^t (x + r1) [1, 1] \in E_2^+$$

and

$$\begin{bmatrix} (2r)1 & x + r1 \\ x + r1 & (2r)1 \end{bmatrix} - \begin{bmatrix} x + r1 & x + r1 \\ x + r1 & x + r1 \end{bmatrix} = \begin{bmatrix} r1 - x & 0 \\ 0 & r1 - x \end{bmatrix} \in E_2^+.$$

Since $\|x + r1\|'_1 - \|x\|'_1 \leq \|r1\|'_1 \leq r$, we conclude that $\|x\|'_1 = 0$, so $\|\cdot\|'_1$ is not a norm.

Finally, (4 \Rightarrow 3) is trivial and (1 \Rightarrow 4) is a special case of (2 \Rightarrow 3). \square

With this lemma now in hand, we see that the axiomatization of abstract operator systems as metric structures is complete. We note in passing that by the Choi-Effros representation theorem for an abstract operator system E each E_n is metrically complete in the $\|\cdot\|'_n$ -norm, so there was nothing lost in insisting upon this in the metric structure language, though nothing is gained by doing so.

3.2. Formulas, Theories, and Models. Now that the language is built we briefly discuss the basic aspects of the continuous model theory of operator systems. We refer the reader to [15, Chapter 2] and [5] for a more in-depth treatment.

Let \mathcal{L} be a language for metric structures as described in [15, Section 2.1]. To each domain of quantification in each sort we assign an infinite number of variables.

Definition 3.8.

- (1) An *atomic formula* $\varphi(x_1, \dots, x_n)$ is an expression in the language, built using finitely many variables and function symbols, which terminates with the application of a predicate. (We will assume all predicates take values in \mathbb{R}^+ .) Since each variable x_i has an assigned domain of quantification \mathcal{D}_i , we have that f has domain

$$\text{domain}(\varphi) := \mathcal{D}_1 \times \dots \times \mathcal{D}_n.$$

Since each variable comes with a (bounded) domain of quantification and each function symbol has a modulus of uniform continuity we can assign a bounded range $[0, K] \subset \mathbb{R}^+$ to φ .

- (2) A *connective* is a uniformly continuous function $f : (\mathbb{R}^+)^k \rightarrow \mathbb{R}^+$.
- (3) If $\varphi_1, \dots, \varphi_k$ are atomic formulas over a common set of variables $x = (x_1, \dots, x_n)$, we can define a *quantifier-free formula* as an expression of the form $g(x) = f(\varphi_1(x), \dots, \varphi_k(x))$ where f is a connective.
- (4) A *formula* $h(x)$ is an expression of the form

$$h(x) = Q_\ell \cdots Q_1 g(x_1, \dots, x_n)$$

where $\ell \leq n$, g is a quantifier-free formula, and Q_i is either the *existential quantifier* $\inf_{x_i \in \mathcal{D}_i}$ or the *universal quantifier* $\sup_{x_i \in \mathcal{D}_i}$. A formula is a *sentence* if all variables are quantified, that is, if $\ell = n$. Similarly to atomic formulas, each formula has an associated domain and bounded range.

- (5) To each formula $h(x) = h(x_1, \dots, x_n)$ and each metric \mathcal{L} -structure M , we can associate an *interpretation* $h(x)^M$ of h which is the function from $\mathcal{D}_1(M) \times \dots \times \mathcal{D}_n(M)$ to \mathbb{R}^+ determined by interpreting all variables, function symbols, domains of quantification, etc., in the language \mathcal{L} in M . If h has range $[0, K]$, then this means that $h(x)^M \in [0, K]$ for all \mathcal{L} -structures M and all $x_1, \dots, x_n \in M$.

Example 3.9. Let \mathcal{L} be the language of operator systems described above. As we will explain below in Remark 3.26 the sorts corresponding to the positive cones do not add any expressive power in terms of defining formulas, so it suffices to build formulas only using the sorts (E_n) . Since the matrix entries of any variable in E_n are interpreted as being in the sort $E = E_1$, it suffices to only consider formulas with variables in this sort alone. Thus, in the language of operator systems, atomic formulas are effectively expressions of the form $\|p(x)\|_d$ where $p(x)$ is a linear matrix $*$ -polynomial of degree d , where each variable is restricted to a domain of quantification in E_1 . It is a bit annoying to have to deal with the domains of quantification at this level, rather than just assigning each variable to the (unbounded) sort E_1 . We will take this view, so each $\|p(x)\|_d$ is technically what we will term an *unbounded atomic formula*.

Definition 3.10. Let \mathcal{L} be a language for metric structures.

- (1) Any collection T of \mathcal{L} -sentences is called a *theory*.
- (2) An \mathcal{L} -structure M *models* T , written $M \models T$, if $h^M = 0$ for all sentences $h \in T$.
- (3) A theory T is said to be *consistent* if there is an \mathcal{L} -structure M so that $M \models T$.
- (4) The *theory* of an \mathcal{L} -structure M , denoted $\text{Theory}(M)$ is the collection of all \mathcal{L} -sentences h so that $h^M = 0$. Clearly, $M \models \text{Theory}(M)$.
- (5) We write $\text{Model}(T)$ for the class of all models of a theory. We say that a class \mathcal{C} of \mathcal{L} -structures is *elementary* if $\mathcal{C} = \text{Model}(T)$ for some theory T .

We will discuss more about elementary classes at the end of the next section.

3.3. Ultraproducts. Let I be an arbitrary set, and let $(E_i)_{i \in I}$ be a collection of (concrete) operator systems indexed by I . We define the *direct product* $\prod_{i \in I} E_i$ to be the set of all bounded functions $x : I \rightarrow \bigsqcup_{i \in I} E_i$, the disjoint union, with $x_i \in E_i$ for all $i \in I$. This is a Banach space $*$ -vector space under pointwise addition, scalar multiplication, and involution and norm $\|x\| = \sup_{i \in I} \|x_i\|_{E_i}$. For representations $E_i \subset B(H_i)$, we see that $\prod_{i \in I} E_i$ is a concrete operator system isometrically represented on the Hilbert space direct sum $\bigoplus_{i \in I} H_i$ via

$$x(\bigoplus_{i \in I} \xi_i) := \bigoplus_{i \in I} x_i \xi_i.$$

It is left as an exercise to check that under this concrete representation we have that $(\prod_{i \in I} E_i)^+ = \prod_{i \in I} E_i^+$ with unit $1 = (1_i)_{i \in I}$ where $1_i \in E_i$ is the unit.

For the rest of the section I will be a fixed directed set and \mathcal{U} be an ultrafilter on I . Our task will be to give a “concrete” definition of the ultraproduct of the operator systems $(E_i)_{i \in I}$ and then explain how this ultraproduct is the ultraproduct at the level of the language of operator systems that we have developed. Consider an arbitrary collection $\phi = (\phi_i)_{i \in I}$ of matrix states $\phi_i \in \mathcal{S}_n(E_i)$. Each such ϕ induces a matrix state

$$\phi_{\mathcal{U}} : \prod_{i \in I} E_i \rightarrow M_n, \text{ defined by } \phi_{\mathcal{U}}(x) := \lim_{\mathcal{U}} \phi_i(x_i).$$

Exercise 3.11. Show that for $\phi = (\phi_i) \in \prod_{i \in I} \mathcal{S}(E_i)$

$$\bigcap_{\phi = (\phi_i)} \ker(\phi_{\mathcal{U}}) = \left\{ (x_i) \in \prod_{i \in I} E_i : \lim_{\mathcal{U}} \|x_i\| = 0 \right\} =: \mathcal{J}.$$

Hence, \mathcal{J} is a kernel.

Exercise 3.12. Show that the class of 1 is an archimedean order unit for the ordered $*$ -vector space quotient; hence, the evaluation map $\hat{\cdot} : \prod_{i \in I} E_i / \mathcal{J} \rightarrow (\prod_{i \in I} E_i / \mathcal{J})^{**}$ is faithful and $\prod_{i \in I} E_i / \mathcal{J}$ itself is equipped with a quotient operator system structure.

Definition 3.13. We define the *ultraproduct*, $\prod_{\mathcal{U}} E_i$ of the operator systems $(E_i)_{i \in I}$ to be the operator system quotient $\prod_{i \in I} E_i / \mathcal{J}$. In other words, $\prod_{\mathcal{U}} E_i$ is the operator system structure defined by the matrix states

$$\text{UCP} \left(\prod_{\mathcal{U}} E_i, M_n \right) = \left\{ \phi_{\mathcal{U}} : \phi \in \prod_{i \in I} \text{UCP}(E_i, M_n) \right\}.$$

To check that this ultraproduct is an ultraproduct of metric structures in the language of operator systems by [5, Chapter 5] it suffices to show that this construction is compatible with the metric ultraproduct for each sort, that is,

$$\mathcal{D}_r \left(\prod_{\mathcal{U}} M_n(E_i) \right) = \prod_{\mathcal{U}} \mathcal{D}_r(M_n(E_i)), \quad \mathcal{D}_r \left((M_n(\prod_{\mathcal{U}} E_i))^+ \right) = \prod_{\mathcal{U}} \mathcal{D}_r(M_n(E_i)^+)$$

for all $r > 0$ and $n = 1, 2, \dots$. In fact, checking this for $r = 1$ suffices due to the existence of a metric compatible scalar multiplication operation, and we may further assume without loss of generality that $n = 1$. For the first equation this follows immediately from Exercises 3.11 and 3.12. For the second equation suppose, by way of contradiction, that there is $x = (x_i) \in \mathcal{D}_1(\prod_{\mathcal{U}} E_i)^+$ so that $\lim_{\mathcal{U}} \text{dist}(x_i, \mathcal{D}_1(E_i^+)) =: \alpha > 0$. We can clearly assume that each x_i is hermitian. Since $\|x_i - 1\| \geq \alpha/2$ for $i \in \mathcal{U}$ generic, we have that $\|x_i\| \geq 1 + \alpha/2$ for $i \in \mathcal{U}$ generic. However, this contradicts that $1 = \|x\| = \lim_{\mathcal{U}} \|x_i\|$, and we have established the second equation holds.

Notation 3.14. In the case that $E_i \equiv E$, we write $E^{\mathcal{U}} := \prod_{\mathcal{U}} E_i$ and refer to $E^{\mathcal{U}}$ as the \mathcal{U} -ultrapower of E .

Let \mathcal{C} be a class of \mathcal{L} -structures. We say that \mathcal{C} is closed under taking *ultraroots* if $E \in \mathcal{C}$ whenever $E^{\mathcal{U}}$ for some ultrafilter \mathcal{U} . The following appears as [15, Theorem 2.4.1]

Proposition 3.15. *A class \mathcal{C} of \mathcal{L} -structures is elementary (in the sense of Definition 3.10) if and only if \mathcal{C} is closed under isomorphisms, ultraproducts, and taking ultraroots.*

Example 3.16. We say an operator system E is *minimal* if it is a subsystem of an abelian C*-algebra; equivalently, if $\widehat{E} \subset C(\mathcal{S}(E))$ is a complete order embedding. Since the operator system ultraproduct of unital (abelian) C*-algebras is again a unital (abelian) C*-algebra under the natural multiplicative structure, we have that the class of minimal operator systems is elementary.

By the same token, if \mathcal{C} is an elementary class of unital C*-algebras, then the class of all subsystems of elements of \mathcal{C} is an elementary class of operator systems.

Question 3.17. Is there an elementary class of operator systems which is closed under taking subsystems which is not the class of all subsystems of some elementary class of unital C*-algebras?

3.4. Definability. From a practical viewpoint, the most important task at hand once a theory T is constructed is to begin to explore the definable sets in the models of T . In short, this is because the definable sets are exactly those which it is permissible to quantify over, so having a large class of natural sets being definable allows for a great range of intuitive constructions of formulas, “breathing life” into the theory. From the point of view of a working analyst, the concept of definability is perhaps the key feature of continuous model theory as to say that a subset of a structure is definable is to say that it possesses stability/rigidity under small perturbations. In this light many important results in the theory of operator systems and C*-algebras can be seen as establishing the

definability of sets elements satisfying certain formulas. Our treatment of definability here is directly taken from [15, Chapter 3], though we will center our discussion here on the practical definition of definability for sets which is afforded by the Beth Definability Theorem: see [15, Section 4.2] or [5, Theorem 9.32].

Let T be a theory and \mathcal{M} an elementary class of models of T . Let F be an assignment to each $A \in \mathcal{M}$ a closed subset $F(A) \subset \prod_{i=1}^d \mathcal{D}_{r_i}(A)$, where $\mathcal{D}_{r_1}(A), \dots, \mathcal{D}_{r_d}(A)$ are sorts in A for d, r_1, \dots, r_d fixed. We write

$$\text{codomain}(F) := \prod_{i=1}^d \mathcal{D}_{r_i}.$$

Definition 3.18. We say that F is a *uniform assignment* if F is a functor, that is, for every homomorphism $\varphi : A \rightarrow B$ with $A, B \in \mathcal{M}$ we have that $\varphi(F(A)) \subset F(B)$.

Remark 3.19. In the language of operator systems defined above we have that the morphisms are always the unital, completely positive maps which are, rather conveniently, all contractions.

When considering the model theory of operator systems (or C^* -algebras) it makes sense to introduce the following unbounded variant of a uniform assignment. Let \mathcal{M} be an elementary class in the theory of operator systems. Suppose that F assigns to every $A \in \mathcal{M}$ a closed subset

$$F(A) \subset M_{r_1}(A) \times \cdots \times M_{r_k}(A) \times M_{s_1}(A)^+ \times \cdots \times M_{s_l}(A)^+ \times \mathbb{C}^d.$$

We say that F is then an *unbounded uniform assignment* if its restriction to every product of sorts is a uniform assignment.

Definition 3.20 (Beth Definability). Let \mathcal{M} be an elementary class of operator systems and F be an unbounded uniform assignment on \mathcal{M} . We say that F is *definable* or that $F(E)$ is a *definable set* for each $E \in \mathcal{M}$ if for every ultraproduct $\prod_{\mathcal{U}} E_i$ of elements $E_i \in \mathcal{M}$ we have that

$$F\left(\prod_{\mathcal{U}} E_i\right) = \prod_{\mathcal{U}} F(E_i).$$

To every (unbounded) formula $f(x)$, there is a natural (unbounded) uniform assignment Z_f which assigns to every $E \in \mathcal{M}$ the *zero set*

$$Z_f(E) = \{x \in \text{domain}(f)^E : f(x) = 0\}.$$

Exercise 3.21. Let f be a formula. We have that Z_f is definable if and only if for every $\epsilon > 0$ there is $\delta > 0$ so that for all $A \in \mathcal{M}$ and $x \in \text{domain}(f)^A$, we have that $f(x)^A \leq \delta$ implies that $\text{dist}(x, Z_f) \leq \epsilon$.

Theorem 3.22 (Beth Definability Theorem). *Let \mathcal{M} be an elementary class of models of a theory T , and let F be a uniform assignment. Writing*

$$p_F(x) := \text{dist}(x, F), \quad x \in \text{codomain}(F),$$

we have that F is definable if and only if there is a sequence of formulas f_1, f_2, \dots with $\text{domain}(f_i) = \text{codomain}(F)$ so that

$$\sup_{A \in \mathcal{M}} \sup \{ |p_F(x)^A - f_n(x)^A| : x \in \text{codomain}(F)^A \} \leq 1/n.$$

We refer the reader to [15, Section 4.2] or [5, Chapter 9] for a proof.

Definition 3.23. If F is an (unbounded) uniform assignment, we refer to $p_F(x) = \text{dist}(x, F)$ as a *definable predicate*. In the unbounded case, by the Beth Definability Theorem p_F restricted to any product of sorts in $\text{codomain}(F)$ is a uniform limit of formulas.

Example 3.24. The hermitian elements in any operator system form a definable set.

Example 3.25. For each n we have that $d_n^+(x) := \text{dist}(x, i_n(C_n))$, $x \in E_n$, is a definable predicate.

Remark 3.26. In fact, something even stronger may be said. Consider the (unbounded) quantifier-free formula

$$p_{t,n}(x) := \|2x - t1_n\|_n \dot{\leq} t$$

for $t \geq 0$ with domain E_n^h . We have that $p_{t,n}(x) = 0$ if and only if $\|x\|_n \leq r$ and $x \in C_n$. This is because for a hermitian element $x \in E_n$ we have that $\|2x - t1_n\|_n \leq t$ if and only if $0 \preceq x \preceq t1_n$. Therefore, every formula in the language of operators systems can be replaced with an equivalent formula where all quantifiers have domain in E_n for some n . That is, it is never necessary to quantify over the positive cones.

Proposition 3.27. For each homogeneous, hermitian linear matrix *-polynomial $p(x)$ of degree d in $x = (x_1, \dots, x_n)$, the set of tuples $X = (X_1, \dots, X_n)$ in E_k or E_k^h satisfying $1_d \otimes 1_k \succeq p(X)$ is a definable set.

Proof. For ease of notation we will write 1 for $1_d \otimes 1_k$. Suppose that $\text{dist}(1 - p(X), C_{dk}) < \epsilon$. Since $1 - p(X)$ is hermitian, we claim

$$(1 + \epsilon)1 \succeq p(X).$$

Indeed, since there exists $Y \in C_{dk}$ so that $\|1 - p(X) - Y\|_{dk} \leq \epsilon$, we have that

$$\epsilon 1 \succeq 1 - p(X) - Y \succeq -\epsilon 1, \text{ so } 1 - p(X) \succeq Y - \epsilon 1 \succeq -\epsilon 1.$$

Setting $X' = \frac{1}{1+\epsilon}X$, by linearity and homogeneity $1 \succeq p(X')$ and $\|X'_i - X_i\|_k \leq \epsilon \|X_i\|_k$ for $i = 1, \dots, n$. The result now follows by Beth Definability and Exercise 3.21. \square

Question 3.28. If $p(x)$ is a homogeneous linear matrix *-polynomial, is the set of all tuples X_1, \dots, X_n in E_k satisfying $1 \succeq p(X)$ definable? If not, is there a natural class of operator systems or C*-algebras where all such sets are definable?

Remark 3.29. We may define a linear operator *-polynomial to be an expression of the form $q(x) = A_1 \otimes x_1 + \dots + A_n \otimes x_n + B_1 \otimes x_1^* + \dots + B_n \otimes x_n^* + C \otimes 1$ where $A_1, \dots, A_n, B_1, \dots, B_n, C \in \mathcal{B}(H)$. All terminology from linear matrix *-polynomials carries over to this more general context equally well. Just as a finite family of linear

matrix $*$ -polynomials can be combined into a single linear matrix $*$ -polynomial, any infinite family of linear matrix $*$ -polynomials can be combined into a single linear operator $*$ -polynomial.

The main point is that by the same proof as Proposition 3.27, we have the following.

Proposition 3.30. *For each homogeneous, hermitian linear operator $*$ -polynomial $q(x)$ in $x = (x_1, \dots, x_n)$, the set of tuples X_1, \dots, X_n in E_k or E_k^h satisfying $1_H \otimes 1_k \succeq q(X)$ is definable.*

Via block diagonal embedding we have the following consequence.

Corollary 3.31. *Let $\{p_i : i \in I\}$ be a collection of homogeneous, hermitian linear matrix $*$ -polynomials in $x = (x_1, \dots, x_n)$. The set of tuples X_1, \dots, X_n in E_k or E_k^h satisfying $1_{d_i} \otimes 1_k \succeq p_i(X)$ for all $i \in I$ is definable.*

For an operator system E we may regard the set $\text{UCP}(M_n, E)$ as a closed subset of $\mathcal{D}_1(E)^{n^2}$ via the correspondence $\varphi \leftrightarrow (\varphi(e_{ij}))_{ij}$. We following result was first observed in [15, Section 5.8].

Proposition 3.32. *For each n we have that $\text{UCP}(M_n, E)$ is a definable set for the class of all operator systems.*

Proof. We have by Lemma 2.24 that $\text{UCP}(M_n, E) \cong \text{UCP}(\mathbb{C}, M_n(E))$. This correspondence identifies $\text{UCP}(M_n, E)$ with the set of all $X \in M_n(E)^+$ with $\text{tr}(X) = \sum_i X_{ii} = 1$. Suppose $E = \prod_{\mathcal{U}} E_i$. We have by the remarks after Definition 3.13 that for any $X \in M_n(E)^+$ there is a net $X_i \in M_n(E_i)^+$ with $X = (X_i)$. Given any $\epsilon > 0$ we have that $\|\text{tr}(X_i) - 1\| \leq \epsilon$ for $i \in \mathcal{U}$ generic. Let $b_i := \text{tr}(X_i) \in E^+$. Since $\|b_i - 1\| \leq \epsilon$ we have that b is invertible with $\|b_i^{-1/2} - 1\| < \epsilon^{1/2}$. Setting

$$[X'_i]_{kl} := b_i^{-1/2} [X_i]_{kl} b_i^{-1/2}$$

we have that $X'_i \in M_n(E_i)^+$ with $\text{tr}(X'_i) = 1$ and $\|X_i - X'_i\| \leq 2n^2\epsilon^{1/2}$. \square

Remark 3.33. Note that the proof above shows that $\text{UCP}(M_n, E)$ is explicitly the zero set of the quantifier-free (positive) formula $f(X) := \|\text{tr}(X) - 1\|$ where $\text{domain}(f) = \mathcal{D}_{n^2}(C_n)$.

Let S_n be the uniform assignment with codomain $\mathcal{D}_1(E)^n \times \mathbb{C}^n$ which assigns to each operator system the (closed) subset of all $(x_1, \dots, x_n, \lambda_1, \dots, \lambda_n)$ so that $x_i \mapsto \lambda_i$ extends to a state. The following result is essentially contained in [15, Section 5.8], though we give a different proof here.

Proposition 3.34. *We have that S_n is quantifier-free definable.*

Proof. We will exhibit S_n as the zero set of a uniform limit of quantifier-free formulas. Let $\mathbb{D} \subset \mathbb{C}$ be the closed unit disk. For convenience we write $x = (x_{-n}, \dots, x_n)$ with $x_0 = 1$ and $x_{-i} = x_i^*$ and $\lambda = (\lambda_{-n}, \dots, \lambda_n)$ with $\lambda_{-i} = \bar{\lambda}_i$ and $\lambda_0 = 1$. Let

$$f_n(x, \lambda) := \sup_{a_i \in \mathbb{D}} \left(\left\| \sum_i a_i \lambda_i \right\| \div \left\| \sum_i a_i x_i \right\| \right)$$

and note that f_n is a uniform limit of quantifier-free formulas. (Recall $x \dot{-} y = \max\{x - y, 0\}$.) Indeed, let (K_p) be a finite $1/p$ -net in \mathbb{D} , and note that f_n is a uniform limit of the formulas

$$f_{n,p}(x, \lambda) = \max_{a_i \in K_p} \left(\left\| \sum_i a_i \lambda_i \right\| \dot{-} \left\| \sum_i a_i x_i \right\| \right)$$

by an application of the triangle inequality.

From now on, let us fix an operator system E . We claim that $f(x, \lambda)^E = 0$ if and only if $x_i \mapsto \lambda_i$ extends to a state on E . By Proposition 2.7, Lemma 2.22, and Theorem 2.26 it suffices to check that $f_n(x, \lambda)^E = 0$ implies that $\varphi : x_i \mapsto \lambda_i$ defines a unital, $*$ -linear contraction on $F := \text{span}\{x_{-n}, \dots, x_n\}$. Observe that $f_n(x, \lambda)^E = 0$ is equivalent to

$$\left\| \sum_i a_i x_i \right\| \geq \left\| \sum_i a_i \lambda_i \right\| \text{ for all } a_{-n}, \dots, a_n \in \mathbb{D}.$$

Suppose that $\sum_i a_i x_i = \sum_i b_i x_i$ so that $\sum_i \frac{1}{2}(a_i - b_i)x_i = 0$; thus, $\sum_i \frac{1}{2}(a_i - b_i)\lambda_i = 0$. This shows that $\varphi : F \rightarrow \mathbb{C}$ is a well-defined, unital, $*$ -linear map. Finally, $\{\sum_i a_i x_i : a_i \in \mathbb{D}\}$ contains a sufficiently small ball about the origin in F , so φ is a contraction. \square

Since $M_n(\mathbb{C}) \subset M_n(E)$ canonically $*$ -isometrically for any operator system E , we have the following corollary.

Corollary 3.35. *Let $S_{n,k}$ be the uniform assignment with codomain $\mathcal{D}_1(E)^n \times \mathcal{D}_1(M_k(E))^n$ which assigns to each operator system the (closed) subset of all $(x_1, \dots, x_n, y_1, \dots, y_n)$ so that $y_1, \dots, y_n \in M_k(\mathbb{C})$ and $x_i \mapsto y_i$ extends to a matrix state. We have that $S_{n,k}$ is quantifier-free definable.*

Remark 3.36. The previous proofs do not show that the $S_{n,k}$ are *positive* quantifier-free definable since $\dot{-}$ is not a positive connective. In fact, it is not possible for $n \geq 5$ for $S_{n,k}$ to be positive quantifier-free definable for all k . This is because such a result would show that any quotient of a nuclear operator system (see section 4.1 below) would be nuclear, essentially following the reasoning of [15, Section 5.14], but there is five-dimensional nuclear operator system with a non-nuclear quotient which is a variant of Example 2.34: see [27].

3.5. Model Theory of C*-Algebras. We will treat the model theory of C*-algebras in a much more abbreviated fashion than operator systems. We refer the reader to Szabo's article in this volume for background on the basic theory of C*-algebras and to [15, 16] for an in-depth treatment of the model theory. In brief, the language of C*-algebras is the language of a normed, complex algebra A with a unary involution function $*$ satisfying the axiom

$$\|x^*x\| = \|x\|^2 \text{ for all } x \in A.$$

The sorts are, naturally, axiomatized to be the balls around the origin. This is all spelled out in [16, Section 3.1] or [15, Section 2.2]. The atomic formulas in the theory of C*-algebras are of the form $\|p(x)\|$ where p is a noncommutative $*$ -polynomial in the variables $x = (x_1, \dots, x_n)$.

Let A be a C^* -algebra. We say that an element $x \in A$ is *positive* if $x = y^*y$ for some $y \in A$. (Importantly, y can be chosen with $\|y\| = \sqrt{\|x\|}$.) It is well known that the set A^+ of positive elements forms a closed cone. In the case that A is a unital C^* -algebra then the unit is an archimedean order unit with respect to A^+ . As in the case of operator systems, this positivity structure is crucial to the theory of C^* -algebras.

Exercise 3.37. We have that A^+ is an (unbounded) definable set in the language of C^* -algebras; thus, $\text{dist}(x, A^+)$ is a definable predicate.

As noted above, every unital C^* -algebra is an operator system under the natural positivity structure imposed on $M_n(A)$ by the fact that $M_n(A)$ is again a C^* -algebra. We would now like to establish that the every property about A that can be expressed in the language of operator systems can be expressed in the language of C^* -algebras. This reduces to proving the following result, due to Lupini, which can be found in [20, Appendix C].

Proposition 3.38. *Let A be a unital C^* -algebra. The norm unit ball of $M_n(A)$ is a definable subset of $\mathcal{D}_1(A)^{n^2}$. Thus, the operator system structure on A is interpretable in the language of C^* -algebras.*

We begin by noting the following lemma.

Lemma 3.39. *The set of all tuples (x_1, \dots, x_n) so that $\|\sum_{i=1}^n x_i^* x_i\| \leq 1$ is definable.*

Proof. Since $\sum_{i=1}^n x_i^* x_i$ is positive, we have that $\|\sum_{i=1}^n x_i^* x_i\| \leq 1$ if and only if

$$\sum_{i=1}^n x_i^* x_i \preceq 1.$$

The proof now follows along the similar lines to the proof of Proposition 3.27. \square

Proof of Proposition 3.38. Let $x \in M_n(A)$. Any unital representation $\pi : A \rightarrow \mathcal{B}(H)$ extends canonically to a unital representation $\tilde{\pi} : M_n(A) \rightarrow \mathcal{B}(H^{\oplus n})$. It is not hard to see that every unital representation of $M_n(A)$ arises this way: for a representation $\rho : M_n(A) \rightarrow \mathcal{B}(K)$, set $H = \rho(e_{11})K$ and $\pi(x) = \rho(xe_{11})$ for $x \in A$. Thus,

$$\|x\|_n = \sup_{(H, \pi)} \sup_{\xi_i, \eta_j \in H} \left\{ \left| \sum_{i,j} \langle x_{ij} \xi_i, \eta_j \rangle \right| : \sum_{i=1}^n \|\xi_i\|^2, \sum_{j=1}^n \|\eta_j\|^2 \leq 1 \right\}$$

over all unital representations $\pi : A \rightarrow \mathcal{B}(H)$. In fact, it suffices to consider cyclic representations, each of which is given up to unitary equivalence by a state ϕ on A by the GNS construction $\phi \rightsquigarrow (H_\phi, \pi_\phi)$. Since $\pi_\phi(A)\hat{1}_\phi$ is dense in H_ϕ for any cyclic vector $\hat{1}_\phi$ corresponding to the unit in A in the GNS construction, we have that

$$\|x\|_n = \sup_{\phi \in \mathcal{S}(A)} \sup_{a_i, b_j \in A} \left\{ \left| \phi \left(\sum_{i,j} a_i^* x_{ij} b_j \right) \right| : \phi \left(\sum_i a_i^* a_i \right), \phi \left(\sum_j b_j^* b_j \right) \leq 1 \right\}$$

which by interchanging the order of the suprema equals

$$\sup_{a_i, b_j \in A} \left\{ \left\| \sum_{i,j} a_i^* x_{ij} b_j \right\| : \left\| \sum_i a_i^* a_i \right\|, \left\| \sum_j b_j^* b_j \right\| \leq 1 \right\}.$$

The result now follows since $\| \sum_i a_i^* a_i \| \leq 1$ and $\| \sum_j b_j^* b_j \| \leq 1$ are definable sets by Lemma 3.39. \square

Remark 3.40. Having just established that every formula in the language of operator systems is interpretable in the language of C*-algebras, it is natural to ask whether the converse is true. The following result shown in [22] verifies that this is indeed the case.

Proposition 3.41. *The elementary class \mathcal{C} of all unital C*-algebras is an elementary class in the language of operator systems.*

By Proposition 3.15, establishing that \mathcal{C} forms an elementary class is equivalent to establishing that if E is an operator system so that the operator system structure on some ultrapower $E^{\mathcal{U}}$ is a reduct of some C*-algebra structure on $E^{\mathcal{U}}$, then the same is true for E . We note that while these languages are thus bi-interpretable (as adding a unit to a C*-algebra is a definable operation), the quantifier complexity of equivalent formulas may differ depending on the language chosen.

4. MODEL THEORY AND FINITE-DIMENSIONAL APPROXIMATION PROPERTIES

The importance of finite-rank approximations to the theory of Banach spaces and its connection with the theory of Banach-space tensor products was first made apparent by the seminal work of Grothendieck [23]. In case of C*-algebras, the theory of finite-dimensional (matrix) approximations and the largely parallel theory of tensor products of C*-algebras were developed in foundational works of Arveson, Choi, Connes, Effros, Kirchberg, Lance, Takesaki, and Tomiyama, among others, and these ideas and concepts continue to form a central part in the theory of operator algebras and operator systems. In this last section, we survey some of the connections that have been made between the model theory of operator systems and their finite-dimensional approximation properties. We refer the reader to the books of Brown and Ozawa [6] and Pisier [40] for an in-depth treatment of the connections between finite-dimensional approximation properties and tensor products of C*-algebras and operator systems.

4.1. Nuclear and Exact Operator Systems.

Definition 4.1. An operator system E is said to be *nuclear* if there are nets $\varphi_i : E \rightarrow M_{n_i}$ and $\psi_i : M_{n_i} \rightarrow E$ so that $\psi \circ \varphi$ converges in the pointwise-norm topology to the identity, that is, $\lim_i \|\psi_i \circ \varphi_i(x) - x\| = 0$ for all $x \in E$.

Remark 4.2. Technically nuclearity of a C*-algebra A refers to the condition that for every C*-algebra B there is exactly one cross norm on the algebraic tensor $A \otimes B$ which can be completed to a C*-algebra: see Szabo’s article in this volume. The equivalence of nuclearity and local completely positive factorization of the identity map through

matrix algebras is due to Choi and Effros [11] and Kirchberg [30] and was extended in the appropriate way to the operator system category in [29].

Remark 4.3.

- (1) It is trivial that M_n is nuclear for each $n = 1, 2, \dots$ and that $M_n(E)$ is nuclear if and only if E is.
- (2) The classes of AF and UHF C^* -algebras as described in Szabo's article in this volume are nuclear. The Cuntz algebras \mathcal{O}_n defined therein are also nuclear.
- (3) $\mathcal{B}(H)$ is not nuclear for H infinite dimensional: see [6, Proposition 2.4.9]
- (4) It is false in general that C^* -subalgebras of nuclear C^* -algebras are nuclear.

Remark 4.4. There is a subsystem of M_3 which is not nuclear. Indeed, let E be the subsystem of all matrices $A \in M_3$ with $A_{13} = A_{31} = 0$. If E was nuclear there would be sequences of unital completely positive maps $\varphi_n : E \rightarrow M_n$ and $\psi_n : M_n \rightarrow E$ so that $\psi_n \circ \varphi_n \rightarrow \text{id}_E$. By Arveson's extension theorem, we may extend each φ_n to a unital completely positive map $\tilde{\varphi}_n : M_3 \rightarrow M_n$, and we see that a cluster point of $\psi_n \circ \tilde{\varphi}_n$ gives a completely positive projection $p : M_3 \rightarrow E$. By [9, Theorem 3.1], this would define a C^* -algebra structure on E under the multiplication $x \cdot y := p(xy)$. There are two distinct, unital $*$ -homomorphisms $M_2 \rightarrow E$,

$$A \mapsto \begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & A \end{bmatrix}$$

which by the proof of the same result must be respected by the C^* -algebra structure on E . This is however a contradiction since E is seven-dimensional and every finite-dimensional C^* -algebra is a direct sum of matrix algebras.

Exercise 4.5. Let X be a compact, Hausdorff space. Show that $C(X)$, the algebra of continuous functions $f : X \rightarrow \mathbb{C}$ under the norm $\|f\|_\infty := \sup_x |f(x)|$ is nuclear.

With the definition and basic examples established, we now turn to understanding the model theory of nuclear operator systems.

Definition 4.6. Let T be a theory of \mathcal{L} -structures, and let $x = (x_1, x_2, \dots, x_n)$ and $y = (y_1, y_2, \dots, y_n)$ be finite collections of symbols. Let \mathcal{F} be a family of (definable) formulas for T , each depending only on x . We say that \mathcal{F} is *uniform* if there is a continuous function $u : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with $u(0) = 0$ so that

$$T \models |f(x) - f(y)| \leq u(\max_i d(x_i, y_i)), \text{ for all } f \in \mathcal{F}.$$

For a class \mathcal{C} of \mathcal{L} -structures so that $C \models T$ for all $C \in \mathcal{C}$, we say that \mathcal{C} is *definable by uniform families of formulas* if there is a collection of formulas $\{f_{n,k} : n, k = 1, 2, \dots\}$ defined on a countable number of symbols $x = (x_1, x_2, \dots)$ so that $\mathcal{F}_k := \{f_{n,k}\}$ is uniform for each k and

$$E \models T \text{ and } \sup_k \inf_n f_{n,k}(x)^E \equiv 0 \text{ if and only if } E \in \mathcal{C}.$$

We refer [15, Section 5.7] for the connection between definability by uniform families of formulas and being definable by omitting (partial) types.

Proposition 4.7. *The property that an operator system E is nuclear is definable by uniform families of existential formulas.*

Proof. The formulas $f_{n,k}$ are given by

$$f_{n,k}(x_1, \dots, x_k) := \inf \left\{ \max_i \|x_i - \psi \circ \varphi(x_i)\| : \varphi \in \text{UCP}(E, M_n), \psi \in \text{UCP}(M_n, E) \right\}$$

which are existential and definable by Propositions 3.32 and 3.34 and Corollary 3.35. Since $\psi \circ \varphi$ is a contraction, we have that each $f_{n,k}$ is 2-Lipschitz, so the families \mathcal{F}_k are uniform. Finally, if $\inf_n f_{n,k}(x)^E = 0$ for all tuples x in E , we can easily construct nets (φ_i) and (ψ_i) indexed over the directed set of pairs (F, ϵ) where F is a finite subset of E and $\epsilon > 0$ so that $\psi_i \circ \varphi_i$ converges to the identity on E in the pointwise-norm topology, so E is nuclear. Conversely, the existence of such a net easily implies that $\inf_n f_{n,k}^E \equiv 0$ for all k . \square

Question 4.8. For the elementary class of unital C*-algebras, is nuclearity defined by a uniform family of *positive* existential formulas?

Since the values of positive existential formulas decrease under surjective morphisms, this would recover the result due to Choi and Effros, [8, Corollary 3.3] and [10, Corollary 4], that nuclear C*-algebras are closed under taking C*-algebra quotients avoiding Connes' [12] celebrated but difficult work on the classification of injective factors. However, as noted in Remark 3.36, this seems like it would require substantially different ideas to avoid the false implication in the operator system category. We refer to [15, Section 5.9] for one such attempt.

Definition 4.9. Let E be a finite-dimensional operator system. We say that E is *exact* if for every $\epsilon > 0$ there is a unital, completely positive map $\varphi : E \rightarrow M_n$ for some n so that $\|\varphi^{-1}|_{\varphi(E)}\|_{\text{cb}} \leq 1 + \epsilon$. We say that an operator system is *exact* if every finite-dimensional subsystem is exact.

Remark 4.10.

- (1) Every nuclear operator system is exact.
- (2) It is clear from the definition that any subsystem of an exact operator system is exact. As a special case, every subsystem of M_n is exact.
- (3) Since every subsystem of any unital abelian C*-algebra A is exact, $\text{MIN}(V)$ is exact for any archimedean ordered *-vector space V .
- (4) A difficult theorem of Kirchberg [32] establishes that C*-algebraic quotients of exact C*-algebras are exact. This is again false in the operator system category by the same counterexample as given in Remark 3.36.

We point out the exactness, similarly to nuclearity, can be phrased as in terms of local completely positive factorization of some (equivalently, every) *inclusion* $E \subset \mathcal{B}(H)$ through matrix algebras, the crucial difference being that the images of the approximating sequence need not land back in the system itself.

Proposition 4.11. *A finite-dimensional operator system $E \subset \mathcal{B}(H)$ is exact if and only if for each $\epsilon > 0$ there are unital, completely positive maps $\varphi : E \rightarrow M_n$ and $\psi : \varphi(E) \rightarrow \mathcal{B}(H)$ so that $\|\psi \circ \varphi - \text{id}_E\| < \epsilon$ where id_E is the restriction of the identity map on $\mathcal{B}(H)$ to E .*

Proof. We have that $\varphi(E) \subset M_n$ and that $\varphi^{-1} : \varphi(E) \rightarrow \mathcal{B}(H)$ is unital and self-adjoint. By [6, Corollary B.9] there is a unital, completely positive map $\psi : \varphi(E) \rightarrow \mathcal{B}(H)$ with $\|\psi - \varphi^{-1}\|_{\text{cb}} \leq 2(\|\varphi^{-1}\|_{\text{cb}} - 1)$; hence,

$$\|\psi \circ \varphi - \text{id}_E\| \leq \|\psi \circ \varphi - \text{id}_E\|_{\text{cb}} = \|\psi \circ \varphi - \varphi^{-1} \circ \varphi\|_{\text{cb}} \leq \|\psi - \varphi^{-1}\|_{\text{cb}} \|\varphi\|_{\text{cb}} \leq 2\epsilon.$$

In the other direction, suppose there are unital, completely positive maps φ, ψ as above, and let $y_1, \dots, y_n, y_1^*, \dots, y_n^*$ be such a basis/dual basis pair for $\varphi(E)$ as given in Lemma 2.29. We have that

$$(\psi - \varphi^{-1})(z) = \sum_{i=1}^n y_i^*(z)(\psi(y_i) - \varphi^{-1}(y_i)),$$

so

$$\|\psi - \varphi^{-1}\| \leq \dim(E) \max_i \|\psi(y_i) - \varphi^{-1}(y_i)\|.$$

Setting $y_i = \varphi(x_i)$ we have that

$$\|x_i\| \leq \|\psi \circ \varphi(x_i) - x_i\| + \|\psi \circ \varphi(x_i)\| \leq \|\varphi(x_i)\| + \epsilon = 1 + \epsilon;$$

hence, $\max_i \|\psi(y_i) - \varphi^{-1}(y_i)\| = \max_i \|\psi \circ \varphi(x_i) - x_i\| \leq (1 + \epsilon)\epsilon$. \square

Given the similarities between nuclearity and exactness it is natural to wonder if exact operator systems have a similar low-complexity description in terms of the model theory. A result of Goldbring and the author [21] shows that this is, however, not the case.

Proposition 4.12. *The class of exact operator systems is not definable by uniform families of existential formulas.*

Question 4.13. Is the class of exact C^* -algebras definable by uniform families of (positive) existential formulas?

4.2. The Local Lifting Property.

Definition 4.14. An operator system E has the *lifting property* (LP) if for every unital C^* -algebra A and every ideal I , for every unital, completely positive map $\varphi : E \rightarrow A/I$ there exists a unital, completely positive map $\tilde{\varphi} : E \rightarrow A$ lifting φ , that is, $q \circ \tilde{\varphi} = \varphi$ where $q : A \rightarrow A/I$ is the quotient map. An operator system has the *local lifting property* (LLP) if every finite-dimensional subsystem has the lifting property.

The following result is due to Robertson and Smith [41].

Proposition 4.15. *Let E be a finite-dimensional operator system and B/J be a quotient C^* -algebra. For each $n = 1, 2, \dots$ every unital, completely positive map $\varphi : E \rightarrow B/J$ admits a unital n -positive lifting $\tilde{\varphi} : E \rightarrow B$.*

The following is immediate from Definition 2.37.

Corollary 4.16. *Every finite-dimensional, k -maximal operator system has the lifting property.*

Definition 4.17. A finite-dimensional operator system E is said to be *CP-stable* if for every $\epsilon > 0$ there exist $\delta > 0$ and n so that for any unital map $\varphi : E \rightarrow A$ into any C*-algebra A satisfying $\|\varphi\|_n \leq 1 + \delta$, there is a unital, completely positive map $\varphi' : E \rightarrow A$ so that $\|\varphi' - \varphi\| \leq \epsilon$.

Parts of the following result are due to Kavruk [26, Theorem 6.6] and Goldbring and the author [20, Proposition 2.42], [21, Section 7], as well as [43].

Theorem 4.18. *For a finite-dimensional operator system E , the following are equivalent:*

- (1) E has the lifting property;
- (2) E^* is exact;
- (3) E is CP-stable;
- (4) for every sequence (A_i) of unital C*-algebras and every non-principal ultrafilter \mathcal{U} , every unital, completely positive map $\varphi : E \rightarrow \prod_{\mathcal{U}} A_i$ admits a unital, completely positive lifting $\tilde{\varphi} : E \rightarrow \prod A_i$;
- (5) for every sequence (M_{n_i}) of matrix algebras and every non-principal ultrafilter, \mathcal{U} every unital, completely positive map $\varphi : E \rightarrow \prod_{\mathcal{U}} M_{n_i}$ admits a unital, completely positive lifting $\tilde{\varphi} : E \rightarrow \prod M_{n_i}$.

This result allows one to give an interesting interpretation of the lifting property for certain maximal-type operator systems in terms of definability via Proposition 3.27.

Example 4.19. Let $p(x)$ with $x = (x_1, \dots, x_n)$ be a homogeneous, hermitian linear matrix *-polynomial of degree d . Consider the closed convex set

$$K := \{x \in \mathbb{R}^n : 1_d \succeq p(x)\}$$

(Any convex set of this form is known as a *spectrahedron*.) Suppose that K is bounded with 0 as an interior point. Recalling the construction from Remark 2.38 of the archimedean ordered *-vector space $V_K = (\mathbb{C}^{n+1}, P_K, e)$ from K , we define an operator system structure on V_K by defining $K_\ell = \text{UCP}(V_K, M_\ell)$ to be given by all linear maps $f : \mathbb{R}^{n+1} \rightarrow M_\ell^h$ with $f(e_1) = I_\ell$ so that the tuple $X = (X_1, \dots, X_n)$ given by $X_i := f(e_{i+1})$ satisfies

$$1_d \otimes 1_\ell \succeq p(X)$$

where e_1, e_2, \dots, e_{n+1} form the standard basis for \mathbb{R}^{n+1} . (Note the order unit e is e_1 .) The collection of sets $(K_\ell)_{\ell \in \mathbb{N}}$ is known as the *free spectrahedron* defined by $p(x)$ [33]. Consequently, we refer to the operator system structure we have just constructed on V_K as the *free operator system* generated by p , which we will denote by $\mathcal{F}(p)$.

Proposition 4.20. *Using the same assumptions and notation as Example 4.19, the free operator system $\mathcal{F}(p)$ has the lifting property.*

Proof. Let $\varphi : \mathcal{F}(p) \rightarrow \mathcal{B}(H)$ be a unital, completely positive map, and define $X_i := \varphi(e_{i+1})$ for $i = 1, \dots, n$ as above. Since $\mathcal{F}(p)$ is finite-dimensional, we may assume without loss of generality that H is separable. (This is just to avoid the slight inconvenience of working with nets rather than sequences, but the subsequent arguments translate to nets practically verbatim.) Let $H_1 \subset H_2 \subset \dots \subset H$ be an increasing sequence of finite-dimensional subspaces whose union is dense in H , and let $p_j : H \rightarrow H_j$ be the orthogonal projection onto H_j . We have that $\varphi^{(j)} : \mathcal{F}(p) \rightarrow M_{m_j}$ defined by $\varphi^{(j)}(x) := p_j \varphi(x) p_j$ is unital, completely positive; thus, for the tuple $X^{(j)} = (X_1^{(j)}, \dots, X_n^{(j)})$ given by $X_i^{(j)} := \varphi^{(j)}(e_{i+1}) = p_j \varphi(e_{i+1}) p_j$ we have that $X^{(j)}$ is hermitian and $1_d \otimes 1_{m_j} \succeq p(X^{(j)})$ holds. Since $X^{(j)}$ converges to X in the strong operator topology, $p(X^{(j)})$ also converges strongly to $p(X)$, so $1_d \otimes 1_H \succeq p(X)$ as well.

Now let $\varphi : \mathcal{F}(p) \rightarrow \prod_{\mathcal{U}} M_{m_j}$ be a unital, completely positive map. Since $\mathcal{F}(p)$ is finite-dimensional, we may assume that $\varphi = (\varphi_j)_{\mathcal{U}}$ where $\varphi_j : \mathcal{F}(p) \rightarrow M_{m_j}$ is unital, $*$ -linear, and sends \mathbb{R}^{n+1} into $M_{m_j}^h$. Setting $Y_i^{(j)} := \varphi_j(e_{i+1})$, we have that

$$\text{dist}(1_d \otimes 1_{m_j} - p(Y^{(j)}), M_{dm_j}^+) < \epsilon \text{ for } j \in \mathcal{U} \text{ generic.}$$

By Proposition 3.27, we may perturb the tuple $Y^{(j)}$ slightly to $Z^{(j)}$ so that $\varphi_j'(e_{i+1}) := Z_i^{(j)}$ defines a unital completely positive map. We have that $\tilde{\varphi} = (\varphi_j') : \mathcal{F}(p) \rightarrow \prod M_{m_j}$ is the required lifting. By the equivalence of (1) and (5) in Theorem 4.18, the result obtains. \square

Remark 4.21. Let $q(x)$ be a homogenous, hermitian linear operator $*$ -polynomial, and assume that the closed, convex set $K = \{x \in \mathbb{R}^n : 1_H \succeq q(x)\}$ is bounded and contains 0 as an interior point. Then, we may construct an operator system $\mathcal{F}(q)$ in exactly the same way as we did in Example 4.19, which we will still refer to as the *free operator system* generated by q . The proof of the previous proposition applies equally well to this situation, using now Proposition 3.30, so $\mathcal{F}(q)$ is seen to have the lifting property. Since every closed, convex set K arises in this way by combining all linear inequalities it satisfies into a single linear operator $*$ -polynomial, this recovers that all maximal operator systems are CP-stable. (The equivalence of CP-stability and the lifting property in Theorem 4.18 itself uses Proposition 4.15.)

4.3. The Weak Expectation Property.

Definition 4.22. We say that an operator system E has the *weak expectation property* if for every unital inclusion $E \subset B$ into another operator system, there is a unital, completely positive map $\varphi : B \rightarrow E^{**}$ so that $\varphi(x) = \hat{x}$ for all $x \in E$. (Recall that $\hat{\cdot} : E \rightarrow E^{**}$ is the evaluation map as in Remark 2.21 and Proposition 2.30.)

Lemma 4.23. *An operator system E has the weak expectation property if and only if for every unital inclusion $E \subset B$ and each hermitian $b \in B^h$, there is a unital, completely positive map $\varphi : E + \mathbb{C}b \rightarrow E^{**}$ so that $\varphi(x) = \hat{x}$ for all $x \in E$.*

Proof. Let us fix $E \subset B$. We first show that for every finite set $b_1, \dots, b_n \in B$ there is a unital, completely positive map $\varphi : E + \mathbb{C}b_1 + \dots + \mathbb{C}b_n \rightarrow E^{**}$ so that $\varphi(x) = \hat{x}$ for all $x \in E$. Inductively, assume that this is true for choice of n elements, and consider $b_1, \dots, b_{n+1} \in B^h$. Let $\varphi : E + \mathbb{C}b_1 + \dots + \mathbb{C}b_n \rightarrow E^{**}$ be such a map. We have that $E^{**} \subset \mathcal{B}(H)$ for some Hilbert space H , so by Arveson's extension theorem, we may extend φ to a unital, completely positive map $\varphi' : B \rightarrow \mathcal{B}(H)$. We have that $E \cong \widehat{E} \subset \mathcal{B}(H)$, so setting $d := \varphi'(b_{n+1})$ by assumption there is a unital, completely positive map $\psi : E + \mathbb{C}d \rightarrow E^{**}$ so that $\psi(x) = \hat{x}$ for all $x \in E$. Let $\psi' : (E + \mathbb{C}d)^{**} \cong E^{**} + \mathbb{C}d \rightarrow E^{**}$ be the continuous extension of ψ in the weak* topology, which is again unital and completely positive. Since $\varphi'(E + \mathbb{C}b_1 + \dots + \mathbb{C}b_{n+1}) \subset E^{**} + \mathbb{C}d$, we have that $\psi' \circ \varphi' : E + \mathbb{C}b_1 + \dots + \mathbb{C}b_{n+1} \rightarrow E^{**}$ is the required map.

Since every $x \in B$ is a linear combination of two hermitian elements, we have demonstrated that for every subsystem $E \subset A \subset B$ with A having finite co-dimension relative to E , that there is a unital, completely positive map $\varphi_A : A \rightarrow E^{**}$ so that $\varphi(x) = \hat{x}$ for all $x \in E$. Consider the directed set \mathcal{F} of all such subsystems of B , ordered by inclusion. We may take a pointwise-weak* cluster point of the net $(\varphi_A)_{A \in \mathcal{F}}$ to define a unital, completely positive map $\varphi : B \rightarrow E^{**}$ so that $\varphi(x) = \hat{x}$ for all $x \in E$. This verifies that E has the weak expectation property. \square

Lemma 4.24. *An operator system E has the weak expectation property if and only if for every unital inclusion $E \subset B$ of operator systems for each finite subsystem $A \subset B$, $n = 1, 2, \dots$, and $\epsilon > 0$, there is a n -contractive *-linear map $\varphi : A \rightarrow E$ so that $\|\varphi(x) - x\| \leq \epsilon\|x\|$ for all $x \in E \cap A$. Moreover, it suffices to check that the second statement holds only for all $A = A_0 + \mathbb{C}b$ where $A_0 \subset E$ is a finite-dimensional subsystem and $b \in B^h$.*

Proof. The proof that the weak expectation property implies the second statement is essentially given in [20, Proposition 2.34] as there was no use of the assumption that E is a C*-algebra therein. By the proof of Proposition 2.30, we have that $M_n(E^{**})$ is completely order isomorphic to $M_n(E)^{**}$; hence, if $\varphi : B \rightarrow E^{**}$ is a unital, completely positive map witnessing the weak expectation property, by the principle of local reflexivity [1, Theorem 12.2.4] for any finite dimensional subsystem $A \subset B$ and $\epsilon > 0$ there is a contraction $\theta : M_n(A) \rightarrow M_n(E)$ so that $\|\theta(x) - \varphi_n(x)\| < \epsilon\|x\|$ for all $x \in M_n(E \cap A)$. Let \mathcal{G} be the group of all signed permutation matrices and set

$$\theta'(x) = \frac{1}{|\mathcal{G}|^2} \sum_{g, h \in \mathcal{G}} g^* \theta(gxh) h^*.$$

Since $g^* \varphi_n(gxh) h^* = \varphi_n(x)$ for all $g, h \in \mathcal{G}$, we have that $\|\theta'(x) - \varphi_n(x)\| < \epsilon\|x\|$ for all $x \in M_n(E \cap A)$ as well. It is not hard to check that $\theta'(e_{ij} \otimes x) = e_{ij} \otimes \psi(x)$, $i, j = 1, \dots, n$, for a unique linear map $\psi : A \rightarrow E$, so that ψ is n -contractive. Since ψ is linear and n -contractive, so is $\psi^*(x) := \psi(x^*)^*$; thus, the average of the two is n -contractive and *-linear, with the other properties being preserved.

In the other direction, let \mathcal{F} be the directed set of all finite-dimensional subsystems of B ordered by inclusion, and let I be the directed set of triples (A, n, ϵ) where $A \in \mathcal{F}$,

n is a positive integer, and $\epsilon > 0$, ordered by $(A', n', \epsilon') \geq (A, n, \epsilon)$ if $A' \supset A$, $n' \geq n$, and $\epsilon' \leq \epsilon$. For each $(A, n, \epsilon) \in I$, find $\varphi_{A, n, \epsilon} : A \rightarrow E$ matching the conditions of the second statement. Equip E^{**} with the weak* topology, and take $\varphi : B \rightarrow E^{**}$ to be a pointwise-weak* cluster point of the net $(\varphi_{A, n, \epsilon})$, where we identify E with $\widehat{E} \subset E^{**}$. We have that φ is unital, *-linear, and completely contractive with $\varphi(x) = \widehat{x}$ for all $x \in E$. By Proposition 2.7, φ is also completely positive.

For the last part, the previous argument can be easily adapted to verify that for each hermitian $b \in B^h$, there is a unital, completely positive map $\varphi : E + \mathbb{C}b \rightarrow E^{**}$ so that $\varphi(x) = \widehat{x}$ for all $x \in E$. Thus, the result follows by Lemma 4.23. \square

We say that a formula f is *positive* if it is built from atomic formulas only using connectives $\theta(x_1, \dots, x_n)$ satisfying $\theta(x_1, \dots, x_n) \leq \theta(y_1, \dots, y_n)$ when $x_i \leq y_i$, $i = 1, \dots, n$.

Definition 4.25. We say that an operator system E is *positively existentially closed* if for every unital inclusion $E \subset B$ of operator systems and every positive, quantifier-free formula $f(x, y)$ we have that $\inf_y f(x, y)^E = \inf_y f(x, y)^B$ for all tuples x chosen from E .

Remark 4.26. We can as easily talk about a specific inclusion $E \subset B$ being *positively existential* if the condition in the definition is met. In the operator system category, this is equivalent to the existence of an ultrafilter \mathcal{U} on some set I so that there is a unital, completely positive map $\varphi : B \rightarrow E^{\mathcal{U}}$ so that $\varphi(x) = \iota(x)$ for all $x \in E$, where $\iota : E \rightarrow E^{\mathcal{U}}$ is the constant embedding, cf. [4, Remark 4.20]. The proof of Lemma 4.24 thus shows the following.

Proposition 4.27. *The canonical inclusion $\widehat{\cdot} : E \rightarrow E^{**}$ is positively existential for any operator system E .*

Remark 4.28. Since $\inf_y f(x, y)^E \geq \inf_y f(x, y)^B$ automatically, checking positive existential closure is equivalent to checking $\inf_y f(x, y)^E \leq \inf_y f(x, y)^B$ where the tuple x is chosen from E . It is straightforward to verify that this in turn is equivalent to checking that for any finite collection $f_1(x, y), \dots, f_k(x, y)$ of atomic formulas (that is, $f_i(x, y) = \|p_i(x, y)\|_{d_i}$ for $p_i(x, y)$ a linear matrix *-polynomial of degree d_i), any $\epsilon > 0$, and any tuples a in E and b in B , there is a tuple b' in E so that $f_i(a, b') \leq f_i(a, b) + \epsilon$ for all $i = 1, \dots, k$.

We say that an element $x \in \mathcal{B}(H)$ is *strictly positive* if $x - \epsilon 1$ is positive for some $\epsilon > 0$. We write $x \succ 0$ to denote the x is strictly positive.

Lemma 4.29. *An operator system is positively existentially closed if and only if for any unital inclusion $E \subset B$ and any finite collection $p_1(x, y), \dots, p_k(x, y)$ of hermitian linear matrix *-polynomials, the following property holds:*

(R) *for any tuples a in E and b in B so that*

$$p_1(a, b) \succ 0, \dots, p_k(a, b) \succ 0,$$

there is a tuple b' in E so that

$$p_1(a, b') \succ 0, \dots, p_k(a, b') \succ 0.$$

Proof. Assume that (R) holds. Let $p_1(x, y), \dots, p_k(x, y)$ be a collection of linear matrix *-polynomials of degree d_1, \dots, d_k . Fixing tuples a in E and b in B and $\epsilon > 0$, define $r_i := \|p_i(a, b)\|_{d_i}$ and consider the hermitian, linear matrix *-polynomials $q_1(x, y), \dots, q_k(x, y)$ where

$$q_i(x, y) := \begin{bmatrix} (r_i + \epsilon)1_{d_i} & p_i(x, y) \\ p_i^*(x, y) & (r_i + \epsilon)1_{d_i} \end{bmatrix}.$$

By Proposition 2.5, we have that $q_i(a, b) \succ 0$ for all $i = 1, \dots, k$. Therefore, by (R) there exists a tuple b' in E so that $q_i(a, b') \succ 0$ for all $i = 1, \dots, k$, and again by Proposition 2.5 we have that $\|p_i(a, b')\|_{d_i} < r_i + \epsilon$ for $i = 1, \dots, k$. That E is positively existentially closed now follows from Remark 4.28.

For the converse, let $p_1(x, y), \dots, p_k(x, y)$ be a collection of hermitian linear matrix *-polynomials of degrees d_1, \dots, d_k , and fix tuples a in E and b in B so that $p_i(a, b) \succeq \epsilon 1_{d_i}$ for all $i = 1, \dots, k$ for some $\epsilon > 0$ sufficiently small. Setting $r_i = \|p_i(a, b)\|_{d_i}$, we define

$$q_i(x, y) := 2p_i(x, y) - (r_i + \epsilon)1_{d_i}.$$

By some basic arithmetic we have that $-(r_i - \epsilon)1_{d_i} \preceq q_i(a, b) \preceq (r_i - \epsilon)1_{d_i}$, which is equivalent to $\|q_i(a, b)\|_{d_i} \leq r_i - \epsilon$. Thus E being positively existentially closed guarantees that there is a tuple b' in E so that $\|q_i(a, b')\|_{d_i} \leq r_i$ for all $i = 1, \dots, k$. Since $q_i(x, y)$ is hermitian it only takes values in the hermitian elements; thus,

$$-r_i 1_{d_i} \preceq q_i(a, b') = 2p_i(a, b') - r_i 1_{d_i} - \epsilon 1_{d_i}$$

for all $i = 1, \dots, k$ which shows that $p_i(a, b') \succ 0$ for $i = 1, \dots, k$. \square

Since a positivity check on a collection of linear matrix *-polynomials is equivalent to a positivity check on a single linear matrix *-polynomial obtained from the collection via block diagonal embedding, the following is immediate from the proof of the previous result.

Corollary 4.30. *An operator system is positively existentially closed if and only if for any unital inclusion $E \subset B$ and any hermitian linear matrix *-polynomial $p(x, y)$ of degree d , one of the following property holds:*

- (R₁) *for any tuples a in E and b in B so that $p(a, b) \succ 0$ there is a tuple b' in E so that $p(a, b') \succ 0$.*
- (R₂) *for any tuples a in E and b in B so that $\|p(a, b)\|_d < 1$ there is a tuple b' in E so that $\|p(a, b')\|_d < 1$.*

The following result was proved by Goldbring and the author, [20, Section 2.4] and [21, Proposition 5.1], and by Lupini [35, Proposition 3.2] in full generality. We note that in light of Remark 4.26 a characterization of the weak expectation property very much in the same spirit appears as [40, Theorem 9.22].

Proposition 4.31. *An operator system has the weak expectation property if and only if it is positively existentially closed.*

Proof. Assume that E has the weak expectation property and that $E \subset B$ is an inclusion of operator systems. Let $p(x, y)$ be a hermitian linear matrix $*$ -polynomial. Suppose there are tuples $a = (a_1, \dots, a_n)$ in E and $b = (b_1, \dots, b_k)$ in B so that $\|p(a, b)\|_d < 1$, and let $A \subset B$ be the subsystem spanned by $a_1, \dots, a_n, b_1, \dots, b_k$. As guaranteed by Lemma 4.24, choose $\varphi : A \rightarrow E$ to be $*$ -linear, d -contractive, and have the property that $\max_i \|\varphi(a_i) - a_i\| < \epsilon$ for $\epsilon > 0$ suitably small, to be determined later. We have that

$$\|p(a, \varphi(b))\|_d \leq \|p(\varphi(a), \varphi(b))\|_d + \Delta(\epsilon) = \|\varphi(p(a, b))\|_d + \Delta(\epsilon) \leq \|p(a, b)\|_d + \Delta(\epsilon),$$

where $\Delta(\epsilon)$ is the modulus of continuity for $x \mapsto \|p(x, \varphi(b))\|_d$. We now choose $\epsilon > 0$ so that $\|p(a, b)\|_d + \Delta(\epsilon) < 1$.

In the other direction, it is clear that for any tuples a in E , b in B , and c, d in E^{**} that the statement, “the $*$ -linear map φ on the span of a and b induced by sending a to c and b to d is n -contractive and satisfies $\max_i \|a_i - c_i\| \leq \epsilon$,” can be verified by a (countably) infinite collection of expressions of the form $\|p_i(a, b)\|_n < r_i \Rightarrow \|p_i(c, d)\|_n < r_i$ for p_i linear matrix $*$ -polynomials with complex rational coefficients; thus, fixing a and b by taking weak $*$ -limits of tuples c_i, d_i in $E = \widehat{E} \subset E^{**}$ satisfying the first i statements we can define an n -contractive $*$ -linear map $\varphi_{a,b}$ from the span of a and b to E^{**} satisfying $\max_i \|\varphi(a_i) - \hat{a}_i\| \leq \epsilon$. The required map $\varphi : B \rightarrow E^{**}$ is then obtained by taking a pointwise-weak $*$ limit of the $\varphi_{a,b}$ ’s over the directed set of all finite tuples and numbers $\epsilon > 0$. \square

In light of Lemmas 4.24 and 4.29, the proof of the previous proposition can be easily adapted to prove the following.

Corollary 4.32. *An operator system is positively existentially closed if and only if for any unital inclusion $E \subset B$ and any hermitian linear matrix $*$ -polynomial $p(x, y) = p(x_1, \dots, x_n, y)$ of degree d , one of the following property holds:*

- (R₃) *for any tuple a in E and $b \in B^h$ so that $p(a, b) \succ 0$ there is $b' \in E$ so that $p(a, b') \succ 0$.*
- (R₄) *for any tuple a in E and $b \in B^h$ so that $\|p(a, b)\|_d < 1$ there is a tuple $b' \in E^h$ so that $\|p(a, b')\| < 1$.*

Question 4.33. Is there a proof of this corollary which uses positive existential closure directly without passing through the equivalence with the weak expectation property?

Remark 4.34. For the (elementary) class of operator systems which are C^* -algebras, there is an explicit, countable collection of universal formulas which verifies the weak expectation property due to Farenick, Paulsen, and Kavruk [18, Theorem 6.1]. Consider the hermitian linear matrix $*$ -polynomial

$$p(x_1, x_2, y_1, y_2) := \begin{bmatrix} y_1 & x_1 & 0 \\ x_1^* & y_2 & x_2 \\ 0 & x_2^* & 1 - y_1 - y_2 \end{bmatrix}.$$

A unital C^* -algebra A has the weak expectation property if and only if for every unital inclusion $A \subset B$ of operator systems and every $n = 1, 2, \dots$ whenever $a_1, a_2 \in M_n(A)$

and $b_1, b_2 \in M_n(B)$ are so that $p(a_1, a_2, b_1, b_2) \succ 0$, there are $b'_1, b'_2 \in M_n(A)$ so that $p(a_1, a_2, b'_1, b'_2) \succ 0$.

Question 4.35. Can this result be extended to the entire class of operator systems?

Remark 4.36. By replacing the entries of p in the Farenick–Kavruk–Paulsen result with matrices, we actually increase the number of variables as each variable is replaced by a collection of variables standing in for the matrix units.

Question 4.37. Is there a countable sequence $p_i(x, y)$ of hermitian linear matrix *-polynomials in a *uniformly bounded* number of variables so that a C*-algebra has the weak expectation property if and only if for all embeddings $A \subset \mathcal{B}(H)$ and all tuples a in A and b in $\mathcal{B}(H)$ so that $p_i(a, b) \succ 0$ for all $i = 1, 2, \dots$ there is a tuple b' in A so that $p_i(a, b') \succ 0$? Would a single p suffice?

It was shown in [21, Section 4] that the weak expectation property is not elementary, but the use of the strict positivity condition in the above seems to preclude its application here.

Remark 4.38. Perhaps the simplest non-trivial hermitian linear matrix *-polynomial is

$$p(x, y) = \begin{bmatrix} 1 - y & x^* \\ x & y \end{bmatrix}.$$

For any C*-algebra $A \subset \mathcal{B}(H)$ it is an exercise in functional calculus to show that if for any $a \in A$ there is some $b \in \mathcal{B}(H)$ so that $p(a, b) \succ 0$, then there is a $b' \in A$ so that $p(a, b') \succ 0$. (It is a bit easier to see that there is $b' \in A^{**}$ since A^{**} is a von Neumann algebra, then use Proposition 4.27.) As is pointed out in the introduction of [18], we have that $\exists b : p(a, b) \succ 0$ is equivalent to the *numerical radius* $\omega(x)$ of x being less than $1/2$. It would be interesting to know if the weak expectation property could be characterized in terms of this simple expression.

Question 4.39. Does a C*-algebra A have the weak expectation property if for some (for all) n for all $a_1, \dots, a_n \in A$ for which there exists $b \in \mathcal{B}(H)$ so that $p(a_i, b) \succ 0$ for all $i = 1, \dots, n$ there is $b' \in A$ so that $p(a_i, b') \succ 0$ for all $i = 1, \dots, n$?

We close this section with a model-theoretic proof of a characterization of the weak expectation property which is an unpublished result due to Kavruk [25]. Kavruk’s proof uses the theory of operator system tensor products [28, 29] along with Kirchberg’s tensorial characterization of the weak expectation property [31].

Notation 4.40. In the following, we write \bowtie to indicate a fixed choice of \prec or \succ , so that, for instance, $a_1 \bowtie b_1, a_2 \bowtie b_2$ means one of four possible expressions with one choice of \prec or \succ for each pair a_i, b_i which remains fixed throughout.

Definition 4.41. We say that any operator system E has the *complete tight Riesz interpolation property* if for any unital inclusion of operator systems $E \subset B$, any $n = 1, 2, \dots$, and any tuple $a = (a_1 \dots, a_k)$ in $M_n(E)^h$ and $b \in M_n(B)^h$ if $b \bowtie a_i$ for all $i = 1, \dots, k$, then there is a $b' \in M_n(E)^h$ so that $b' \bowtie a_i$ for all $i = 1, \dots, k$.

Theorem 4.42 (Kavruk [25, Theorem 7.4]). *An operator system E has the complete tight Riesz interpolation property if and only if it has the weak expectation property.*

Proof. Proposition 4.31 and Lemma 4.29 show that the weak expectation property implies the complete tight Riesz interpolation property.

For the converse, we will show that the complete tight Riesz interpolation property implies property (R_3) , therefore the weak expectation property by Proposition 4.31 and Corollary 4.32. Let $E \subset B$ be a unital inclusion of operator systems and $p(x_1, \dots, x_n, y)$ be a hermitian linear matrix $*$ -polynomial of degree d . Assume we have chosen $a_1, \dots, a_n \in E$ and $b \in B^h$ so that $p(a_1, \dots, a_n, b) \succ 0$. Since b is hermitian, we have that

$$p(a_1, \dots, a_n, b) = X \otimes b - A \succ 0$$

where $X \in M_d^h$ and $A \in M_n(E)^h$. By perturbing X by ϵI_d for ϵ sufficiently small, we may assume without loss of generality that X is invertible. Therefore, we have that $X = Y \Sigma Y^*$ for Y invertible and Σ a diagonal matrix with ± 1 entries on the main diagonal. We may assume the first f entries of the main diagonal of Σ are 1 with the remaining being -1 . Let \mathcal{G} be the group of signed permutation matrices in M_d preserving the set $\{\pm e_1, \dots, \pm e_f\}$ where e_1, \dots, e_d is the standard basis for \mathbb{C}^d .

For each $g \in \mathcal{G}$, define $A_g := gY^{-1}A(Y^{-1})^*g^* \in M_d(E)^h$, and consider the system of inequalities

$$\{\Sigma \otimes b \succ A_g : g \in \mathcal{G}\},$$

noting that this system is consistent if and only if $X \otimes b \succ A$ since $g(\Sigma \otimes b)g^* = \Sigma \otimes b$ for all $g \in \mathcal{G}$. By the complete tight Riesz interpolation property, there is $b' \in M_d(E)^h$ so that $b' \succ A_g$ for all $g \in \mathcal{G}$. We have that $gb'g^* \succ Y^{-1}A(Y^{-1})^*$ for all $g \in \mathcal{G}$; thus,

$$b'' := \frac{1}{|\mathcal{G}|} \sum_{g \in \mathcal{G}} gb'g^* \succ Y^{-1}A(Y^{-1})^*.$$

It is not hard to check that b'' is block diagonal of the form

$$b'' = \begin{bmatrix} I_f \otimes b_1 & 0 \\ 0 & I_{d-f} \otimes b_2 \end{bmatrix}$$

for some $b_1, b_2 \in E^h$. It now suffices to find $c \in E^h$ with $c \succ b_1 - \epsilon 1$ and $-c \succ b_2 - \epsilon 1$ for $\epsilon > 0$ sufficiently small as this would imply

$$\Sigma \otimes c \succ b'' - \epsilon 1_d \succ Y^{-1}A(Y^{-1})^*, \text{ therefore } X \otimes c \succ A.$$

In order to do so, we choose $\epsilon =: 2\delta > 0$ sufficiently small so that the previous equation line holds. Realizing $E \subset \mathcal{B}(H)$ and setting $b_i^\delta = b_i - \delta 1$, we have that $b_i^\delta = [b_i^\delta]_+ - [b_i^\delta]_-$ for some $[b_i^\delta]_+, [b_i^\delta]_- \in \mathcal{B}(H)^+$, $i = 1, 2$. Setting $c' := [b_2^\delta]_+ - [b_1^\delta]_+ \in \mathcal{B}(H)^h$, we can check that

$$b_1 - \epsilon 1 \prec b_1 - \delta 1 \preceq c' \preceq -b_2 + \delta 1 \prec -b_2 + \epsilon 1.$$

Hence by the complete tight Riesz interpolation property there is $c \in E^h$ so that $b_1 - \epsilon 1 \prec c \prec -(b_2 - \epsilon 1)$ verifying the claim. \square

ACKNOWLEDGEMENTS

The author was supported by NSF grant DMS-2055155.

REFERENCES

- [1] Fernando Albiac and Nigel J. Kalton, *Topics in Banach space theory*, Graduate Texts in Mathematics, vol. 233, Springer, New York, 2006. MR2192298
- [2] Erik M. Alfsen, *Compact convex sets and boundary integrals*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 57, Springer-Verlag, New York-Heidelberg, 1971. MR0445271
- [3] Guillaume Aubrun and Stanisław J. Szarek, *Alice and Bob meet Banach*, Mathematical Surveys and Monographs, vol. 223, American Mathematical Society, Providence, RI, 2017. The interface of asymptotic geometric analysis and quantum information theory. MR3699754
- [4] Selçuk Barlak and Gábor Szabó, *Sequentially split $*$ -homomorphisms between C^* -algebras*, Internat. J. Math. **27** (2016), no. 13, 1650105, 48, DOI 10.1142/S0129167X16501056. MR3589655
- [5] Itai Ben Yaacov, Alexander Berenstein, C. Ward Henson, and Alexander Usvyatsov, *Model theory for metric structures*, Model theory with applications to algebra and analysis. Vol. 2, London Math. Soc. Lecture Note Ser., vol. 350, Cambridge Univ. Press, Cambridge, 2008, pp. 315–427, DOI 10.1017/CBO9780511735219.011. MR2436146
- [6] Nathaniel P. Brown and Narutaka Ozawa, *C^* -algebras and finite-dimensional approximations*, Graduate Studies in Mathematics, vol. 88, American Mathematical Society, Providence, RI, 2008. MR2391387
- [7] Man Duen Choi, *Completely positive linear maps on complex matrices*, Linear Algebra Appl. **10** (1975), 285–290, DOI 10.1016/0024-3795(75)90075-0.
- [8] Man Duen Choi and Edward G. Effros, *Separable nuclear C^* -algebras and injectivity*, Duke Math. J. **43** (1976), no. 2, 309–322. MR405117
- [9] Man Duen Choi and Edward Effros, *Injectivity and operator spaces*, Journal of functional analysis **24** (1977), no. 2, 156–209.
- [10] Man Duen Choi and Edward G. Effros, *Nuclear C^* -algebras and injectivity: the general case*, Indiana Univ. Math. J. **26** (1977), no. 3, 443–446, DOI 10.1512/iumj.1977.26.26034. MR430794
- [11] ———, *Nuclear C^* -algebras and the approximation property*, Amer. J. Math. **100** (1978), no. 1, 61–79, DOI 10.2307/2373876. MR482238
- [12] A. Connes, *Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$* , Ann. of Math. (2) **104** (1976), no. 1, 73–115, DOI 10.2307/1971057. MR454659
- [13] John B. Conway, *A course in functional analysis*, 2nd ed., Graduate Texts in Mathematics, vol. 96, Springer-Verlag, New York, 1990. MR1070713
- [14] Edward G. Effros, *Aspects of noncommutative order*, C^* -algebras and applications to physics (Proc. Second Japan-USA Sem., Los Angeles, Calif., 1977), Lecture Notes in Math., vol. 650, Springer, Berlin, 1978, pp. 1–40. MR504750
- [15] Ilijas Farah, Bradd Hart, Martino Lupini, Leonel Robert, Aaron Tikuisis, Alessandro Vignati, and Wilhelm Winter, *Model theory of C^* -algebras*, Mem. Amer. Math. Soc. **271** (2021), no. 1324, viii+127, DOI 10.1090/memo/1324. MR4279915
- [16] Ilijas Farah, Bradd Hart, and David Sherman, *Model theory of operator algebras II: model theory*, Israel J. Math. **201** (2014), no. 1, 477–505, DOI 10.1007/s11856-014-1046-7. MR3265292
- [17] Douglas Farenick and Vern I. Paulsen, *Operator system quotients of matrix algebras and their tensor products*, Math. Scand. **111** (2012), no. 2, 210–243, DOI 10.7146/math.scand.a-15225. MR3023524
- [18] Douglas Farenick, Ali S. Kavruk, and Vern I. Paulsen, *C^* -algebras with the weak expectation property and a multivariable analogue of Ando’s theorem on the numerical radius*, J. Operator Theory **70** (2013), no. 2, 573–590, DOI 10.7900/jot.2011oct07.1938. MR3138370
- [19] Bernd Gärtner and Jiří Matoušek, *Approximation algorithms and semidefinite programming*, Springer, Heidelberg, 2012. MR3015090

- [20] Isaac Goldbring and Thomas Sinclair, *On Kirchberg's embedding problem*, J. Funct. Anal. **269** (2015), no. 1, 155–198, DOI 10.1016/j.jfa.2015.02.016. MR3345606
- [21] ———, *Omitting types in operator systems*, Indiana Univ. Math. J. **66** (2017), no. 3, 821–844, DOI 10.1512/iumj.2017.66.6019. MR3663327
- [22] ———, *On the axiomatizability of C^* -algebras as operator systems*, Glasg. Math. J. **61** (2019), no. 3, 629–635, DOI 10.1017/s001708951800040x. MR3991361
- [23] Alexandre Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Mem. Amer. Math. Soc. **16** (1955), Chapter 1: 196 pp.; Chapter 2: 140 (French). MR75539
- [24] Richard V. Kadison, *A representation theory for commutative topological algebra*, Mem. Amer. Math. Soc. **7** (1951), 39. MR44040
- [25] Ali Ş. Kavruk, *The Weak Expectation Property and Riesz Interpolation*, arXiv e-prints (2012), arXiv:1201.5414, available at 1201.5414.
- [26] ———, *Nuclearity related properties in operator systems*, J. Operator Theory **71** (2014), no. 1, 95–156, DOI 10.7900/jot.2011nov16.1977. MR3173055
- [27] A. Samil Kavruk, *On a non-commutative analogue of a classical result of Namioka and Phelps*, J. Funct. Anal. **269** (2015), no. 10, 3282–3303, DOI 10.1016/j.jfa.2015.09.002. MR3401618
- [28] Ali Kavruk, Vern I. Paulsen, Ivan G. Todorov, and Mark Tomforde, *Tensor products of operator systems*, J. Funct. Anal. **261** (2011), no. 2, 267–299, DOI 10.1016/j.jfa.2011.03.014. MR2793115
- [29] Ali S. Kavruk, Vern I. Paulsen, Ivan G. Todorov, and Mark Tomforde, *Quotients, exactness, and nuclearity in the operator system category*, Adv. Math. **235** (2013), 321–360, DOI 10.1016/j.aim.2012.05.025. MR3010061
- [30] Eberhard Kirchberg, *C^* -nuclearity implies CPAP*, Math. Nachr. **76** (1977), 203–212, DOI 10.1002/mana.19770760115. MR512362
- [31] ———, *On nonsemisplit extensions, tensor products and exactness of group C^* -algebras*, Invent. Math. **112** (1993), no. 3, 449–489, DOI 10.1007/BF01232444. MR1218321
- [32] ———, *Commutants of unitaries in UHF algebras and functorial properties of exactness*, J. Reine Angew. Math. **452** (1994), 39–77, DOI 10.1515/crll.1994.452.39. MR1282196
- [33] Tom-Lukas Kriel, *An introduction to matrix convex sets and free spectrahedra*, Complex Anal. Oper. Theory **13** (2019), no. 7, 3251–3335, DOI 10.1007/s11785-019-00937-8. MR4020034
- [34] L. Lovász, *Semidefinite programs and combinatorial optimization*, Recent advances in algorithms and combinatorics, CMS Books Math./Ouvrages Math. SMC, vol. 11, Springer, New York, 2003, pp. 137–194, DOI 10.1007/0-387-22444-0_6. MR1952986
- [35] Martino Lupini, *An intrinsic order-theoretic characterization of the weak expectation property*, Integral Equations Operator Theory **90** (2018), no. 5, Paper No. 55, 17, DOI 10.1007/s00020-018-2479-x. MR3829543
- [36] Vern Paulsen, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Mathematics, vol. 78, Cambridge University Press, Cambridge, 2002.
- [37] Vern I. Paulsen, Ivan G. Todorov, and Mark Tomforde, *Operator system structures on ordered spaces*, Proc. Lond. Math. Soc. (3) **102** (2011), no. 1, 25–49, DOI 10.1112/plms/pdq011. MR2747723
- [38] Vern I. Paulsen and Mark Tomforde, *Vector spaces with an order unit*, Indiana Univ. Math. J. **58** (2009), no. 3, 1319–1359, DOI 10.1512/iumj.2009.58.3518. MR2542089
- [39] Gilles Pisier, *Introduction to operator space theory*, London Mathematical Society Lecture Note Series, vol. 294, Cambridge University Press, Cambridge, 2003. MR2006539
- [40] ———, *Tensor products of C^* -algebras and operator spaces—the Connes-Kirchberg problem*, London Mathematical Society Student Texts, vol. 96, Cambridge University Press, Cambridge, 2020. MR4283471
- [41] A. G. Robertson and R. R. Smith, *Liftings and extensions of maps on C^* -algebras*, J. Operator Theory **21** (1989), no. 1, 117–131. MR1002124

- [42] Zhong-Jin Ruan, *On matricially normed spaces associated with operator algebras*, University of California, Los Angeles, 1987.
- [43] Thomas Sinclair, *CP-stability and the local lifting property*, New York J. Math. **23** (2017), 739–747. MR3665586
- [44] M. Takesaki, *Theory of operator algebras. I*, Encyclopaedia of Mathematical Sciences, vol. 124, Springer-Verlag, Berlin, 2002. Reprint of the first (1979) edition; Operator Algebras and Non-commutative Geometry, 5. MR1873025
- [45] Corran Webster and Soren Winkler, *The Krein-Milman theorem in operator convexity*, Trans. Amer. Math. Soc. **351** (1999), no. 1, 307–322, DOI 10.1090/S0002-9947-99-02364-8. MR1615970
- [46] Blerina Xhabli, *Universal operator system structures on ordered spaces and their applications*, ProQuest LLC, Ann Arbor, MI, 2009. Thesis (Ph.D.)—University of Houston. MR2736708
- [47] ———, *The super operator system structures and their applications in quantum entanglement theory*, J. Funct. Anal. **262** (2012), no. 4, 1466–1497, DOI 10.1016/j.jfa.2011.11.009. MR2873847

MATHEMATICS DEPARTMENT, PURDUE UNIVERSITY, 150 N. UNIVERSITY STREET, WEST LAFAYETTE, IN 47907-2067

Email address: tsincla@purdue.edu

URL: <http://www.math.purdue.edu/~tsincla/>