

SOLUTIONS TO HOMEWORK 11

1. (a) One has

$$[\sqrt{5}] = 2, \quad 1/(\sqrt{5} - 2) = \sqrt{5} + 2,$$

$$[\sqrt{5} + 2] = 4, \quad 1/((\sqrt{5} + 2) - 4) = 1/(\sqrt{5} - 2) = \sqrt{5} + 2,$$

and we obtain repetition. Thus $\sqrt{5} = [2; \overline{4}]$.

Also, one has

$$[\sqrt{6}] = 2, \quad 1/(\sqrt{6} - 2) = (\sqrt{6} + 2)/2,$$

$$[(\sqrt{6} + 2)/2] = 2, \quad 1/((\sqrt{6} + 2)/2 - 2) = 2/(\sqrt{6} - 2) = \sqrt{6} + 2,$$

$$[\sqrt{6} + 2] = 4, \quad 1/((\sqrt{6} + 2) - 4) = 1/(\sqrt{6} - 2) = (\sqrt{6} + 2)/2,$$

and we obtain repetition. Thus $\sqrt{6} = [2; \overline{2, 4}]$.

(b) One has

$$[\sqrt{54}] = 7, \quad 1/(\sqrt{54} - 7) = (\sqrt{54} + 7)/5,$$

$$[(\sqrt{54} + 7)/5] = 2, \quad 1/((\sqrt{54} + 7)/5 - 2) = 5/(\sqrt{54} - 3) = (\sqrt{54} + 3)/9,$$

$$[(\sqrt{54} + 3)/9] = 1, \quad 1/((\sqrt{54} + 3)/9 - 1) = 9/(\sqrt{54} - 6) = (\sqrt{54} + 6)/2,$$

$$[(\sqrt{54} + 6)/2] = 6, \quad 1/((\sqrt{54} + 6)/2 - 6) = 2/(\sqrt{54} - 6) = (\sqrt{54} + 6)/9,$$

$$[(\sqrt{54} + 6)/9] = 1, \quad 1/((\sqrt{54} + 6)/9 - 1) = 9/(\sqrt{54} - 3) = (\sqrt{54} + 3)/5,$$

$$[(\sqrt{54} + 3)/5] = 2, \quad 1/((\sqrt{54} + 3)/5 - 2) = 5/(\sqrt{54} - 7) = \sqrt{54} + 7,$$

$$[\sqrt{54} + 7] = 14, \quad 1/((\sqrt{54} + 7) - 14) = 1/(\sqrt{54} - 7) = (\sqrt{54} + 7)/5,$$

and we obtain repetition. Thus $\sqrt{54} = [7; \overline{2, 1, 6, 1, 2, 14}]$.

2. One has

$$[\sqrt{69}] = 8, \quad 1/(\sqrt{69} - 8) = (\sqrt{69} + 8)/5,$$

$$[(\sqrt{69} + 8)/5] = 3, \quad 1/((\sqrt{69} + 8)/5 - 3) = 5/(\sqrt{69} - 7) = (\sqrt{69} + 7)/4,$$

$$[(\sqrt{69} + 7)/4] = 3, \quad 1/((\sqrt{69} + 7)/4 - 3) = 4/(\sqrt{69} - 5) = (\sqrt{69} + 5)/11,$$

$$[(\sqrt{69} + 5)/11] = 1, \quad 1/((\sqrt{69} + 5)/11 - 1) = 11/(\sqrt{69} - 6) = (\sqrt{69} + 6)/3,$$

$$[(\sqrt{69} + 6)/3] = 4, \quad 1/((\sqrt{69} + 6)/3 - 4) = 3/(\sqrt{69} - 6) = (\sqrt{69} + 6)/11,$$

$$[(\sqrt{69} + 6)/11] = 1, \quad 1/((\sqrt{69} + 6)/11 - 1) = 11/(\sqrt{69} - 5) = (\sqrt{69} + 5)/4,$$

$$[(\sqrt{69} + 5)/4] = 3, \quad 1/((\sqrt{69} + 5)/4 - 3) = 4/(\sqrt{69} - 7) = (\sqrt{69} + 7)/5,$$

$$[(\sqrt{69} + 7)/5] = 3, \quad 1/((\sqrt{69} + 7)/5 - 3) = 5/(\sqrt{69} - 8) = \sqrt{69} + 8,$$

$$[\sqrt{69} + 8] = 16, \quad 1/(\sqrt{69} - 8) = (\sqrt{69} + 8)/5,$$

and we obtain repetition. Thus $\sqrt{69} = [8; \overline{3, 3, 1, 4, 1, 3, 3, 16}]$.

Also, one has

$$[(24 - \sqrt{15})/7] = 2, \quad 1/((24 - \sqrt{15})/7 - 2) = 7/(10 - \sqrt{15}) = 7(10 + \sqrt{15})/85,$$

$$[7(10 + \sqrt{15})/85] = 1, \quad 1/(7(10 + \sqrt{15})/85 - 1) = 85/(-15 + 7\sqrt{15}) = (15 + 7\sqrt{15})/6,$$

$[(15 + 7\sqrt{15})/6] = 7$, $1/((15 + 7\sqrt{15})/6 - 7) = 6/(-27 + 7\sqrt{15}) = 27 + 7\sqrt{15}$,
 $[27 + 7\sqrt{15}] = 54$, $1/((27 + 7\sqrt{15}) - 54) = 1/(-27 + 7\sqrt{15}) = (27 + 7\sqrt{15})/6$,
 $[(27 + 7\sqrt{15})/6] = 9$, $1/((27 + 7\sqrt{15})/6 - 9) = 6/(-27 + 7\sqrt{15}) = 27 + 7\sqrt{15}$,
 and we obtain repetition. Thus $(24 - \sqrt{15})/7 = [2; 1, 7, \overline{54, 9}]$.

3. Write $\theta = \sum_0^\infty 2023^{-n!}$. For each natural number j , write $q_j = 2023^{j!}$ and

$$a_j = 2023^{j!} \sum_{n=0}^j 2023^{-n!}.$$

Then both a_j and q_j are natural numbers with $(a_j, q_j) = 1$, and

$$|\theta - a_j/q_j| = \sum_{n=j+1}^\infty 2023^{-n!} < 2023^{1-(j+1)!} < q_j^{-j}.$$

If θ were algebraic, then it would be algebraic of some degree $d \geq 1$. By Liouville's theorem, for some positive number c , one would have $|\theta - a/q| > c/q^d$ for every pair of natural numbers a and q with $(a, q) = 1$ and q large enough. But the above upper bound contradicts this lower bound as soon as $j > d$ and j is large enough in terms of c . Hence θ is transcendental.

4. Write $\Theta = \sum_1^\infty 2^{-p_n\#}$. For each natural number j , write $q_j = 2^{p_j\#}$ and

$$a_j = 2^{p_j\#} \sum_{n=1}^j 2^{-p_n\#}.$$

Then both a_j and q_j are natural numbers with $(a_j, q_j) = 1$, and

$$|\Theta - a_j/q_j| = \sum_{n=j+1}^\infty 2^{-p_n\#} < 2^{1-p_{j+1}\#} < q_j^{-j}.$$

Notice here that we use the trivial lower bound $p_{j+1} \geq j + 1$ to derive the last of these inequalities. If Θ were algebraic, then it would be algebraic of some degree $d \geq 1$. By Liouville's theorem, for some positive number c , one would have $|\Theta - a/q| > c/q^d$ for every pair of natural numbers a and q with $(a, q) = 1$ and q large enough. But the above upper bound contradicts this lower bound as soon as $j > d$ and j is large enough in terms of c . Hence Θ is transcendental.

©Trevor D. Wooley, Purdue University 2023. This material is copyright of Trevor D. Wooley at Purdue University unless explicitly stated otherwise. It is provided exclusively for educational purposes at Purdue University, and is to be downloaded or copied for your private study only.