

New estimates for smooth Weyl sums.

Trevor D. Wooley.

1. Introduction.

Since the early part of this century, estimates for Weyl sums (or generalisations thereof) have been central to the treatment of many problems in the additive theory of numbers. For over forty years, the strongest such estimates have stemmed from a method due to Vinogradov [8], the argument having been somewhat simplified recently by the use of the large sieve (see [4, Lemma 5.4]). During this period, improvements in estimates for generalisations of Weyl sums have arisen from improved bounds on mean values of such sums, very recently with the arrival of Vaughan's new iterative method (see [5, Theorems 1.5 and 1.8]). In contrast, this paper will be devoted to improvements at the core of this circle of ideas, within Vinogradov's method itself. Our ideas, which here we shall investigate in the context of smooth Weyl sums, would seem to be applicable elsewhere, and this is a matter which we intend to pursue in the future. We now describe our conclusions in some detail.

Let k be a natural number, and P be a large real number. When $2 \leq R \leq P$, we define the set of R -smooth numbers, $\mathcal{A}(P, R)$, by

$$\mathcal{A}(P, R) = \{n \in [1, P] \cap \mathbb{Z} : p \text{ prime, } p|n \Rightarrow p \leq R\},$$

and for each real number α , we define the corresponding *smooth Weyl sum*, $f(\alpha; P, R)$, by

$$f(\alpha; P, R) = \sum_{x \in \mathcal{A}(P, R)} e(\alpha x^k),$$

where here, and throughout, we write $e(\alpha) = e^{2\pi i \alpha}$. The precise form of our results is to be found in section 4, the following upper bound being a simple corollary.

Theorem 1.1. *Let \mathfrak{m} denote the set of real numbers α such that whenever $a \in \mathbb{Z}$, $q \in \mathbb{N}$, $(a, q) = 1$ and $|\alpha - a/q| \leq q^{-1}P^{1-k}$, one has $q > P$. Then when $\eta = \eta(\varepsilon, k)$ is a sufficiently small positive number, and $2 \leq R \leq P^\eta$, we have*

$$\sup_{\alpha \in \mathfrak{m}} |f(\alpha; P, R)| \ll_{\varepsilon, k} P^{1-\rho(k)+\varepsilon},$$

where, when k is large, $\rho(k)^{-1} = k(\log k + O(\log \log k))$.

1991 Mathematics Subject Classification: 11L07, 11P05.

For comparison, Vaughan [5, Theorem 1.8] obtains a similar result with the exponent satisfying $\rho(k)^{-1} = (4 + o(1))k \log k$, this having been improved, by means of superior mean value estimates, by Wooley [9, Theorem 1.4] to the extent that “4” can be replaced by “2” in the latter conclusion. We note that when $R = P$, the exponential sum $f(\alpha; P, R)$ represents a classical Weyl sum, for which the best estimate corresponding to Theorem 1.1 currently has exponent satisfying $\rho(k)^{-1} = (2 + o(1))k^2 \log k$ (see [10, Corollary 1.1]). Thus, since $\text{card}(\mathcal{A}(P, P^\eta)) \gg_\eta P$ when η is positive, for points in \mathfrak{m} , estimates for smooth Weyl sums are considerably sharper than those for classical Weyl sums.

As far as applications of our new estimate in additive number theory are concerned, we shall restrict ourselves to a cursory consideration of two basic problems. First we improve localised estimates for the fractional parts of αn^k .

Theorem 1.2. *Let $k \in \mathbb{N}$, $\alpha \in \mathbb{R}$ and $\varepsilon > 0$. Then there is a real number $N(\varepsilon, k)$ such that whenever $N \geq N(\varepsilon, k)$,*

$$\min_{1 \leq n \leq N} \|\alpha n^k\| \leq N^{\varepsilon - \tau(k)},$$

where when k is large, $\tau(k)$ satisfies $\tau(k)^{-1} = k(\log k + O(\log \log k))$.

This may be compared with [11, Corollary to Theorem 1.2], where a similar result is established with $\tau(k)^{-1} = 2k(\log k + \log \log k + 2 + o(1))$. Our improvements in Waring’s problem are, unfortunately, rather small. This is because improvements in “minor arc” estimates for exponential sums tend to have a less significant impact, within the circle method, than reduced upper bounds for mean values. As usual, we define $G(k)$ to be the smallest number s such that every sufficiently large natural number is the sum of, at most, s k th powers of natural numbers. The precise form of our new estimate (see Corollary 1 to Theorem 4.2) leads to the following bounds for $G(k)$ when $10 \leq k \leq 20$.

Theorem 1.3. *We have $G(10) \leq 62$, $G(12) \leq 78$, $G(13) \leq 86$, $G(14) \leq 94$, $G(15) \leq 102$, $G(16) \leq 110$, $G(17) \leq 118$, $G(18) \leq 127$, $G(19) \leq 135$, $G(20) \leq 144$.*

For comparison, the respective bounds $G(10) \leq 63$, $G(12) \leq 79$, $G(13) \leq 87$, $G(14) \leq 95$, $G(15) \leq 103$, $G(16) \leq 112$, $G(17) \leq 120$, $G(18) \leq 129$, $G(19) \leq 138$, $G(20) \leq 146$ were obtained in [9, Theorem 1.1]. When $k = 11$, our improvements are not sufficiently large to lead to a visible reduction in $G(k)$. For large k our methods now lead to the following upper bound on $G(k)$.

Theorem 1.4. *We have*

$$G(k) \leq k \left(\log k + \log \log k + 2 + O \left(\frac{\log \log k}{\log k} \right) \right).$$

This conclusion may be compared with [11, Theorem 1.3], where a similar result is given with “ $2 + \log 2$ ” replacing “2”.

Our estimate for $f(\alpha; P, R)$ is based on an application of the large sieve inequality, in essence, at least in the initial stages of the argument, following the treatment of Vaughan [5, section 10]. We let M be a real number with $P^{1/2} \leq M \leq P$ to be chosen later. By Dirichlet’s Theorem, we may find $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ with $(a, q) = 1$, $|q\alpha - a| \leq \frac{1}{2}(MR)^{-k}$ and $q \leq 2(MR)^k$. Then, by a suitable combinatorial lemma combined with Hölder’s inequality, when $s \in \mathbb{N}$ we are able to bound the smooth Weyl sum $f(\alpha; P, R)$ in the form

$$f(\alpha; P, R)^{2s} \ll M^{2s+\varepsilon} + (MR)^{2s-1} \sum_{\substack{M < v \leq MR \\ (v, q) = 1}} \left| \sum_{1 \leq y \leq s(P/M)^k} b_y e(\alpha v^k y) \right|^2, \quad (1.1)$$

where b_y denotes the number of solutions of the diophantine equation

$$u_1^k + \dots + u_s^k = y,$$

with $u_i \in \mathcal{A}(P/M, R)$ ($1 \leq i \leq s$). We may classify the values of v into $O(P^\varepsilon)$ sets $\mathcal{V}_1, \dots, \mathcal{V}_L$ so that $v_1, v_2 \in \mathcal{V}_j$ and $v_1^k \equiv v_2^k \pmod{q}$ together imply that $v_1 \equiv v_2 \pmod{q}$. Thus, provided that M is chosen suitably, there is a j satisfying

$$f(\alpha; P, R)^{2s} \ll (MR)^{2s-1+\varepsilon} \sum_{v \in \mathcal{V}_j} \left| \sum_{1 \leq y \leq s(P/M)^k} b_y e(\alpha v^k y) \right|^2.$$

When α lies in a set of type similar to the set \mathfrak{m} defined in the statement of Theorem 1.1, an analysis of the αv^k shows that they are spaced at least $(2q)^{-1}$ apart modulo 1. Then by the large sieve inequality,

$$f(\alpha; P, R)^{2s} \ll (MR)^{2s-1+\varepsilon} (q + s(P/M)^k) \sum_{1 \leq y \leq Y} b_y^2,$$

the last sum being bounded by using a suitable mean value estimate for smooth Weyl sums. The strength of the ensuing bound for $f(\alpha; P, R)$ now depends on the relative magnitudes of M and $q + (P/M)^k$, our estimate improving as the former parameter grows and the latter expression decreases. Vaughan takes $M = P^{1/2}$, which leads to the bound $q + (P/M)^k \ll P^{k/2}$. We take $M = P^\lambda$ with λ a parameter satisfying $\frac{1}{2} < \lambda < 1$. As it stands, it is possible that q is as large as $P^{k\lambda}$, and this would lead to weak bounds on $f(\alpha; P, R)$. However, by modifying an argument of Heath-Brown [2, section 5], it is possible to obtain a complementary bound on $f(\alpha; P, R)$ which is useful only when q is

large. Thus we are able to discard those q with $q > P^{k(1-\lambda)}$, and thereby achieve a suitable bound on $q + (P/M)^k$.

The above argument is effective for the set of α satisfying the property that whenever $a \in \mathbb{Z}$, $q \in \mathbb{N}$, $(a, q) = 1$ and $|\alpha - a/q| \leq q^{-1}MP^{-k}$, one has $q > M$. In Vaughan's analysis one is constrained to take $M \leq P^{1/2}$, which limits the strength of the ensuing bounds. The basic advantage of our new method is the relaxation of this condition, and indeed for large k we may now take $M = P^{1+o(1)}$. We note that Thanigasalam [3] has obtained a variant of Vinogradov's method which can be applied to smooth exponential sums (see [4, section 9]). The latter method permits one to take $M = P^{\frac{k}{2k-1}} = P^{\frac{1}{2}+O(1/k)}$, so that it is asymptotically of no greater strength than that due to Vaughan.

In section 2 we record some basic estimates for mean values of smooth Weyl sums, and also provide a suitable combinatorial lemma for our later arguments. Section 3 is devoted to the task of establishing an asymmetric estimate for $f(\alpha; P, R)$ of value for large moduli q , which in combination with a suitable estimate for small moduli in section 4, leads to the desired estimate for $f(\alpha; P, R)$. In sections 5 and 6 we then draw corollaries concerning Waring's problem and the fractional parts of polynomials.

The author thanks the referee for several useful comments which have improved the exposition of this paper.

2. Preliminary observations.

We start by recalling some of the salient features of the new iterative method in Waring's problem. Throughout, s , t and u will denote positive integers, and ε and η will denote sufficiently small positive numbers. We take P to be a large positive real number depending at most on k , s , t , u , ε and η . We use \ll and \gg to denote Vinogradov's well-known notation, implicit constants depending at most on the latter numbers. Also, we write $[x]$ for the greatest integer not exceeding x , and write $\|x\|$ for $\min_{y \in \mathbb{Z}} |x - y|$. In order to simplify our analysis, we adopt the following convention concerning the numbers ε and R . Whenever ε or R appear in a statement, either implicitly or explicitly, we assert that for each $\varepsilon > 0$, there exists a positive number $\eta_0 = \eta_0(\varepsilon, s, t, u, k)$ such that the statement holds whenever $R = P^\eta$, with $0 < \eta \leq \eta_0$. Note that the "value" of ε , and η_0 , may change from statement to statement, and hence also the dependency of implicit constants on ε and η .

We define $S_s(P, R)$ to be the number of solutions of the diophantine equation

$$x_1^k + \dots + x_s^k = y_1^k + \dots + y_s^k,$$

with $x_i, y_i \in \mathcal{A}(P, R)$ ($1 \leq i \leq s$). Thus

$$S_s(P, R) = \int_0^1 |f(\alpha; P, R)|^{2s} d\alpha.$$

We shall say that an exponent $\Delta_s = \Delta_{s,k}$ is *permissible* whenever the exponent has the property that $S_s(P, R) \ll P^{\lambda_{s,k} + \varepsilon}$, with $\lambda_{s,k} = 2s - k + \Delta_{s,k}$. It follows easily that any permissible exponent $\Delta_{s,k}$ is non-negative, and moreover, without loss of generality, that $\Delta_{s,k} \leq k$. The following lemma provides us with permissible exponents when k is large.

Lemma 2.1. *Let $k \geq 4$ and $t \in \mathbb{N}$. For each $s \in \mathbb{N}$ with $2 \leq s \leq t$, define the real number $\Delta_s = \Delta_{s,k}$ to be the unique positive solution of the equation*

$$\Delta_s e^{\Delta_s/k} = k e^{1-2s/k}.$$

Then $\Delta_{s,k}$ is permissible, and consequently the exponent $\Delta_{s,k}^ = k e^{1-2s/k}$ is permissible.*

Proof. This is the corollary to [11, Theorem 2.1], which simplifies [9, Lemma 3.2].

We shall require a result on the density of integers with a given square-free kernel. Given an integer v with canonical prime factorisation $\prod_{i=1}^t p_i^{r_i}$, we denote by $s_0(v)$ the square-free kernel of v , that is $\prod_{i=1}^t p_i$. Furthermore, we define the set $\mathcal{C}_q(Q)$ by

$$\mathcal{C}_q(Q) = \{x \in \mathbb{Z} \cap [1, Q] : s_0(x) | s_0(q)\}.$$

Lemma 2.2. *Suppose that L is a positive real number and r is a positive integer with $\log r \ll \log L$. Then for each $\varepsilon > 0$,*

$$\text{card}(\mathcal{C}_r(L)) \ll L^\varepsilon.$$

Proof. For each divisor d of r , it follows from [9, Lemma 2.1] that

$$\text{card} \{1 \leq y \leq L : s_0(y) = s_0(d)\} \ll L^\varepsilon.$$

The lemma now follows by using standard estimates for the divisor function.

We conclude this section by providing a means of decomposing the smooth Weyl sum $f(\alpha; P, R)$ into a form in which we can apply the arguments of sections 3 and 4. For this purpose we apply essentially the same argument as that used by Vaughan [5, p67].

Lemma 2.3. *Let $\alpha \in \mathbb{R}$ and $r \in \mathbb{N}$, and suppose that Q , M and R satisfy $2 \leq R \leq M < Q$. Then*

$$\sum_{\substack{x \in \mathcal{A}(Q, R) \\ (x, r) = 1}} e(\alpha x^k) \ll R \log Q \max_{\substack{\pi \leq R \\ \pi \text{ prime}}} \sup_{\theta \in [0, 1)} V_r(\alpha; Q, M, R; \pi, \theta) + M,$$

where

$$V_r(\alpha; Q, M, R; \pi, \theta) = \sum_{\substack{v \in \mathcal{B}(M, \pi, R) \\ (v, r) = 1}} \left| \sum_{\substack{u \in \mathcal{A}(Q/M, \pi) \\ (u, r) = 1}} e(\alpha(uv)^k + \theta u) \right|,$$

and

$$\mathcal{B}(M, \pi, R) = \{v \in \mathbb{N} : M < v \leq M\pi, \pi | v, \text{ and } p \text{ prime, } p | v \Rightarrow \pi \leq p \leq R\}.$$

Proof. On following the argument leading to equation (10.9) of Vaughan [5], it is apparent that the conclusion remains valid with first summation in equation (10.10) adjusted so that the condition $M < v \leq MR$ is replaced by $v \in \mathcal{B}(M, p, R)$. The lemma therefore follows immediately from equations (10.4), (10.6), (10.9) and (10.10) of Vaughan [5].

3. Upper bounds for large moduli.

When α is close to a rational a/q with q large, we use an asymmetric form of an argument similar to one used by Heath-Brown [2, section 5] in the estimation of the fractional part of αn^k .

Lemma 3.1. *Suppose that λ satisfies $\frac{1}{2} < \lambda < 1$, and write $M = P^\lambda$. Let $\alpha \in \mathbb{R}$, and suppose that there exist $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ satisfying $(a, q) = 1$ and $|\alpha - a/q| \leq q^{-2}$. Then when $t, w \in \mathbb{N}$, and Δ_t and Δ_w are permissible,*

$$f(\alpha; P, R) \ll q^\varepsilon P^{1+\varepsilon} (M^{\Delta_w} (P/M)^{\Delta_t} (q^{-1} + M^{-k} + (P/M)^{-k} + qP^{-k}))^{\frac{1}{2tw}} + M.$$

Proof. By applying Lemma 2.3 with $r = 1$, we deduce that there exists a prime π with $\pi \leq R$, and $\theta \in [0, 1)$ such that

$$f(\alpha; P, R) \ll P^\varepsilon R \sum_{v \in \mathcal{A}(MR, R)} |h(\alpha; v, \theta)| + M, \quad (3.1)$$

where

$$h(\alpha; v, \theta) = \sum_{u \in \mathcal{A}(P/M, \pi)} e(\alpha(uv)^k + \theta u).$$

Define the complex numbers of unit modulus, $\varepsilon(v, \theta)$, by

$$|h(\alpha; v, \theta)|^t = \varepsilon(v, \theta) h(\alpha; v, \theta)^t. \quad (3.2)$$

Also, let r_c denote the number of solutions of the diophantine equation

$$u_1^k + \dots + u_t^k = c, \quad (3.3)$$

with $u_i \in \mathcal{A}(P/M, \pi)$, in which each solution \mathbf{u} is counted with weight $e(\theta(u_1 + \dots + u_t))$.

Thus

$$h(\alpha; v, \theta)^t = \sum_{1 \leq c \leq t(P/M)^k} r_c e(\alpha c v^k),$$

and hence an application of Hölder's inequality, together with (3.2), yields

$$\begin{aligned} \left(\sum_{v \in \mathcal{A}(MR, R)} |h(\alpha; v, \theta)| \right)^t &\leq (MR)^{t-1} \sum_{v \in \mathcal{A}(MR, R)} |h(\alpha; v, \theta)|^t \\ &= (MR)^{t-1} \sum_{1 \leq c \leq t(P/M)^k} r_c \sum_{v \in \mathcal{A}(MR, R)} \varepsilon(v, \theta) e(\alpha c v^k). \end{aligned}$$

Next we note that $|r_c| \leq n_c$, where n_c denotes the number of solutions of the equation (3.3) with $u_i \in \mathcal{A}(P/M, \pi)$. A further application of Hölder's inequality therefore shows that

$$\left(\sum_{v \in \mathcal{A}(MR, R)} |h(\alpha; v, \theta)| \right)^{2tw} \leq (MR)^{2w(t-1)} \left(\sum_c n_c \right)^{2w-2} \left(\sum_c n_c^2 \right) J_w(\alpha),$$

where

$$J_w(\alpha) = \sum_{1 \leq c \leq t(P/M)^k} |g(\alpha; c, \theta)|^{2w}, \quad (3.4)$$

and

$$g(\alpha; c, \theta) = \sum_{v \in \mathcal{A}(MR, R)} \varepsilon(v, \theta) e(\alpha c v^k).$$

But by considering the underlying diophantine equations, we have $\sum_c n_c \leq (P/M)^t$ and $\sum_c n_c^2 \leq S_t(P/M, R)$, and hence

$$\left(\sum_{v \in \mathcal{A}(MR, R)} |h(\alpha; v, \theta)| \right)^{2tw} \leq (PR)^{2tw} (P/M)^{-2t} (MR)^{-2w} J_w(\alpha) S_t(P/M, R). \quad (3.5)$$

Let \tilde{n}_d denote the number of solutions of the equation

$$\sum_{i=1}^w v_i^k - \sum_{i=w+1}^{2w} v_i^k = d,$$

with $v_i \in \mathcal{A}(MR, R)$ ($1 \leq i \leq 2w$), each solution \mathbf{v} counted with weight

$$\prod_{i=1}^w \varepsilon(v_i, \theta) \overline{\varepsilon(v_{w+i}, \theta)}.$$

Then by considering the underlying diophantine equation,

$$\tilde{n}_d = \int_0^1 |g(\beta; c, \theta)|^{2w} e(-\beta d) d\beta.$$

Therefore, on making a trivial estimate, $|\tilde{n}_d| \leq \tilde{n}_0 \leq S_w(MR, R)$, since $|\varepsilon(v, \theta)| = 1$. On recalling equation (3.4), we have

$$\begin{aligned} J_w(\alpha) &= \sum_{1 \leq c \leq t(P/M)^k} \sum_{|d| \leq w(MR)^k} \tilde{n}_d e(\alpha cd) \\ &\leq S_w(MR, R) \sum_{|d| \leq w(MR)^k} \left| \sum_{1 \leq c \leq t(P/M)^k} e(\alpha cd) \right| \\ &\ll S_w(MR, R) \Upsilon(q), \end{aligned} \quad (3.6)$$

where

$$\Upsilon(q) = \sum_{|d| \leq w(MR)^k} \min \{ (P/M)^k, \|\alpha d\|^{-1} \}.$$

But by using [4, Lemma 2.2] we obtain

$$\Upsilon(q) \ll (PR)^{k+\varepsilon} q^\varepsilon (q^{-1} + (P/M)^{-k} + M^{-k} + qP^{-k}).$$

Therefore, by (3.5), (3.6), and recalling the discussion of $S_s(P, R)$ in section 2, we deduce that

$$\begin{aligned} \sum_{v \in \mathcal{A}(MR, R)} |h(\alpha; v, \theta)| &\ll ((PR)^{2tw+\varepsilon} (P/M)^{-k+\Delta_t} (MR)^{-k+\Delta_w} \Upsilon(q))^{\frac{1}{2tw}} \\ &\ll q^\varepsilon P^{1+\varepsilon} ((P/M)^{\Delta_t} M^{\Delta_w} (q^{-1} + (P/M)^{-k} + M^{-k} + qP^{-k}))^{\frac{1}{2tw}}. \end{aligned}$$

The lemma now follows immediately from (3.1).

4. Upper bounds for small moduli.

When α is close to a rational a/q with q small, we are able to adapt a variant of Vinogradov's method given by Vaughan [5, section 10] to provide an upper bound for $f(\alpha; P, R)$. Our proof will differ in detail from that of Vaughan, since we are able to provide some technical simplifications which lead to a more precise result.

Lemma 4.1. *Suppose that $\frac{1}{2} < \lambda < 1$, and write $M = P^\lambda$. Let $\alpha \in \mathbb{R}$, and suppose that $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ satisfy $(a, q) = 1$, $|q\alpha - a| \leq \frac{1}{2}(MR)^{-k}$, $q \leq 2(MR)^k$ and either $|q\alpha - a| > MP^{-k}$ or $q > MR$. Then if s is a natural number with $2s \geq k + 1$, and Δ_s is permissible,*

$$f(\alpha; P, R) \ll M^{1+\varepsilon} + P^{1+\varepsilon} (M^{-1}(P/M)^{\Delta_s} (1 + q(P/M)^{-k}))^{1/2s}.$$

Proof. Observe that

$$f(\alpha; P, R) = \sum_{x \in \mathcal{A}(P, R)} e(\alpha x^k) = \sum_{d \in \mathcal{C}_q(P) \cap \mathcal{A}(P, R)} \sum_{\substack{x \in \mathcal{A}(P/d, R) \\ (x, q) = 1}} e(\alpha (xd)^k).$$

Thus, on applying Lemma 2.2,

$$\begin{aligned} f(\alpha; P, R) &\ll \sum_{d \in \mathcal{C}_q(P/M)} \left| \sum_{\substack{x \in \mathcal{A}(P/d, R) \\ (x, q) = 1}} e(\alpha (xd)^k) \right| + \sum_{\substack{d \in \mathcal{C}_q(P) \\ d > P/M}} P/d \\ &\ll P^\varepsilon \max_{d \in \mathcal{C}_q(P/M)} \left| \sum_{\substack{x \in \mathcal{A}(P/d, R) \\ (x, q) = 1}} e(\alpha (xd)^k) \right| + M^{1+\varepsilon}. \end{aligned}$$

Next, by Lemma 2.3, it follows that for some $d \in \mathcal{C}_q(P/M)$, $\theta \in [0, 1)$, and prime $\pi \leq R$, we have

$$f(\alpha; P, R) \ll M^{1+\varepsilon} + P^\varepsilon Rg(\alpha; d, \pi, \theta), \quad (4.1)$$

where

$$g(\alpha; d, \pi, \theta) = \sum_{\substack{v \in \mathcal{B}(M/d, \pi, R) \\ (v, q) = 1}} \left| \sum_{\substack{u \in \mathcal{A}(P/M, \pi) \\ (u, q) = 1}} e(\alpha(uvd)^k + \theta u) \right|.$$

Notice here that $M/d \geq M^2/P = P^{2\lambda-1}$, so that M/d is at least as large as some positive power of P .

Let $J(q, d, h)$ denote the number of solutions of the congruence $(xd)^k \equiv h \pmod{q}$ with $1 \leq x \leq q$. Then when $(h, q) | d^k$, a simple estimate gives $J(q, d, h) \ll q^\varepsilon d^k$. Hence there is an $L \ll q^\varepsilon d^k$ such that the v with $M/d < v \leq MR/d$ and $(v, q) = 1$ can be divided into L classes $\mathcal{V}_1, \dots, \mathcal{V}_L$ such that for any two distinct elements v_1, v_2 in a given class \mathcal{V}_j , we have $(v_1d)^k \equiv (v_2d)^k \pmod{q}$ if and only if $v_1 \equiv v_2 \pmod{q}$. On writing c_y for the number of solutions of the diophantine equation

$$u_1^k + \dots + u_s^k = y$$

with $u_i \in \mathcal{A}(P/M, \pi)$, we may apply Hölder's inequality to obtain

$$g(\alpha; d, \pi, \theta)^{2s} \ll P^\varepsilon d^k (MR/d)^{2s-1} \max_{1 \leq j \leq L} \sum_{v \in \mathcal{V}_j} \left| \sum_{1 \leq y \leq s(P/M)^k} b_y e(\alpha(vd)^k y) \right|^2, \quad (4.2)$$

where $|b_y| \leq c_y$.

Now recall that $|q\alpha - a| \leq \frac{1}{2}(MR)^{-k}$. Thus, if $v_1, v_2 \in \mathcal{V}_j$ and $v_1 \not\equiv v_2 \pmod{q}$, then we have $(v_1d)^k \not\equiv (v_2d)^k \pmod{q}$, and hence

$$\|\alpha((v_1d)^k - (v_2d)^k)\| \geq \left\| \frac{a}{q}((v_1d)^k - (v_2d)^k) \right\| - \frac{1}{2}q^{-1} \geq \frac{1}{2}q^{-1}. \quad (4.3)$$

We divide into cases.

(i) Suppose that $q > MR/d$. Then the elements of \mathcal{V}_j are distinct mod q , and hence it follows from (4.3) that for $v \in \mathcal{V}_j$, the $\alpha(vd)^k$ are spaced at least $\frac{1}{2}q^{-1}$ apart modulo 1.

(ii) Suppose that $q \leq MR/d$. Then since $q \leq MR$, by hypothesis we have $|q\alpha - a| > MP^{-k}$. Given any two distinct elements v_1, v_2 of \mathcal{V}_j with $v_1 \not\equiv v_2 \pmod{q}$, we may conclude as in case (i) that $\alpha(v_1d)^k$ and $\alpha(v_2d)^k$ are spaced at least $\frac{1}{2}q^{-1}$ apart modulo 1. Thus we are left to consider the situation in which $v_1 \equiv v_2 \pmod{q}$, but $v_1 \neq v_2$. Then

$$\|\alpha((v_1d)^k - (v_2d)^k)\| = \|(\alpha - a/q)((v_1d)^k - (v_2d)^k)\| = |\alpha - a/q| \cdot |(v_1d)^k - (v_2d)^k|.$$

Since $v_1 - v_2$ is a non-zero multiple of q , and $v_1 d > M$ and $v_2 d > M$, we have

$$\|\alpha((v_1 d)^k - (v_2 d)^k)\| \geq |\alpha - a/q| M^{k-1} q > (P/M)^{-k}.$$

Therefore, in this case, for $v \in \mathcal{V}_j$ the $\alpha(vd)^k$ are spaced at least $\frac{1}{2} \min\{q^{-1}, (P/M)^{-k}\}$ apart modulo 1.

Then in either case, by the large sieve inequality (see, for example, [4, Lemma 5.3]) we have

$$\begin{aligned} \sum_{v \in \mathcal{V}_j} \left| \sum_{1 \leq y \leq s(P/M)^k} b_y e(\alpha(vd)^k y) \right|^2 &\ll (q + (P/M)^k) \sum_{1 \leq y \leq s(P/M)^k} |b_y|^2 \\ &\ll (q + (P/M)^k) S_s(P/M, R). \end{aligned}$$

Then since $2s \geq k + 1$, it follows from (4.1) and (4.2) that

$$f(\alpha; P, R) \ll M^{1+\varepsilon} + (P^\varepsilon (MR)^{2s-1} (P/M)^{2s-k+\Delta_s} (q + (P/M)^k))^{1/2s}.$$

The lemma now follows immediately.

Theorem 4.2. *Suppose that $\frac{1}{2} < \lambda < 1$. Let $\alpha \in \mathbb{R}$, and suppose that whenever $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ satisfy $(a, q) = 1$ and $|\alpha - a/q| \leq q^{-1} P^{\lambda-k}$, then one has $q > P^\lambda R$. Then if $s, t, w \in \mathbb{N}$ satisfy $2s \geq k + 1$, and Δ_s, Δ_t and Δ_w are permissible exponents,*

$$f(\alpha; P, R) \ll P^\varepsilon \left(P^\lambda + P^{1-\mu(k)} + P^{1-\nu(k)} \right),$$

where

$$\mu(k) = \frac{k(1-\lambda) - \lambda\Delta_w - (1-\lambda)\Delta_t}{2tw}$$

and

$$\nu(k) = \frac{\lambda - (1-\lambda)\Delta_s}{2s}.$$

Proof. Write $M = P^\lambda$. By Dirichlet's Theorem, there exist $a \in \mathbb{Z}$ and $q \in \mathbb{N}$ with $(a, q) = 1$, $q \leq 2(MR)^k$ and $|q\alpha - a| \leq \frac{1}{2}(MR)^{-k}$. If $q > (P/M)^k$, then we may apply Lemma 3.1 to obtain the bound

$$\begin{aligned} f(\alpha; P, R) &\ll P^{1+\varepsilon} \left(P^{\lambda\Delta_w + (1-\lambda)\Delta_t} \left(P^{-k(1-\lambda)} + P^{-k\lambda} \right) \right)^{\frac{1}{2tw}} + P^\lambda \\ &\ll P^{\lambda+\varepsilon} + P^{1-\mu(k)+\varepsilon}. \end{aligned}$$

Thus we may assume that $q \leq (P/M)^k$. In this second case we apply Lemma 4.1 to establish the estimate

$$\begin{aligned} f(\alpha; P, R) &\ll P^{\lambda+\varepsilon} + P^{1+\varepsilon} \left(P^{-\lambda+(1-\lambda)\Delta_s} \right)^{1/2s} \\ &\ll P^{\lambda+\varepsilon} + P^{1-\nu(k)+\varepsilon}. \end{aligned}$$

This completes the proof of the theorem.

By simply choosing λ optimally in Theorem 4.2, we obtain the following conclusion.

Corollary 1. *Let s, t, w be natural numbers satisfying $2s \geq k + 1$, and suppose that Δ_n ($n = s, t, w$) are permissible exponents. Define*

$$\sigma(k) = \frac{k - \Delta_t - \Delta_s \Delta_w}{2(s(k + \Delta_w - \Delta_t) + tw(1 + \Delta_s))},$$

and

$$\lambda(k) = \frac{s(k - \Delta_t) + tw\Delta_s}{s(k + \Delta_w - \Delta_t) + tw(1 + \Delta_s)}.$$

Suppose that $\frac{1}{2} < \lambda(k) < 1 - \sigma(k)$. Then when α satisfies the hypotheses of Theorem 4.2,

$$f(\alpha; P, R) \ll P^{1-\sigma(k)+\varepsilon}.$$

We now explore the consequences of our new estimate when k is large.

Corollary 2. *Let \mathfrak{m}_λ denote the set of $\alpha \in \mathbb{R}$ such that whenever $a \in \mathbb{Z}$, $q \in \mathbb{N}$, $(a, q) = 1$ and $|\alpha - a/q| \leq q^{-1}P^{\lambda-k}$, then $q > P^\lambda R$. Then there is a natural number $k_0(\varepsilon)$ with the following property. When $k \geq k_0(\varepsilon)$, there are real numbers $\lambda = \lambda(k)$ and $\sigma(k)$ with*

$$\frac{\log \log k}{\log k} \ll 1 - \lambda \ll \frac{\log \log k}{\log k},$$

and

$$\sigma(k)^{-1} = k(\log k + O(\log \log k)),$$

and such that

$$\sup_{\alpha \in \mathfrak{m}_\lambda} |f(\alpha; P, R)| \ll P^{1-\sigma(k)+\varepsilon}.$$

Proof. We put

$$s = \left[\frac{1}{2} k \log k \right], \quad t = \left[\frac{k \log \log k}{\log k} \right], \quad w = [k(\log \log k - \log \log \log k)].$$

Then on solving the equation $\xi e^\xi = e^{1-2t/k}$, we find from Lemma 2.1 that the exponent Δ_t is permissible, where

$$\frac{\Delta_t}{k} = 1 - \frac{t}{k} + O\left(\frac{t^2}{k^2}\right).$$

Hence

$$k - \Delta_t = k \frac{\log \log k}{\log k} \left(1 + O\left(\frac{\log \log k}{\log k}\right)\right).$$

Similarly, the exponents Δ_w and Δ_s are permissible, where by the concluding remark of Lemma 2.1,

$$\Delta_w = ke^{1-2w/k} \ll k \left(\frac{\log \log k}{\log k}\right)^2,$$

and

$$\Delta_s \leq ke^{1-2s/k} \ll 1.$$

We now note that, in the notation of Corollary 1 to Theorem 4.2,

$$\sigma(k)^{-1} = 2s + \frac{2(s\Delta_w + tw)(1 + \Delta_s)}{k - \Delta_t - \Delta_s\Delta_w}.$$

Thus

$$\sigma(k)^{-1} - k \log k \ll k \log \log k. \quad (4.4)$$

Furthermore, again in the notation of Corollary 1 to Theorem 4.2,

$$\lambda = 1 - \frac{s\Delta_w + tw}{s(k + \Delta_w - \Delta_t) + tw(1 + \Delta_s)}.$$

Thus

$$\frac{\log \log k}{\log k} \ll 1 - \lambda \ll \frac{\log \log k}{\log k},$$

and so, in particular, when k is sufficiently large we have $\frac{1}{2} < \lambda < 1 - \sigma(k)$, and so the hypotheses of Corollary 1 to Theorem 4.2 are satisfied. Therefore, in view of (4.4), the corollary follows immediately from Corollary 1 to Theorem 4.2.

We note that Theorem 1.1 is merely a simplification of Corollary 2 to Theorem 4.2.

5. The estimation of $G(k)$.

The estimation of $G(k)$ is now relatively routine. It should be noted that our “minor” arcs will be slightly different from those used in previous analyses (see, in particular, [6, section 9]). Thus we shall take this opportunity to record the new results stemming from our analysis in the form of the following theorem.

Theorem 5.1. *Let $\lambda(k)$ and $\sigma(k)$ be the real numbers defined in Corollary 1 to Theorem 4.2. Under the same hypotheses as in that corollary, for each integer v with $v \geq k - 1$, and each permissible exponent Δ_v ,*

$$G(k) \leq 2v + 3 + \left\lceil \frac{\Delta_{v+1}}{\sigma(k)} \right\rceil.$$

Proof. Let n be a large positive number, and write $P = \lceil n^{1/k} \rceil$. Let $v, w \in \mathbb{N}$ satisfy $v \geq k - 1$, and write $s = 2v + w$. We consider the number of representations, $R(n)$, of n in the form

$$n = x_1^k + x_2^k + y_1^k + \dots + y_s^k,$$

with $x_1, x_2 \in \mathbb{Z} \cap [1, P]$ and $y_i \in \mathcal{A}(P, R)$ ($1 \leq i \leq s$). On writing

$$g(\alpha) = \sum_{1 \leq x \leq P} e(\alpha x^k),$$

we therefore have

$$R(n) = \int_0^1 g(\alpha)^2 f(\alpha; P, R)^s e(-\alpha n) d\alpha.$$

Let \mathfrak{m} denote the set of real numbers $\alpha \in [0, 1)$ with the property that, whenever $a \in \mathbb{Z}$, $q \in \mathbb{N}$, $(a, q) = 1$ and $|\alpha - a/q| \leq q^{-1}P^{1-k}$, one has $q > P$. Then by Corollary 1 to Theorem 4.2, together with the remark at the end of section 3 of [9], it follows that

$$\begin{aligned} \int_{\mathfrak{m}} g(\alpha)^2 f(\alpha; P, R)^{2v+w} e(-\alpha n) d\alpha &\ll \left(\sup_{\alpha \in \mathfrak{m}} |f(\alpha; P, R)| \right)^w \int_0^1 |g(\alpha)^2 f(\alpha; P, R)^{2v}| d\alpha \\ &\ll \left(P^{1-\sigma(k)+\varepsilon} \right)^w P^{2v+2-k+\Delta_{v+1}+\varepsilon}. \end{aligned}$$

Thus, provided that $w\sigma(k) > \Delta_{v+1}$, then

$$\int_{\mathfrak{m}} g(\alpha)^2 f(\alpha; P, R)^{2v+w} e(-\alpha n) d\alpha \ll P^{s+2-k-\delta}$$

for some positive number δ . The major arcs $\mathfrak{M} = [0, 1) \setminus \mathfrak{m}$ may be dealt with by means of the same pruning argument as was used in [5, section 5], owing to the presence of the factor $g(\alpha)^2$ (consisting of exponential sums over complete intervals). Thereby, one may obtain

$$\int_{\mathfrak{M}} g(\alpha)^2 f(\alpha; P, R)^{2v+w} e(-\alpha n) d\alpha \gg P^{s+2-k},$$

and the desired conclusion follows immediately.

In order to prove Theorem 1.4, we have merely to apply Theorem 5.1 with $v = [\frac{1}{2}k(\log k + \log \log k + 1)]$. From the concluding remark of Lemma 2.1, the exponent Δ_{v+1} is permissible, where

$$\Delta_{v+1} \leq k e^{1-2(v+1)/k} \leq 1/\log k.$$

Meanwhile, from Theorem 1.1 we have $\sigma(k)^{-1} = k \log k + O(k \log \log k)$. Thus, in Theorem 5.1 we deduce that

$$G(k) \leq k(\log k + \log \log k + 1) + k + O\left(\frac{k \log \log k}{\log k}\right).$$

For Theorem 1.3 we must work a little harder. We calculate $\sigma(k)$ by applying Corollary 1 to Theorem 4.2, using the values of Δ_s , Δ_t and Δ_w listed in the table in the Appendix. The latter values of Δ_n may be shown, after some calculations, to be permissible by using the methods of [9] (we should point out that superior estimates should arise, albeit after much greater effort, from the methods of [7]). Having checked that the corresponding value of $\lambda(k)$, which is also listed in the table, satisfies the necessary hypotheses, we then apply Theorem 5.1 using the value of v indicated in the Appendix. The estimates for $G(k)$ detailed in Theorem 1.3 then follow immediately.

6. Localised estimates for fractional parts of polynomials.

We can dispose of the proof of Theorem 1.2 swiftly with a standard appeal to [1, Lemma 5]. We take $\lambda = \lambda(k)$ and $\sigma(k)$ to be as defined in the statement of Corollary 2 to Theorem 4.2. We let ϕ be any real number with $\varepsilon < \phi < \sigma(k)$. Let P be a large positive number, and put $H = P^{\sigma(k)-\phi}$. Define $T(\alpha)$ by

$$T(\alpha) = \sum_{1 \leq h \leq H} |f(h\alpha; P, R)|.$$

Then provided that we can establish the bound $T(\alpha) = o(P)$, by [1, Lemma 5], as in [11, section 5], it follows that

$$\min_{1 \leq n \leq P} \|\alpha n^k\| < P^{\phi-k}.$$

Suppose first that there is a triple h, a, q with $1 \leq h \leq H$, $a \in \mathbb{Z}$, $q \in \mathbb{N}$, $(a, q) = 1$, $|qh\alpha - a| \leq P^{\lambda-k}$ and $q \leq P^\lambda R$. Then

$$\|\alpha(qh)^k\| \leq |(qh)^k \alpha - a(qh)^{k-1}| < (HP^\lambda R)^{k-1} P^{\lambda-k} \ll P^{-\sigma(k)}.$$

Then in this case

$$\min_{1 \leq n \leq P} \|\alpha n^k\| \leq \|\alpha(qh)^k\| < P^{\phi-\sigma(k)}.$$

In the alternative case, for each triple h, a, q with $1 \leq h \leq H$, $a \in \mathbb{Z}$, $q \in \mathbb{N}$, $(a, q) = 1$ and $|qh\alpha - a| \leq P^{\lambda-k}$, we have $q > P^\lambda R$. Then by Corollary 2 to Theorem 4.2,

$$\max_{1 \leq h \leq H} |f(h\alpha; P, R)| \ll P^{1-\sigma(k)+\varepsilon}.$$

Consequently,

$$T(\alpha) \ll HP^{1-\sigma(k)+\varepsilon} = o(P),$$

and the desired conclusion follows once again.

Appendix. Numerical values for parameters.

In this appendix we display in tabular form the numerical values of the parameters arising in the method discussed in section 5. The displayed figures were calculated to 12 significant figures on a computer, and then the values of the permissible exponents Δ_n and $\sigma(k)^{-1}$ were rounded up in the last displayed figure.

k	s	w	t	Δ_s	Δ_w	Δ_t	$\sigma(k)^{-1}$	$\lambda(k)$	v	Δ_{v+1}
10	22	13	7	0.22849	1.46021	4.06022	97.962	0.55161	25	0.09524
11	25	15	8	0.22308	1.44899	4.32206	110.137	0.55358	29	0.08311
12	28	16	9	0.22111	1.70161	4.59367	122.577	0.55521	32	0.09005
13	31	18	9	0.22065	1.69429	5.45150	134.996	0.55702	35	0.09680
14	35	20	10	0.19066	1.69881	5.71309	147.591	0.55847	39	0.08897
15	38	21	11	0.19444	1.93796	5.97304	160.133	0.56014	43	0.08311
16	41	22	12	0.19883	2.18309	6.23839	172.870	0.56153	46	0.08993
17	44	24	12	0.20344	2.18066	7.09136	185.636	0.56296	50	0.08538
18	47	26	13	0.20843	2.18618	7.35059	198.499	0.56436	54	0.08188
19	51	27	14	0.19133	2.42272	7.60987	211.371	0.56567	57	0.08837
20	54	29	14	0.19729	2.42874	8.46768	224.370	0.56682	61	0.08546

References.

- [1] G.Harman. Trigonometric sums over primes I. *Mathematika* 28 (1981), 249-254.
- [2] D.R.Heath-Brown. On the fractional part of αn^k . *Mathematika* 35 (1988), 28-37.
- [3] K.Thanigasalam. Some new estimates for $G(k)$ in Waring's problem. *Acta Arith.* 42 (1982), 73-78.
- [4] R.C.Vaughan. *The Hardy-Littlewood method*. Cambridge University Press, Cambridge 1981.
- [5] R.C.Vaughan. A new iterative method in Waring's problem. *Acta Math.* 162 (1989), 1-71.
- [6] R.C.Vaughan & T.D.Wooley. On Waring's problem: some refinements. *Proc. Lond. Math. Soc.* (3) 63 (1991), 35-68.
- [7] R.C.Vaughan & T.D.Wooley. Further improvements in Waring's problem. (*Submitted for publication*).
- [8] I.M.Vinogradov. The method of trigonometrical sums in the theory of numbers. *Trav. Inst. Steklov* 23 (1947).
- [9] T.D.Wooley. Large improvements in Waring's problem. *Annals of Math.* 135 (1992), 131-164.

- [10] T.D.Wooley. On Vinogradov's mean value theorem. *Mathematika* 39 (1992), 379-399.
- [11] T.D.Wooley. The application of a new mean value theorem to the fractional parts of polynomials. *Acta Arith.* (to appear).

*T.D. Wooley,
Department of Mathematics,
University of Michigan,
Ann Arbor,
Michigan 48109-1003.
U.S.A.*