

COHEN–MACAULAYNESS AND COMPUTATION OF NEWTON GRADED TORIC RINGS

MATHIAS SCHULZE AND ULI WALTHER

ABSTRACT. Let $H \subseteq \mathbb{Z}^d$ be a positive semigroup generated by $\mathcal{A} \subseteq H$, and let $\mathbb{K}[H]$ be the associated semigroup ring over a field \mathbb{K} . We investigate heredity of the Cohen–Macaulay property from $\mathbb{K}[H]$ to both its \mathcal{A} -Newton graded ring and to its face rings. We show by example that neither one inherits in general the Cohen–Macaulay property. On the positive side we show that for every H there exist generating sets \mathcal{A} for which the Newton graduation preserves Cohen–Macaulayness. This gives an elementary proof for an important vanishing result on A -hypergeometric Euler–Koszul homology. As a tool for our investigations we develop an algorithm to compute algorithmically the Newton filtration on a toric ring.

CONTENTS

1.	Introduction	1
2.	A necessary criterion for Cohen–Macaulayness of toric rings	3
3.	An algorithm to compute the Newton filtration	7
4.	Cohen–Macaulayness of the Newton graded toric ring	15
	References	19

1. INTRODUCTION

The motivation for this article arises from the study of the solutions of A -hypergeometric systems. These systems of linear partial differential equations have been introduced by Gel’fand, Graev, Kapranov, and Zelevinskiĭ [GGZ87, GZK89] as a general framework including the classical hypergeometric functions.

Given an integer $d \times n$ -matrix $A = ((a_{i,j})) \in \mathbb{Z}^{d \times n}$, let D_A be the (complex) Weyl algebra in the variables $x = x_1, \dots, x_n$ with corresponding differential operators $\partial = \partial_1, \dots, \partial_n$ where $\partial_i = \frac{\partial}{\partial x_i}$ for $i = 1, \dots, n$. Writing $(\mathbf{u}_+)_j = \max(0, \mathbf{u}_j)$ and $\mathbf{u}_- = \mathbf{u}_+ - \mathbf{u}$ for $\mathbf{u} \in \mathbb{Z}^n$, the *toric relations* are defined by

$$\square_{\mathbf{u}} = \partial^{\mathbf{u}_+} - \partial^{\mathbf{u}_-}$$

for $\mathbf{u} \in \ker A \cap \mathbb{Z}^n$. For a complex parameter vector $\beta \in \mathbb{C}^d$, the induced A -hypergeometric system is the system partial differential equations for $f(x_1, \dots, x_n)$

1991 *Mathematics Subject Classification.* 14M25, 16W70.

Key words and phrases. toric ring, Newton filtration, Cohen–Macaulay.

MS was supported by the College of Arts & Sciences at Oklahoma State University through a FY 2008 Dean’s Incentive Grant.

UW was supported by the NSF under grant DMS 0555319, and by the NSA under grant H98230-06-1-0012.

given by all *Euler equations* $E_i \bullet (f) = \beta_i \cdot f$ where

$$E_i = \sum_{j=1}^n a_{i,j} x_j \partial_i$$

for $i = 1, \dots, d$, and all *toric equations* $\square_{\mathbf{u}} \bullet (f) = 0$ for $\mathbf{u} \in \ker A \cap \mathbb{Z}^n$. More intrinsically, one considers the A -hypergeometric D_A -module

$$M_A(\beta) = D_A / \langle E - \beta, I_A \rangle$$

where we abbreviate $E = E_1, \dots, E_d$ and

$$I_A = \langle \square_{\mathbf{u}} \mid \mathbf{u} \in \ker A \cap \mathbb{Z}^n \rangle \subseteq R_A = \mathbb{C}[\partial_1, \dots, \partial_n].$$

is the *toric ideal*. The quotient of R_A by the toric ideal defines the *toric ring*

$$S_A = R_A / I_A \cong \mathbb{C}[t^{\mathbf{a}_1}, \dots, t^{\mathbf{a}_n}] \subseteq \mathbb{C}[t_1^{\pm 1}, \dots, t_d^{\pm 1}].$$

A fundamental tool for studying $M_\beta(A)$ is to consider it as the 0-th homology of the *Euler–Koszul complex* of $E = E_1, \dots, E_d$ on $D \otimes_{R_A} S_A$, [MMW05]. Considered in a commutative polynomial ring $\mathbb{C}[x, \partial]$, the elements E form a subset of a system of parameters on $S_A[x]$. There are two main approaches to understanding the relationship of Euler–Koszul homology (and hence $M_A(\beta)$) with the commutative Koszul complex: Adolphson [Ado94] applied grading techniques with respect to the *Newton filtration* on S_A using ideas of Kouchnirenko [Kou76]; on the other hand, [MMW05] introduces homological methods on the category of *toric modules*, a natural \mathbb{Z}^d -graded category that contains all face rings of S_A and their \mathbb{Z}^d -shifts. Two natural questions arise:

- (Q1) Is the Newton graded ring of a Cohen–Macaulay toric ring Cohen–Macaulay?
- (Q2) Are the face rings of a Cohen–Macaulay toric ring Cohen–Macaulay?

Closer inspection shows that (Q1) does not quite make sense as stated: speaking of a Newton filtration requires a distinguished (finite) set of generators on the toric ring in order to form the Newton polyhedron. The (historic) default choice for S_A is the column set of A . In that case, Okuyama claims a positive answer to (Q1) in [Oku06b, Prop. 3.1] and [Oku06a, Prop. 3.4] and derives certain vanishing properties for Euler–Koszul homology. We shall show that these claims are incorrect. However, we also show that for suitable choices of generating sets the answer to (Q1) is positive, we prove the vanishing of Euler–Koszul homology on Cohen–Macaulay rings, and we describe an algorithm to compute the Newton filtration induced by arbitrary (finite) generating sets.

Regarding (Q2) we will show that in general the answer is negative as well, and elaborate on a criterion discussed in [TH86] regarding Cohen–Macaulayness of toric rings.

The following notation will be used throughout this note.

Notation 1.1. We let $\mathbb{N} = \{0, 1, 2, \dots\}$ be the natural numbers and denote by \mathbb{Q}_+ the non-negative rational numbers.

Let H be an *affine semigroup*. By this we mean a finitely generated, commutative, torsion-free, cancelative monoid. We assume that H is *positive* which means that 0 is the only invertible element. By Hochster’s transformation [Hoc72, §2], H may be viewed as a submonoid of \mathbb{N}^d for some $d \in \mathbb{N}$. The Grothendieck group $\mathbb{Z}H$ of H is isomorphic to \mathbb{Z}^r where $r = \text{rk } H$ and we can view $H \subseteq \mathbb{Z}H$. We write

$\mathbb{K}H$ for the \mathbb{K} -vector space $\mathbb{K} \otimes_{\mathbb{Z}} \mathbb{Z}H$ and identify $h \in H$ with $1 \otimes h \in \mathbb{K}H$ if \mathbb{K} has characteristic zero.

Let \mathbb{K} be an arbitrary field and $\mathbb{K}[H]$ the semigroup ring of H over \mathbb{K} . The inclusion $H \subseteq \mathbb{Z}H \cong \mathbb{Z}^r$ (or $H \subseteq \mathbb{N}^d$) gives rise to an inclusion $\mathbb{K}[H] \subseteq \mathbb{K}[\mathbb{Z}^r] \cong \mathbb{K}[t_1^{\pm 1}, \dots, t_r^{\pm 1}]$ (or $\mathbb{K}[H] \subseteq \mathbb{K}[\mathbb{N}^d] \cong \mathbb{K}[t_1, \dots, t_d]$). This makes $\text{Spec}(\mathbb{K}[H])$ a toric variety and $\mathbb{K}[H]$ a *toric ring* endowed with a canonical $\mathbb{Z}H$ -grading given by $\deg(t^h) = h$ for $h \in H$. Generally, for a subset $S \subseteq \mathbb{Z}^d$, we let $\mathbb{K}[S] := \bigoplus_{s \in S} \mathbb{K} \cdot t^s$. We denote by $\mathfrak{m} = \mathfrak{m}_H = \mathbb{K}[H \setminus \{0\}]$ the unique maximal $\mathbb{Z}H$ -graded ideal which defines the vertex of $\text{Spec}(\mathbb{K}[H])$.

Generally, \mathcal{A} will be a finite set of semigroup generators for H . Then \mathcal{A} defines a presentation

$$R_{\mathcal{A}} = \mathbb{K}[(y_a)_{a \in \mathcal{A}}] = \mathbb{K}[\mathbb{N}^{\mathcal{A}}] \twoheadrightarrow \mathbb{K}[H] = S_{\mathcal{A}}$$

with kernel $I_{\mathcal{A}}$ and we write $y_i = y_{a_i}$ for $i = 1, \dots, n$ if $\mathcal{A} = \{a_1, \dots, a_n\}$. In the situation of hypergeometric systems, the given matrix $A = (a_1, \dots, a_n)$ replaces the generating set \mathcal{A} and we write ∂_i rather than y_i .

Consider the rational projective r -space

$$\mathbb{P}_{\mathbb{Q}}^H = \mathbb{P}_{\mathbb{Q}}(\mathbb{Q}H \times \mathbb{Q})$$

containing $\mathbb{Q}H$ via the embedding $qh \mapsto (qh : 1)$. Since H is positive, there is a linear functional $h \in \text{Hom}_{\mathbb{Q}}(\mathbb{Q}H, \mathbb{Q})$ such that $h^{-1}(0) \cap \mathbb{Q}_+H = \{0\}$ while $h(H) \geq 0$. Pick a rational weight vector $L: \mathcal{A} \rightarrow \mathbb{Q}$ and choose $0 > \varepsilon \in \mathbb{Q}$ such that $L(a) < 0$ implies $\varepsilon > h(a)/L(a)$. This can be done since \mathcal{A} is finite, cf. [SW06]. The convex hull in $\mathbb{P}_{\mathbb{Q}}^H \setminus h^{-1}(-\varepsilon)$ of the origin $0 = (0 : 1)$ and the points $a^L := (a : L(a))$, $a \in \mathcal{A}$, is the (\mathcal{A}, L) -polyhedron. Its simplicial complex $\Phi_{\mathcal{A}}^L$ of faces not containing $(0 : 1)$ is the (\mathcal{A}, L) -umbrella, cf. [SW06, §2.3]. The dimension $\dim(\tau)$ of $\tau \in \Phi_{\mathcal{A}}^L$ is its topological dimension as a boundary component of $\Delta_{\mathcal{A}}^L$. We denote by $\Phi_{\mathcal{A}}^{L,k} \subseteq \Phi_{\mathcal{A}}^L$ the subcollection of faces of dimension k , hence a *facet* is of dimension $r - 1$. We reserve σ for elements of $\Phi_{\mathcal{A}}^{L,r-1}$ and τ for general elements in $\Phi_{\mathcal{A}}^L$.

The *interior* (\mathcal{A}, L) -umbrella $\dot{\Phi}_{\mathcal{A}}^L$ consists of the faces $\tau \in \Phi_{\mathcal{A}}^L$ for which there is no facet σ of $\Delta_{\mathcal{A}}^L$ with $\tau \subseteq \sigma \ni 0$. One may view $\dot{\Phi}_{\mathcal{A}}^L$ as the quotient complex of $\Phi_{\mathcal{A}}^L$ by the subcomplex of faces contained in its boundary as a piecewise linear $(r - 1)$ -manifold.

Consider the special weight vectors $\mathbf{0} = (0)_{a \in \mathcal{A}}$ and $\mathbf{1} = (1)_{a \in \mathcal{A}}$. Then $\Delta_{\mathcal{A}}^{\mathbf{1}}$ is the convex hull of \mathcal{A} and 0 , while $\Delta_{\mathcal{A}}^{\mathbf{0}}$ is the closure in $\mathbb{P}_{\mathbb{Q}}^H$ of the positive rational cone $C_H = \mathbb{Q}_+H$. One can identify $\dot{\Phi}_{\mathcal{A}}^{\mathbf{0}}$ with the simplicial complex of a cross section of the cone C_H . If \mathcal{A} is contained in a hyperplane in $\mathbb{Q}H$ then $\mathbb{K}[H]$ is the coordinate ring of a projective variety and we say that \mathcal{A} is *projective*.

Let $\tau \in \Phi_{\mathcal{A}}^L$. We call the cone $C_{\tau} = \mathbb{Q}_+\tau$ over τ a *face cone* in general and *facet cone* if τ is a facet. We abbreviate $H_{\tau} = H \cap C_{\tau}$ and $\mathcal{A}_{\tau} = \mathcal{A} \cap C_{\tau}$, and we denote by \mathfrak{m}_{τ} the maximal $\mathbb{Z}H$ -graded ideal $\mathbb{K}[H_{\tau} \setminus \{0\}]$ in $\mathbb{K}[H_{\tau}]$. For a facet $\sigma \in \Phi_{\mathcal{A}}^{L,r-1}$, we denote by ℓ_{σ} the unique linear form on $\mathbb{Q}H$ such that $\max \ell_{\sigma}(\Delta_{\mathcal{A}}^{\mathbf{1}}) = 1$ and $\ell_{\sigma}^{-1}(1) \cap \Delta_{\mathcal{A}}^{\mathbf{1}} = \sigma$.

2. A NECESSARY CRITERION FOR COHEN–MACAULAYNESS OF TORIC RINGS

In this section, we discuss Cohen–Macaulayness of affine semigroup rings following Hoa and Trung [TH86, HT88]. We give a simplified proof for one of their results and use it to find a negative answer to Question (Q2).

The following result is classical and a special case of Proposition 3.6.

Lemma 2.1. *The set of $\mathbb{Z}H$ -graded primes in $\mathbb{K}[H]$ is in bijection with $\Phi_{\mathcal{A}}^{\mathbf{0}}$. Each $\tau \in \Phi_{\mathcal{A}}^{\mathbf{0}}$ corresponds to a prime ideal $\mathfrak{p}_{\tau} = \mathbb{K}[H \setminus C_{\tau}]$. \square*

Definition 2.2. Let \bar{H} be the set of $h \in \mathbb{Z}H$ for which there exists $\mathcal{B} \subseteq H$, $0 \notin \mathcal{B}$, such that $\mathcal{B}_{\tau} \neq \emptyset$ for all $\tau \in \Phi_{\mathcal{A}}^{\mathbf{0}, r-2}$ while $h + \mathcal{B} \subseteq H$.

For example, if $H = 2\mathbb{N} + 3\mathbb{N} \subseteq \mathbb{N}$ then $\bar{H} = \mathbb{Z}$, where $k \in \bar{H}$ is witnessed by $\mathcal{B}_k = \{3 - k\}$ and the equation $k + (3 - k) = 3 \in H$. Note that these relations encode cosets $t^k + \mathbb{K}[H]$ in $H_{\langle \mathcal{B} \rangle}^1(\mathbb{K}[H]) = \mathbb{K}[\mathbb{Z}H]/\mathbb{K}[H]$ which are non-zero for $k \leq 1$.

Note that $\bar{H} \supseteq H$ and if $r \geq 2$ then $\bar{H} \subseteq C_H$. Indeed, for all $h \in \bar{H}$ and for any $\tau \in \Phi_{\mathcal{A}}^{\mathbf{0}, r-2}$ there is a $b \in \mathcal{B} \cap C_{\tau}$ with $h + b \in H$. So any linear functional that is non-negative on H and vanishes on C_{τ} must be non-negative on h .

Part (3) of the following result (cf. [TH86, Cor. 2.2], [HT88, Lem. 4.3]) states a necessary condition for $\mathbb{K}[H]$ to be Cohen–Macaulay; our contribution is a simplified proof. When combined with the topological description of the Cohen–Macaulayness of $\mathbb{K}[\bar{H}]$ from [TH86, HT88], this characterizes Cohen–Macaulayness of $\mathbb{K}[H]$.

Proposition 2.3. *Let $\mathcal{B} \subseteq H$ with $0 \notin \mathcal{B} \neq \emptyset$ and put $\mathfrak{b} = \langle t^b \mid b \in \mathcal{B} \rangle \subseteq \mathbb{K}[H]$.*

- (1) $H_{\mathfrak{m}}^1(\mathbb{K}[H]) \subseteq \mathbb{K}[\bar{H}]/\mathbb{K}[H]$ and $H_{\mathfrak{m}}^1(\mathbb{K}[H]) \subseteq H_{\mathfrak{b}}^1(\mathbb{K}[H])$.
- (2) For $h \in \mathbb{Z}H \setminus H$ with $h + \mathcal{B} \subseteq H$, h induces a non-zero element in $H_{\mathfrak{b}}^1(\mathbb{K}[H])$.
In particular, $\mathbb{K}[\bar{H}]/\mathbb{K}[H] \subseteq \sum_{\mathcal{B}} H_{\mathfrak{b}}^1(\mathbb{K}[H])$ with \mathcal{B} as in Definition 2.2.
- (3) If $\mathbb{K}[H]$ is Cohen–Macaulay of dimension $r \geq 2$ then $H = \bar{H}$.
- (4) For $\mathbb{K}[H]$ of dimension $r = 2$, $\bar{H}/H = H_{\mathfrak{m}}^1(\mathbb{K}[H])$.

Proof. Any $\mathbb{Z}H$ -graded element of $H_{\mathfrak{m}}^1(\mathbb{K}[H])$ is the coset modulo $\mathbb{K}[H]$ of an element $(t^{h_1 - c_1 a_1}, \dots, t^{h_k - c_k a_k})$ where $a_1, \dots, a_k \in \mathcal{A}$ generate H and where $c_i \in \mathbb{N}$ and $h_i \in H$ with $h_1 - c_1 a_1 = h_j - c_j a_j$ for all j . So $h_1 - c_1 a_1 \in \bar{H}$ with $\mathcal{B} = \{a_1, \dots, a_k\}$ which proves the first inclusion.

As $\mathbb{K}[H]$ is a domain, $H_{\mathfrak{b}}^1(\mathbb{K}[H]) = (\bigcap_{b \in \mathcal{B}} \mathbb{K}[H + \mathbb{Z}b]) / \mathbb{K}[H]$ while $H_{\mathfrak{m}}^1(\mathbb{K}[H]) = (\bigcap_{a \in \mathcal{A}} \mathbb{K}[H + \mathbb{Z}a]) / \mathbb{K}[H]$. Pick $b \in \mathcal{B}$. Then $0 \neq b = \sum_{a \in \mathcal{A}} k_a a$ with $k_a \in \mathbb{N}$ implies $H + \mathbb{Z}b \supseteq H + \mathbb{Z}a$ for any a with $k_a > 0$. Thus, $\bigcap_{b \in \mathcal{B}} \mathbb{K}[H + \mathbb{Z}b]$ contains $\bigcap_{a \in \mathcal{A}} \mathbb{K}[H + \mathbb{Z}a]$ and the first claim follows.

By definition $(\mathbb{K}[H] + \mathbb{K}[H] \cdot t^h) / \mathbb{K}[H]$ is \mathfrak{b} -torsion. Applying $R\Gamma_{\mathfrak{b}}$ to the short exact sequence

$$0 \rightarrow \mathbb{K}[H] \rightarrow \mathbb{K}[H] + \mathbb{K}[H] \cdot t^h \rightarrow (\mathbb{K}[H] + \mathbb{K}[H] \cdot t^h) / \mathbb{K}[H] \rightarrow 0$$

yields

$$\underbrace{\Gamma_{\mathfrak{b}}(\mathbb{K}[H] + \mathbb{K}[H] \cdot t^h)}_{=0} \rightarrow \underbrace{(\mathbb{K}[H] + \mathbb{K}[H] \cdot t^h) / \mathbb{K}[H]}_{\ni t^h + \mathbb{K}[H]} \rightarrow H_{\mathfrak{b}}^1(\mathbb{K}[H]),$$

which proves the second claim.

By definition, $\mathbb{K}[\bar{H}]/\mathbb{K}[H]$ is \mathfrak{b} -torsion. Applying $R\Gamma_{\mathfrak{b}}$ to the short exact sequence

$$0 \rightarrow \mathbb{K}[H] \rightarrow \mathbb{K}[\bar{H}] \rightarrow \mathbb{K}[\bar{H}]/\mathbb{K}[H] \rightarrow 0$$

yields

$$0 = \Gamma_{\mathfrak{b}}(\mathbb{K}[\bar{H}]) \rightarrow \mathbb{K}[\bar{H}]/\mathbb{K}[H] \rightarrow H_{\mathfrak{b}}^1(\mathbb{K}[H]),$$

which proves the second claim.

For the third claim assume that $\mathbb{K}[H]$ is Cohen–Macaulay with $\dim(\mathbb{K}[H]) \geq 2$. Let $\mathcal{B} \subseteq H$ with $0 \notin \mathcal{B} \neq \emptyset$. Then $\mathfrak{b} \neq 0$ and hence $\dim(\mathbb{K}[H]/\mathfrak{b}) < r = \dim(\mathbb{K}[H])$ as $\mathbb{K}[H]$ is a domain. Since \mathfrak{b} is monomial, it is \mathbb{Z}^d -graded. So by Lemma 2.1 any associated prime of \mathfrak{b} is of the form \mathfrak{p}_τ with $\tau \in \Phi_{\mathcal{A}}^{\mathbf{0}}$. Hence there is an inclusion

$$\mathbb{K}[H_\tau] = \mathbb{K}[H]/\mathfrak{p}_\tau \hookrightarrow \mathbb{K}[H]/\mathfrak{b}.$$

Now let $h \in \bar{H}$ and consider a \mathcal{B} corresponding to h in Definition 2.2. Then \mathfrak{b} contains a non-zerodivisor on $\mathbb{K}[H_{\tau'}]$ for every $\tau' \in \Phi_{\mathcal{A}}^{\mathbf{0}, r-2}$. On the other hand, \mathfrak{b} kills $\mathbb{K}[H_\tau] \subseteq \mathbb{K}[H]/\mathfrak{b}$, so $\dim(\tau) \leq r - 3$. This being so for all associated primes of \mathfrak{b} , $\text{height } \mathfrak{b} \geq 2$. As $\mathbb{K}[H]$ is Cohen–Macaulay, $2 \leq \text{height } \mathfrak{b} = \text{depth } \mathfrak{b}$ and hence $H_{\mathfrak{b}}^1(\mathbb{K}[H]) = 0$. It follows from (2) that we therefore must have $h \in H$ and so $\bar{H} = H$ as claimed.

Finally assume that $r = 2$. Then any \mathcal{B} as in Definition 2.2 meets every C_τ , $\tau \in \Phi_{\mathcal{A}}^{\mathbf{0}, 0}$, and in particular the extremal rays of C_H . This means that \mathfrak{b} is \mathfrak{m} -primary and hence $\mathbb{K}[\bar{H}]/\mathbb{K}[H] \subseteq H_{\mathfrak{m}}^1(\mathbb{K}[H])$ by (2). Then the last claim follows from the first inclusion in (1). \square

Remark 2.4. In fact, \bar{H} is a subsemigroup of $\mathbb{Z}H$ and if $r = 2$ then $\mathbb{K}[\bar{H}]$ is the ideal transform of $\mathbb{K}[H]$ relative to \mathfrak{m} , cf. [BS98].


Investigating cases where the criterion of Proposition 2.3 applies to facets, but not all, of H , we found the following example showing that the answer to Question (Q2) is negative: the facet rings of a Cohen–Macaulay toric ring are not Cohen–Macaulay in general, even in the projective case.

Example 2.5. Consider the affine semigroup H generated by

$$(2.1) \quad \mathcal{A} = \{a_1, \dots, a_6\} = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 4 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 4 \\ 1 \end{pmatrix} \right\}.$$

Note that $\ell = (1, 0, 0)$ equals 1 on \mathcal{A} , so H is projective. The linear form

FIGURE 1. The $(\mathcal{A}, \mathbf{1})$ -polyhedron for Example 2.5



$\ell_1 = (0, 0, 1)$ defines the face $\tau_1 \in \Phi_{\mathcal{A}}^{\mathbf{0}, 1}$ spanned by $\mathcal{A}' = \{a_1, a_2, a_3, a_4\}$; the corresponding submonoid H_{τ_1} of H is $\mathbb{N}\mathcal{A}'$. With $\mathcal{B} = \{a_1, a_4\}$,

$$2a_3 - a_4 = 2a_2 - a_1 = (1, 2, 0) \in \bar{H}_{\tau_1} \setminus H_{\tau_1}$$

and so $\mathbb{K}[H_{\tau_1}]$ is not Cohen–Macaulay, as is well-known.

We now test whether $(1, 2, 0) \in \bar{H}$. Inspection shows that $\Phi_{\mathcal{A}}^{\mathbf{0}, 1} = \{\tau_1, \dots, \tau_4\}$ where

$$\tau_1 \ni a_1, a_2, a_3, a_4, \quad \tau_2 \ni a_1, a_5, \quad \tau_3 \ni a_4, a_6, \quad \tau_4 \ni a_5, a_6$$

are defined by the functionals

$$\ell_1 = (0, 0, 1), \ell_2 = (0, 1, -1), \ell_3 = (1, -1/4, 0), \ell_4 = (1, 0, -1).$$

So any set \mathcal{B} to be used in Definition 2.2 must contain an element $b \in \mathbb{N}a_5 + \mathbb{N}a_6$. Consider a relation $(1, 2, 0) + kb \in H$ with $k \in \mathbb{N}$. Note that $\ell_1(a_5) = \ell_1(a_6)$ while $\ell_1(\tau_1) = 0$ and so a_5 and a_6 must appear with opposite coefficients. Moving the a_5 - and a_6 -terms to one side, one obtains a relation $(1, 2, 0) + k(a_6 - a_5) \in H_{\tau_1}$ where now $k \in \mathbb{Z}$. As $H_{\tau_1} \cap \ell_1^{-1}(1) = \mathcal{A}'$ we find that $(1, 2, 0) + k(a_6 - a_5) = (1, 2, 0) + k(0, 3, 0) \in \mathcal{A}'$ which is clearly impossible. It follows that $(1, 2, 0) \notin \bar{H}$ and so $H_{\mathfrak{m}}^1(\mathbb{K}[H])$ is zero in degree $(1, 2, 0)$.

We now show that $\mathbb{K}[H]$ is actually a Cohen–Macaulay ring. To this end consider the ideal J of $\mathbb{K}[H]$ given by

$$(2.2) \quad J = \langle I_{\mathcal{A}'}, y_4y_5 - y_2y_6, y_3y_5 - y_1y_6 \rangle.$$

Obviously, J is contained in the toric ideal $I_{\mathcal{A}}$, and $J + \langle y_1, y_4 + y_5, y_6 \rangle$ is \mathfrak{m} -primary since its residue ring is spanned by the monomials

$$1, \quad y_2, \quad y_3, \quad y_5, \quad y_2^2, \quad y_2y_5, \quad y_3^2.$$

Hence $\dim(\mathbb{K}[H]/J) = 3$ and so $I_{\mathcal{A}}$ is a minimal prime of J . Below we will show that $R_{\mathcal{A}}/J$ is Cohen–Macaulay. This implies that every associated prime of J is of dimension 3. Thus, $J = I_{\mathcal{A}}$ provided that the two ideals have the same degree. The simplicial volume of $\mathcal{A} \cup \{0\}$ is 7 and equals the degree of $I_{\mathcal{A}}$. On the other hand, $\deg(R_{\mathcal{A}}/J) = \deg(R_{\mathcal{A}}/(J + \langle y_1, y_4 + y_5, y_6 \rangle))$ is also 7, and it follows that $J = I_{\mathcal{A}}$.

We proceed to showing that J is a Cohen–Macaulay ideal. The following is certainly Gröbner folklore but we don't know a reference.

Lemma 2.6. *Suppose J is an ideal in a polynomial ring $R = \mathbb{K}[y_1, \dots, y_n]$ and let \leq be a term order on R . Write $\text{in}_{\leq}(J)$ for the initial ideal of J under \leq . If y_n is not a zerodivisor on $R/\text{in}_{\leq}(J)$ then y_n is a non-zerodivisor on R/J .*

Proof. Let G be a reduced \leq -Gröbner basis for J and suppose $y_n f \in J$ for some $f \in R$.

If $f - f' \in J$ for some second element $f' \in R$ then $y_n f \in J$ if and only if $y_n f' \in J$ while of course $f \in J$ if and only if $f' \in J$. In particular, we may assume that f is equal to its \leq -normal form relative to G .

As $y_n f \in J$ we have $\text{in}_{\leq}(y_n f) \in \text{in}_{\leq}(J)$. The hypothesis implies that $\text{in}_{\leq}(f) \in \text{in}_{\leq}(J)$. Hence either $f = 0$, or f can be \leq -reduced relative to G . As f is in normal form, $f = 0$. \square

The lemma implies that y_6 is a non-zerodivisor on $R_{\mathcal{A}}/J$. Indeed, one may verify by hand that the four generators

$$y_2y_3 - y_1y_4, \quad y_2^3 - y_1^2y_3, \quad y_3^3 - y_2y_4^2, \quad y_3^2y_1 - y_2^2y_4$$

of $I_{\mathcal{A}'}$ together with the two further generators of J in (2.2) form a Gröbner basis for J under the graded reverse-lexicographic order. Since the initial terms of J do not involve y_6 the desired conclusion follows.

It hence suffices to show that the two-dimensional quotient

$$S' = R_{\mathcal{A}}/(J + \langle y_6 \rangle) \cong \mathbb{K}[H_{\tau_1}][y_5]/\langle y_3y_5, y_4y_5 \rangle$$

has vanishing 0-th and 1-st local cohomology with respect to the maximal ideal $\mathfrak{m}' = \langle y_1, \dots, y_5 \rangle$ of S' . The decomposition $\langle y_3 y_5, y_4 y_5 \rangle = \langle y_5 \rangle \cap \langle y_3, y_4 \rangle$ in $\mathbb{K}[H_{\sigma_1}][y_5]$ gives a short exact sequence

$$0 \rightarrow \frac{\mathbb{K}[H_{\tau_1}][y_5]}{\langle y_3 y_5, y_4 y_5 \rangle} \rightarrow \frac{\mathbb{K}[H_{\tau_1}][y_5]}{\langle y_5 \rangle} \oplus \frac{\mathbb{K}[H_{\tau_1}][y_5]}{\langle y_3, y_4 \rangle} \rightarrow \frac{\mathbb{K}[H_{\tau_1}][y_5]}{\langle y_3, y_4, y_5 \rangle} \rightarrow 0.$$

Note that $\mathbb{K}[H_{\tau_1}][y_5]/\langle y_3, y_4 \rangle \cong \mathbb{K}[y_1, y_2, y_5]/\langle y_2^3 \rangle$ so that the two rightmost displayed rings are complete intersections. As $\mathbb{K}[H_{\tau_1}]$ is a domain, the long exact local cohomology sequence with support in \mathfrak{m}' reads

$$\begin{aligned} 0 \rightarrow H_{\mathfrak{m}'}^0(S') \rightarrow 0 \oplus 0 \rightarrow \\ 0 \rightarrow H_{\mathfrak{m}'}^1(S') \rightarrow H_{\mathfrak{m}'}^1\left(\frac{\mathbb{K}[H_{\tau_1}][y_5]}{\langle y_5 \rangle}\right) \oplus 0 \rightarrow H_{\mathfrak{m}'}^1\left(\frac{\mathbb{K}[H_{\tau_1}][y_5]}{\langle y_3, y_4, y_5 \rangle}\right) \rightarrow \dots \end{aligned}$$

Our claim then follows if we can show that

$$H_{\mathfrak{m}'}^1(\mathbb{K}[H_{\tau_1}][y_5]/\langle y_5 \rangle) \rightarrow H_{\mathfrak{m}'}^1(\mathbb{K}[H_{\tau_1}][y_5]/\langle y_3, y_4, y_5 \rangle)$$

is injective. However, $H_{\mathfrak{m}'}^1(\mathbb{K}[H_{\tau_1}][y_5]/\langle y_5 \rangle) \cong \mathbb{K} \cdot (y_2^2/y_1, y_3^2/y_4, 0)$, generated by the indicated 1-cocycle in the Čech complex to y_1, y_4, y_5 on $\mathbb{K}[H_{\tau_1}][y_5]/\langle y_5 \rangle$. Modulo y_3 and y_4 this becomes the class of y_2^2/y_1 in the Čech complex of y_1 on $\mathbb{K}[H_{\tau_1}]/\langle x_3, x_4 \rangle \cong \mathbb{K}[y_1, y_2]/\langle y_2^3 \rangle$, and that is clearly non zero.

Of course, all claims made are corroborated by computer results.

3. AN ALGORITHM TO COMPUTE THE NEWTON FILTRATION

A rational polyhedron $0 \in \Delta \subseteq C_H$ defines a Newton filtration $k \cdot \Delta \cap H$, $k \in \mathbb{Q}$, on H and hence on $\mathbb{K}[H]$. This filtration is separated if Δ is bounded; it is exhaustive if $k \cdot \Delta \cap H$ generates H for some k . We shall impose both conditions and assume that $\mathcal{A} = \Delta \cap H$ is a finite set of generators of H . To recover the Newton polytope $\Delta = \Delta_{\mathcal{A}}^1$ from \mathcal{A} we shall use Gröbner methods.

The main goal of this section to develop an algorithm to compute the associated graded ring to such a Newton filtration on $\mathbb{K}[H]$. This algorithm is based on Gröbner bases and represents our computational tool to approach Question (Q1).

We start with a formal definition of the Newton filtration relative to \mathcal{A} . Recall that ℓ_{σ} is the linear functional associated to the facet $\sigma \in \Phi_{\mathcal{A}}^{1,r-1}$.

Definition 3.1. For $h \in H$ its (Newton) \mathcal{A} -degree is

$$\deg_{\mathcal{A}}(h) = \max\{\ell_{\sigma}(h) \mid \sigma \in \Phi_{\mathcal{A}}^{1,r-1}\} \in \mathbb{Q}.$$

Note that $\deg_{\mathcal{A}}(h)$ is the rational number k for which h is precisely on the boundary of $k \cdot \Delta_{\mathcal{A}}^1$, and that $\deg_{\mathcal{A}}$ is subadditive on H :

$$\deg_{\mathcal{A}}(h + h') \leq \deg_{\mathcal{A}}(h) + \deg_{\mathcal{A}}(h') \text{ for all } h, h' \in H.$$

The \mathcal{A} -degree defines the increasing Newton filtration $N_{\bullet} = N_{\bullet}^{\mathcal{A}}$ on H by

$$N_k H = \{h \in H \mid \deg_{\mathcal{A}}(h) \leq k\} \text{ for all } k \in \mathbb{Q}_+.$$

Definition 3.2. Let $H \subseteq \mathbb{Z}^d$ be any affine semigroup with increasing \mathbb{Q} -indexed discrete exhaustive filtration F_{\bullet} such that $F_i + F_j \subseteq F_{i+j}$. We let $\deg_F(h)$ denote the smallest i with $h \in F_i(h)$. The graded associated semigroup $\text{gr}^F(H)$ is the set

$$\{[h] \mid h \in H\} \sqcup \{-\infty\}$$

subject to the rules

$$[h] + [h'] = \begin{cases} -\infty & \text{if } h = -\infty \text{ or } h' = -\infty \\ -\infty & \text{if } \deg_F(h + h') < \deg_F(h) + \deg_F(h') \\ [h + h'] & \text{otherwise.} \end{cases}$$

In the dictionary between semigroup operations and semigroup ring operations, sums equal to $-\infty$ encode zerodivisors in the associated graded ring.

If the filtration in question is the Newton filtration to \mathcal{A} we denote the associated graded semigroup by $\text{gr}^{\mathcal{A}}(H)$ rather than $\text{gr}^{N^{\mathcal{A}}}(H)$. In contrast to filtrations induced by an additive weight, the Newton filtration may have an associated graded object whose generators are not immediately obvious.

Example 3.3. Let $H = \mathbb{N}\mathcal{A}$ with $\mathcal{A} = \{(1, 0), (2, 2), (0, 1)\} \subseteq \mathbb{Z}^2$, so $H = \mathbb{N}(1, 0) + \mathbb{N}(0, 1)$. Although $(1, 0) + (0, 1)$ equals $(1, 1) \in H$, the corresponding sum is $-\infty$ in the associated graded semigroup $\text{gr}^{\mathcal{A}}(H)$. To see this, note that $\deg_{\mathcal{A}}((1, 1)) = 1/2 < 2 = \deg_{\mathcal{A}}((1, 0)) + \deg_{\mathcal{A}}((0, 1))$. This also implies that $\text{gr}_{\geq 1}^{\mathcal{A}}(H)$ cannot contain a generating set for $\text{gr}^{\mathcal{A}}(H)$ and in particular the cosets of $(1, 0)$, $(2, 2)$ and $(0, 1)$ are not generators of the graded object.

The following fundamental fact is stated, but not proved, in [Kou76, (6)]. It puts the above example in perspective and determines the additive structure of $\text{gr}^{\mathcal{A}}(H)$. In consequence, it describes the ring structure of $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$, which as $\mathbb{Z}H$ -graded \mathbb{K} -vector space can be identified with $\mathbb{K}[H]$. In particular, one finds that $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ contains a copy of $\mathbb{K}[H_{\tau}]$ for all $\tau \in \Phi_{\mathcal{A}}^1$, cf. Section 4). For convenience of the reader we provide a proof.

Lemma 3.4. *The equality $\deg_{\mathcal{A}}(h + h') = \deg_{\mathcal{A}}(h) + \deg_{\mathcal{A}}(h')$ holds if and only if h, h' share a $\mathbf{1}$ -facet cone.*

Proof. If $h \in H_{\sigma}$ with $\sigma \in \Phi_{\mathcal{A}}^{1, r-1}$ then $\deg_{\mathcal{A}}(h) = \ell_{\sigma}(h)$ and hence $\ell_{\sigma'}(h) \leq \ell_{\sigma}(h)$ for all $\sigma' \in \Phi_{\mathcal{A}}^{1, r-1}$. If in addition $h \notin C_{\sigma'}$ then $h/\ell_{\sigma}(h) \in \Delta_{\mathcal{A}}^1 \setminus \sigma'$ and hence $\ell_{\sigma'}(h) < \ell_{\sigma}(h)$. Now let $h \in H_{\sigma}$, $h' \in H_{\sigma'}$, and $h + h' \in C_{\sigma''}$ for some $\sigma, \sigma', \sigma'' \in \Phi_{\mathcal{A}}^{1, r-1}$. Then

$$\begin{aligned} \deg_{\mathcal{A}}(h + h') &= \ell_{\sigma''}(h + h') = \ell_{\sigma''}(h) + \ell_{\sigma''}(h') \\ &\leq \ell_{\sigma}(h) + \ell_{\sigma'}(h') = \deg_{\mathcal{A}}(h) + \deg_{\mathcal{A}}(h'). \end{aligned}$$

If $h, h' \in C_{\sigma''}$ then one can choose $\sigma = \sigma' = \sigma''$ so that the \mathcal{A} -degree of h, h' and $h + h'$ is evaluated by the same linear functional $\ell_{\sigma''}$. So equality in the above display follows in this case. If conversely either $h \notin C_{\sigma''}$ or $h' \notin C_{\sigma''}$ then the inequality is strict by the remarks before the display. \square

Let $\mathcal{A}' \supseteq \mathcal{A}$ be a second set of generators for H . It defines a free presentation of the monoid H ,

$$(3.1) \quad \varphi_{\mathcal{A}'}: F_{\mathcal{A}'} = \bigoplus_{a' \in \mathcal{A}'} \mathbb{N} \cdot e_{a'} \rightarrow H, \quad e_{a'} \mapsto a'.$$

where $\{e_{a'} \mid a' \in \mathcal{A}'\}$ is the distinguished monoid basis of $F_{\mathcal{A}'} = \mathbb{N}^{\mathcal{A}'}$. The equalizer of $\varphi_{\mathcal{A}'}$,

$$E_{\mathcal{A}'} = \{(p, q) \in F_{\mathcal{A}'} \times F_{\mathcal{A}'} \mid \varphi_{\mathcal{A}'}(p) = \varphi_{\mathcal{A}'}(q)\},$$

defines an equivalence relation $\sim_{\varphi_{\mathcal{A}'}}$ on $F_{\mathcal{A}'}$ by $[p \sim_{\varphi_{\mathcal{A}'}} q] \Leftrightarrow [(p, q) \in E_{\mathcal{A}'}]$. Of course, the set of cosets with induced additive structure is precisely H .

Let $L = (L_{a'})_{a' \in \mathcal{A}'}$ be the vector of Newton degrees relative to \mathcal{A}' :

$$(3.2) \quad L_{a'} = \deg_{\mathcal{A}'}(a').$$

The corresponding linear form $\deg_L = \sum_{a' \in \mathcal{A}'} L_{a'} e_{a'}^*$ on $\mathbb{Q}F_{\mathcal{A}'}$ defines an additive degree and an increasing filtration $L_{\bullet} = L_{\bullet}^{\mathcal{A}'}$ on $F_{\mathcal{A}'}$ by

$$(3.3) \quad L_k F_{\mathcal{A}'} = \{p \in F_{\mathcal{A}'} \mid \deg_L(p) \leq k\} \text{ for all } k \in \mathbb{Q}_+.$$

The equalizer $E_{\mathcal{A}'}$ is equipped with an induced L -filtration via the inclusion $E_{\mathcal{A}'} \subseteq F_{\mathcal{A}'} \times F_{\mathcal{A}'}$. This endows H with a second rational increasing filtration besides the \mathcal{A} -Newton filtration: the filtration given by the degree function

$$(3.4) \quad \deg_L(\varphi_{\mathcal{A}'}(p)) = \min\{\deg_L(q) \mid p \sim_{\varphi_{\mathcal{A}'}} q\}.$$

In order to keep the notation straight we denote this second incarnation of H with the corresponding filtration by $F_{\mathcal{A}'}/\sim_{\varphi_{\mathcal{A}'}}$. Both the \mathcal{A} -Newton filtration and the filtration $L_{\bullet}^{\mathcal{A}'}$ are equivalent to one with index set \mathbb{N} ; in particular, their index sets are well-ordered.

By (3.2), the additivity of \deg_L on $F_{\mathcal{A}'}$ and the subadditivity of $\deg_{\mathcal{A}'}$ on H imply $\deg_L(p) \geq \deg_{\mathcal{A}'}(\varphi_{\mathcal{A}'}(p))$ for all $p \in F_{\mathcal{A}'}$. Thus, $\varphi_{\mathcal{A}'}(L_k F_{\mathcal{A}'}) \subseteq N_k H$ and

$$(3.5) \quad L_k(F_{\mathcal{A}'}/\sim_{\varphi_{\mathcal{A}'}}) \subseteq N_k H$$

for all $k \in \mathbb{Q}$ which yields a morphism of filtered semigroups

$$\bar{\varphi}_{\mathcal{A}', \mathcal{A}'}: (F_{\mathcal{A}'}/\sim_{\varphi_{\mathcal{A}'}} , L_{\bullet}) \rightarrow (H, N_{\bullet}^{\mathcal{A}'}).$$

To (3.1) corresponds the \mathbb{K} -algebra morphism

$$(3.6) \quad \varphi_{\mathcal{A}'}^{\mathbb{K}}: \mathbb{K}[F_{\mathcal{A}'}] \rightarrow \mathbb{K}[H], \quad y^p \mapsto t^{\varphi_{\mathcal{A}'}(p)},$$

whose kernel is the toric ideal

$$(3.7) \quad I_{\mathcal{A}'} = \langle y^p - y^q \mid (p, q) \in E_{\mathcal{A}'} \rangle.$$

The L -filtration on $F_{\mathcal{A}'}$ induces an L -filtration on $\mathbb{K}[F_{\mathcal{A}'}]$ that descends to the filtration on $\mathbb{K}[F_{\mathcal{A}'}]/I_{\mathcal{A}'}$ given by the L -filtration on $F_{\mathcal{A}'}/\sim_{\varphi_{\mathcal{A}'}}$ from (3.4). By (3.5) we hence have a morphism of filtered \mathbb{K} -algebras

$$(3.8) \quad \bar{\varphi}_{\mathcal{A}', \mathcal{A}'}^{\mathbb{K}}: (\mathbb{K}[F_{\mathcal{A}'}]/I_{\mathcal{A}'}, L_{\bullet}) \rightarrow (\mathbb{K}[H], N_{\bullet}^{\mathcal{A}'}).$$

Note that

$$(3.9) \quad \text{gr}^L(\mathbb{K}[F_{\mathcal{A}'}]/I_{\mathcal{A}'}) = \mathbb{K}[F_{\mathcal{A}'}]/\text{gr}^L(I_{\mathcal{A}'}).$$

While (3.6) is a surjection with kernel $I_{\mathcal{A}'}$, (3.8) is not necessarily an epimorphism of filtered algebras. In particular, it may fail to induce a surjection of associated graded algebras; for the most elementary example see Example 3.3 with $\mathcal{A} = \mathcal{A}'$.

The following result describes precisely which sets \mathcal{A}' produce an isomorphism in (3.8). In this proposition, and hereafter, we make use of the fact that $\Delta_{\mathcal{A}'}^{\mathbf{1}, k} = \Delta_{\mathcal{A}'}^{L, k}$ (cf. Notation 1.1 and (3.2)) and hence $\Phi_{\mathcal{A}'}^{\mathbf{1}, k} = \Phi_{\mathcal{A}'}^{L, k}$ for all k .

Theorem 3.5. *For $\mathcal{A}' \supseteq \mathcal{A}$ the following conditions are equivalent:*

- (1) For each $\sigma \in \Phi_{\mathcal{A}'}^{\mathbf{1}, r-1}$, \mathcal{A}'_{σ} contains a generating set for the monoid H_{σ} .
- (2) For each $\tau \in \Phi_{\mathcal{A}'}^{\mathbf{1}}$, \mathcal{A}'_{τ} contains a generating set for the monoid H_{τ} .
- (3) The surjection (3.6) maps $L_k(F_{\mathcal{A}'})$ onto $N_k(H)$ for all $k \in \mathbb{Q}$.
- (4) The morphism (3.8) is an isomorphism of filtered semigroups.

Proof. It is clear that the last two conditions are equivalent. As faces are the intersection of the facets they are contained in, the first two conditions are equivalent as well.

Pick $h \in H$ with $\deg_{\mathcal{A}}(h) = l$, and let C_σ be any cone in $\Phi_{\mathcal{A}}^{1,r-1}$ that contains h . If condition (1) holds then there is a relation $h = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} a'$ and by Lemma 3.4 we have

$$\deg_{\mathcal{A}}(h) = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} \deg_{\mathcal{A}}(a') = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} \deg_L(e_{a'}) = \deg_L\left(\sum_{a' \in \mathcal{A}'_\sigma} k_{a'} e_{a'}\right).$$

Hence, $h \in \varphi_{\mathcal{A}'}(L_1 F_{\mathcal{A}'})$ and (3) follows.

Conversely, assume condition (1) fails and let, for suitable $\sigma \in \Phi_{\mathcal{A}}^{1,r-1}$, $h \in H_\sigma$ be an element not contained in $\mathbb{N}_{\mathcal{A}'_\sigma}$. Then any relation $h = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} a'$ involves at least one $a' \in \mathcal{A}'$ outside C_σ . As $h \in C_\sigma$ but $a' \notin C_\sigma$, Lemma 3.4 asserts that $\deg_{\mathcal{A}}(h)$ is strictly less than $\deg_L(\sum_{a' \in \mathcal{A}'_\sigma} k_{a'} e_{a'})$. This being so for all presentations for h , h cannot be in $\varphi_{\mathcal{A}'}(L_1 F_{\mathcal{A}'})$ and (3) cannot hold. \square

Algorithm 3.9 below computes the Newton filtration relative to \mathcal{A} by enlarging $\mathcal{A}' \supseteq \mathcal{A}$ until the conditions in Theorem 3.5 are fulfilled. The approach is to compare not the filtrations L_\bullet and $N_\bullet^{\mathcal{A}'}$ or the corresponding graded objects but the defining relations of the latter and to systematically add generators to \mathcal{A}' to reach equality of these relations. While Lemma 3.4 determines the relations of H , the relations $\text{gr}^L(E_{\mathcal{A}'})$ have been studied in general in [SW06] from where we shall extract Corollary 3.8 for our purposes. In Lemma 3.11 it will turn out that the above equality of relations enforces condition 3.5.(1) which will justify our procedure.

The following is a reformulation of [SW06, Thm. 2.15] which is valid for general L . Recall that Lemma 2.1 contains the special case $L = \mathbf{0}$.

Proposition 3.6. *For any \mathcal{A} and any $L = (L_a)_{a \in \mathcal{A}}$, any $\mathbb{Z}H$ -graded prime ideal in $\text{gr}^L(\mathbb{K}[F_{\mathcal{A}}]/I_{\mathcal{A}})$ is of the form*

$$I_\tau^L = I_{\{a \in \mathcal{A} \mid a^L \in \tau\}} + \langle y_a \mid a \in \mathcal{A}, a^L \notin \tau \rangle$$

for some $\tau \in \Phi_{\mathcal{A}}^L$. In particular, the radical of $\text{gr}^L(I_{\mathcal{A}'})$ is $\bigcap_{\sigma \in \Phi_{\mathcal{A}'}^{L,r-1}} I_\sigma^L$. \square

The following corollary adapts Proposition 3.6 to our special choice of L and yields the core of our procedure. Its first part relates relations in $\text{gr}^L(E_{\mathcal{A}'})$ to those of H defined by $\mathbf{1}$ -facets in Lemma 3.4. Its second part serves to determine all $\mathbf{1}$ -facets from a Gröbner basis in Steps 1 and 2 of our algorithm.

Definition 3.7. We call an L -leading term of $E_{\mathcal{A}'}$ an element $p \in F_{\mathcal{A}'}$ such that $\deg_L(p) > \deg_L(q)$ for some $(p, q) \in E_{\mathcal{A}'}$.

Corollary 3.8. *For $a'_1, a'_2 \in \mathcal{A}'$, the following conditions are equivalent:*

- (1) *For some $k \geq 1$, $k(e_{a'_1} + e_{a'_2})$ is an L -leading term of $E_{\mathcal{A}'}$.*
- (2) *The elements a'_1, a'_2 do not share a $\mathbf{1}$ -facet cone.*

If a_1, a_2 are actually in \mathcal{A} , the following conditions are equivalent:

- (3) *One has $\deg_{\mathcal{A}}(a_1) = 1 = \deg_{\mathcal{A}}(a_2)$, and a_1, a_2 do not share a $\mathbf{1}$ -facet cone.*
- (4) *For some $k \geq 1$, $k(e_{a_1} + e_{a_2})$ is a $\mathbf{1}$ -leading term of $E_{\mathcal{A}}$, but neither ke_{a_1} nor ke_{a_2} is a $\mathbf{1}$ -leading term of $E_{\mathcal{A}}$ for any $k \geq 1$.*

Proof. By (3.7) and (3.9), condition 3.8.(1) holds if and only if $t^{a'_1} t^{a'_2}$ is nilpotent in $\text{gr}^L(\mathbb{K}[F_{\mathcal{A}'})/I_{\mathcal{A}'})$. By Proposition 3.6 this happens exactly when $t^{a'_1} t^{a'_2}$ is contained

in $\bigcap_{\tau \in \Phi_{\mathcal{A}'}^L} I_{\tau}^L$. This is in turn equivalent to $(a'_1)^L, (a'_2)^L \in \tau$ for no $\tau \in \Phi_{\mathcal{A}'}^L = \Phi_{\mathcal{A}'}^1$. By definition, $\deg_{\mathcal{A}'}((a')^L) = 1$ for all $a' \in \mathcal{A}'$ and hence $(a')^L \in \tau$ is equivalent to $a' \in C_{\tau}$ for all $\tau \in \Phi_{\mathcal{A}'}^L = \Phi_{\mathcal{A}'}^1$. This proves equivalence of 3.8.(1) and 3.8.(2).

By the equivalence of conditions 3.8.(1) and 3.8.(2) above, the first condition in 3.8.(4) means that $a_1, a_2 \in \sigma$ for no $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$. On the other hand, the second condition means that neither t^{a_1} nor t^{a_2} is nilpotent in $\text{gr}^1(\mathbb{K}[H])$, which in turn, by Proposition 3.6, is equivalent to $\deg_{\mathcal{A}'}(a_1) = 1 = \deg_{\mathcal{A}'}(a_2)$. But under this latter condition, $a_1, a_2 \in \sigma$ for no $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$ is equivalent to $a_1, a_2 \in C_{\sigma}$ for no $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$. The equivalence of 3.8.(3) and 3.8.(4) follows. \square

We now state the algorithm and explain how to carry out its steps in practice, and then prove termination and correctness.

Algorithm 3.9 (Newton filtration on an affine semigroup).

INPUT: $\mathcal{A} \subseteq \mathbb{Z}^d$ such that $H = \mathbb{N}\mathcal{A}$ is positive.

OUTPUT: $H \supseteq \mathcal{A}' \supseteq \mathcal{A}$ and $L = (L_a)_{a \in \mathcal{A}'}$ such that (3.5) is an equality for all k .

1. Compute the set \mathcal{B} of $e_{a_1} + e_{a_2}$ where $a_1, a_2 \in \mathcal{A}$ such that $k(e_{a_1} + e_{a_2})$ is a **1**-leading term of $E_{\mathcal{A}}$ for some $k \geq 1$ but neither ke_{a_1} nor ke_{a_2} is a **1**-leading term of $E_{\mathcal{A}}$ for any $k \geq 1$.
2. Determine all $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$ by the rule: $a_1, a_2 \notin \sigma$ iff $e_{a_1} + e_{a_2} \in \mathcal{B}$.
3. For each $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$, compute the linear form ℓ_{σ} that equals 1 on σ .
4. For every pair $a_1, a_2 \in \mathcal{A}$ with $\deg_{\mathcal{A}'}(a_1) \deg_{\mathcal{A}'}(a_2) < 1$ update \mathcal{B} to include $e_{a_1} + e_{a_2}$ if a_1, a_2 do not share a **1**-facet cone.
5. Initialize $\mathcal{A}' = \mathcal{A}$ and $L = (L_{a'})_{a' \in \mathcal{A}'}$ by $L_{a'} = \deg_{\mathcal{A}'}(a')$.
6. Remove from \mathcal{B} all L -leading terms of $E_{\mathcal{A}'}$.
7. For each $e_{b_1} + e_{b_2} \in \mathcal{B}$ do the following:
 - 7.1 Update \mathcal{A}' to include $b_1 + b_2$.
 - 7.2 Update L with $L_{b_1+b_2} = \deg_{\mathcal{A}'}(b_1 + b_2)$.
 - 7.3 Update \mathcal{B} to include $e_{a'_1} + e_{a'_2}$ whenever $a'_1, a'_2 \in \mathcal{A}'$ do not share a **1**-facet cone.
8. If $\mathcal{B} \neq \emptyset$ continue with Step 6.
9. Return \mathcal{A}' and L .

Remark 3.10.

(1) By the correspondence (3.7) between $E_{\mathcal{A}'}$ and $I_{\mathcal{A}'}$, the set of all L -leading terms of $E_{\mathcal{A}'}$ is computable by Gröbner basis methods in rings of polynomials. Such computations may be carried out with standard programs such as [M2, GPS05].

(2) By the second part of Corollary 3.8, Step 1 adds expressions $e_{a_1} + e_{a_2}$ to the queue \mathcal{B} where $a_1, a_2 \in \mathcal{A}$ are in different facets of the **1**-umbrella of \mathcal{A} . The conditions in Step 1 translate to $y_{a_1}y_{a_2} \in \sqrt{\text{gr}^1(I_{\mathcal{A}})} \not\equiv y_{a_1}, y_{a_2}$ which may be tested via `decompose` in [M2] or with `radical` in [GPS05]. Alternatively, in large examples, one can use the estimate in Lemma 3.14 below to test this membership.

(3) The condition $a \in \sigma$ is, via Definition 3.1, equivalent to $\deg_{\mathcal{A}'}(a) = \ell_{\sigma}(a)$.

(4) Step 4 adds expressions $e_{a_1} + e_{a_2}$ to the queue \mathcal{B} where $a_1, a_2 \in \mathcal{A}$ are in different **1**-facet cones and at least one of them is in the interior of $\Delta_{\mathcal{A}'}^1$. Starting with the first passage of Step 6, the queue \mathcal{B} indicates pairs $\{a'_1, a'_2\}$ for which $[a'_1] + [a'_2] = -\infty$ in $\text{gr}^{\mathcal{A}'}(H)$ but not in $\text{gr}^L(F_{\mathcal{A}'}/\sim_{\varphi_{\mathcal{A}'}})$.

(5) By (3.9), the L -leading terms of an L -Gröbner basis of $I_{\mathcal{A}'}$ give a presentation of $\text{gr}^{\mathcal{A}'}(\mathbb{K}[H]) = \text{gr}^L(\mathbb{K}[F_{\mathcal{A}'}/I_{\mathcal{A}'}])$, cf. (3.13) in Example 3.15.

Our first task is to assure convergence of the algorithm. To this end fix a facet $\sigma \in \Phi_{\mathcal{A}}^{1,r-1}$. Any increasing sequence of subsets $\mathcal{A}_\sigma = \mathcal{A}_\sigma^0 \subseteq \mathcal{A}_\sigma^1 \subseteq \mathcal{A}_\sigma^2 \subseteq \dots \subseteq H_\sigma$ leads to a stabilizing sequence of semigroups

$$(3.10) \quad \mathbb{N}\mathcal{A}_\sigma = \mathbb{N}\mathcal{A}_\sigma^0 \subseteq \mathbb{N}\mathcal{A}_\sigma^1 \subseteq \mathbb{N}\mathcal{A}_\sigma^2 \subseteq \dots \subseteq H_\sigma.$$

Namely, one obtains an increasing sequence of $\mathbb{K}[\mathbb{N}\mathcal{A}_\sigma]$ -modules

$$(3.11) \quad \mathbb{K}[\mathbb{N}\mathcal{A}_\sigma] = \mathbb{K}[\mathbb{N}\mathcal{A}_\sigma^0] \subseteq \mathbb{K}[\mathbb{N}\mathcal{A}_\sigma^1] \subseteq \mathbb{K}[\mathbb{N}\mathcal{A}_\sigma^2] \subseteq \dots \subseteq \mathbb{K}[H_\sigma].$$

Essentially by Gordan's lemma, H_σ is finitely generated. Since \mathcal{A} contains elements on the extremal rays of C_σ , $\mathbb{K}[H_\sigma]$ is a finite integral extension of $\mathbb{K}[\mathcal{A}_\sigma]$ and hence a Noetherian $\mathbb{K}[\mathcal{A}_\sigma]$ -module. Thus, (3.11) stabilizes and hence so does (3.10). By finiteness of $\Phi_{\mathcal{A}}^{1,r-1}$, it follows that eventually any new element $b_1 + b_2$ of \mathcal{A}' suggested by Step 7 is already in $\mathbb{N}\mathcal{A}'_\sigma$ for some $\sigma \in \Phi_{\mathcal{A}}^{1,r-1}$.

Suppose the algorithm has reached this stage and let $e_{b_1} + e_{b_2} \in \mathcal{B}$. By Remark 3.10.(2) and Step 4, b_1, b_2 do not share a $\mathbf{1}$ -facet cone. By Lemma 3.4, with $a'' = b_1 + b_2$,

$$\deg_L(e_{b_1} + e_{b_2}) = \deg_L(e_{b_1}) + \deg_L(e_{b_2}) = \deg_{\mathcal{A}}(b_1) + \deg_{\mathcal{A}}(b_2) > \deg_{\mathcal{A}}(a'').$$

By the stability hypothesis on (3.10), $a'' \in \mathbb{N}\mathcal{A}'_\sigma$ for some $\sigma \in \Phi_{\mathcal{A}}^{1,r-1}$ and one can write $a'' = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} a'$ where

$$\deg_{\mathcal{A}}(e_{a''}) = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} \deg_{\mathcal{A}}(e_{a'}) = \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} \deg_L(e_{a'}) = \deg_L(e_{a''})$$

by Lemma 3.4 again. Thus, $(e_{b_1} + e_{b_2}, \sum_{a' \in \mathcal{A}'_\sigma} k_{a'} e_{a'}) \in E_{\mathcal{A}'}$ with L -leading term $e_{b_1} + e_{b_2}$ in contradiction to Step 6. We conclude that $\mathcal{B} = \emptyset$ and the algorithm terminates.

We now prove that the algorithm computes what we want. So we assume that \mathcal{B} has been reduced to the empty set and we let from now on \mathcal{A}' denote the stable value of the generating set for H that forms the output of Algorithm 3.9.

Lemma 3.11. *If $\mathcal{B} = \emptyset$ then for $a'_1, a'_2 \in \mathcal{A}'$ the following are equivalent:*

- (1) $e_{a'_1} + e_{a'_2}$ is an L -leading term of $E_{\mathcal{A}'}$.
- (2) a'_1, a'_2 do not share a $\mathbf{1}$ -facet cone.

In particular, $\mathbb{N}\mathcal{A}'_\sigma = H_\sigma$ for all $\sigma \in \Phi_{\mathcal{A}}^{1,r-1}$.

Proof. In order to show the announced equivalence it suffices by Corollary 3.8 to show that condition 3.8.(1) implies condition 3.11.(1). So assume that $k(e_{a'_1} + e_{a'_2})$ is an L -leading term of $E_{\mathcal{A}'}$, k being minimal in that respect. By Corollary 3.8, a'_1 and a'_2 do not share a $\mathbf{1}$ -facet cone and hence $e_{a'_1} + e_{a'_2}$ was added to \mathcal{B} at some point in the algorithm. This is obvious from Step 7 if not both a'_1, a'_2 are in \mathcal{A} and from (2) and (4) in Remark 3.10 otherwise. As $\mathcal{B} = \emptyset$, $e_{a'_1} + e_{a'_2}$ was eliminated from the queue in Step 6 and hence must be an L -leading term of $E_{\mathcal{A}'}$ itself.

To prove the second claim, let $a \in H_\sigma$. Since \mathcal{A}' generates H , one can write $a = \sum_{a' \in \mathcal{A}'} k_{a'} a'$. Of all such expressions pick one for which $\deg_L(\sum_{a' \in \mathcal{A}'} k_{a'} e_{a'})$ is minimized. Suppose a'_1, a'_2 make a contribution to the sum and 3.11.(2) and hence 3.11.(1) holds. Then $a'_1 + a'_2 = \sum_{a'' \in \mathcal{A}'} k_{a''} a''$ with $\deg_L(e_{a'_1} + e_{a'_2}) > \deg_L(\sum_{a'' \in \mathcal{A}'} k_{a''} e_{a''})$. So $\sum_{a' \in \mathcal{A}'} k_{a'} e_{a'} - e_{a'_1} - e_{a'_2} + \sum_{a'' \in \mathcal{A}'} k_{a''} e_{a''}$ maps to a

under $\varphi_{\mathcal{A}'}$ in display (3.1) but has smaller L -degree than $\sum_{a' \in \mathcal{A}'} k_{a'} e_{a'}$. By contradiction, there must be a $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$ for which C_σ contains all terms in $\sum_{a' \in \mathcal{A}'} k_{a'} a'$ and so $a \in \mathbb{N}\mathcal{A}'_\sigma$ as required. \square

Corollary 3.8, Theorem 3.5, Lemma 3.11, and the arguments after Remark 3.10 combine to the main theorem of this section.

Theorem 3.12. *Algorithm 3.9 terminates and is correct.* \square

The following bound for the torsion order k in Corollary 3.8 can be useful in practice, cf. Remark 3.10.(2).

Definition 3.13. We denote by $M(\mathcal{A})$ be the largest absolute value of a maximal minor of a matrix A whose columns are a $\mathbb{Z}H$ basis representation of the elements of \mathcal{A} .

Lemma 3.14. *For any $p \in F_{\mathcal{A}}$, if kp is an L -leading term of $E_{\mathcal{A}}$ for some $k \geq 1$ then also for $k = M(\mathcal{A})$.*

Proof. Proposition 3.6 spells out when t^p is nilpotent in $\text{gr}^L(\mathbb{K}[H])$. We may reduce to the cases $p = e_a$ where $a^L \notin \sigma$ for any $\sigma \in \Phi_{\mathcal{A}}^{L, r-1}$, and $p = e_{a_1} + e_{a_2}$ where $a_1^L, a_2^L \in \sigma$ for no $\sigma \in \Phi_{\mathcal{A}}^{L, r-1}$.

In the first case, a^L lies in the interior of $\Delta_{\mathcal{A}}^L$ and we choose $\sigma \in \Phi_{\mathcal{A}}^{L, r-1}$ with $a \in C_\sigma$. Then there is a subset $\mathcal{B} \subseteq \mathcal{A}$ of r linearly independent elements such that $\mathcal{B}^L = \{b^L \mid b \in \mathcal{B}\} \subseteq \sigma$ and $a \in \mathbb{Q}_+\mathcal{B}$. Let B be a matrix whose columns are the elements of \mathcal{B} , denote the adjoint matrix of B by \check{B} , and let $k = \det(B) \neq 0$. Then $ka = B\check{B}a$ implies that $ka = \sum_{b \in \mathcal{B}} k_b b$ for some $k_b \in \mathbb{Z}$. Since $ka \in \mathbb{Q}_+\mathcal{B}$ and since \mathcal{B} is linearly independent it follows that each $k_b \in \mathbb{N}$. Rewriting this as $kL_a a^L = \sum_{b \in \mathcal{B}} k_b L_b b^L$, the above conditions on a^L imply that

$$\deg_L(ke_a) = kL_a > \sum_{b \in \mathcal{B}} k_b L_b = \deg_L\left(\sum_{b \in \mathcal{B}} k_b e_b\right).$$

Thus, $(ke_a, \sum_{b \in \mathcal{B}} k_b e_b) \in E_{\mathcal{A}}$ and ke_a is an L -leading term of $E_{\mathcal{A}}$.

The proof of the second case is analogous: the convex combination $\frac{L_1 a_1^L + L_2 a_2^L}{L_1 + L_2} = \frac{a_1 + a_2}{L_1 + L_2}$ of a_1^L and a_2^L lies in the interior of $\Delta_{\mathcal{A}}^L$. As above, choose a linearly independent subset $\mathcal{B} \subseteq \mathcal{A}$ such that $a_1 + a_2 \in \mathbb{Q}_+\mathcal{B}$. This time, an equality

$$(L_1 + L_2)k \frac{L_1 a_1^L + L_2 a_2^L}{L_1 + L_2} = k(a_1 + a_2) = \sum_{b \in \mathcal{B}} k_b b = \sum_{b \in \mathcal{B}} k_b L_b b^L,$$

with $k_b \in \mathbb{N}$, implies that $\deg_L(k(e_{a_1} + e_{a_2})) > \deg_L(\sum_{b \in \mathcal{B}} k_b e_b)$ and hence that $k(e_{a_1} + e_{a_2})$ is an L -leading term of $E_{\mathcal{A}}$.

By Definition 3.13, $M(\mathcal{A}) \geq \det(B) = k$ which completes the proof. \square

Example 3.15. Consider the affine semigroup H generated by

$$\mathcal{A} = \{a_1, \dots, a_6\} = \left\{ \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 3 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 2 \\ 1 \end{pmatrix} \right\}.$$




FIGURE 2. \mathcal{A}' as generated by Algorithm 3.9 applied to Example 3.15 and its central projection onto the $(\mathcal{A}, \mathbf{1})$ -umbrella

We follow the steps in Algorithm 3.9. Using SINGULAR [GPS05], we compute a $\mathbf{1}$ -Gröbner basis

$$\begin{aligned} y_2y_3 - y_1y_4, \quad y_4^2y_5 - y_1^2y_6, \quad y_3y_4y_5 - y_2y_6, \quad y_3^2y_5 - y_1y_6, \\ y_1y_3^2 - y_4^2, \quad y_1^2y_3 - y_2y_4, \quad y_1^3 - y_2^2 \end{aligned}$$

of $I_{\mathcal{A}}$ and thus a Gröbner basis

$$(3.12) \quad y_2y_3 - y_1y_4, \quad y_4^2y_5 - y_1^2y_6, \quad y_3y_4y_5, \quad y_3^2y_5, \quad y_1y_3^2, \quad y_1^2y_3, \quad y_1^3$$

of $\text{gr}^{\mathbf{1}}(I_{\mathcal{A}})$. Obviously, $e_{a_3} + e_{a_5} \in \mathcal{B}$ after Step 1 while $e_{a_1} + e_{a_3}$ is not added to \mathcal{B} since y_3^3 is a $\mathbf{1}$ -leading term. To complete Step 1, compute $M(\mathcal{A}) = 6$ and reduce all 6-th powers of y_iy_j and y_k with respect to the Gröbner basis (3.12) to find

$$\mathcal{B} = \{e_{a_2} + e_{a_3}, e_{a_3} + e_{a_5}, e_{a_4} + e_{a_5}\}.$$

By the rule in Step 2,

$$\Phi_{\mathcal{A}}^{\mathbf{1}, r-1} = \{\sigma_1, \sigma_2, \sigma_3\} = \{\{a_2, a_4, a_6\}, \{a_2, a_5, a_6\}, \{a_3, a_4, a_6\}\}$$

and the corresponding linear forms from Step 3 are

$$\ell_1 = \left(\frac{1}{3}, \frac{2}{3}, -\frac{1}{3}\right), \quad \ell_2 = \left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\right), \quad \ell_3 = (0, 1, -1).$$

Note that $2a_2 = 3a_1$, and hence $a_1 \in C_{\sigma_1} \cap C_{\sigma_2}$. According to Step 4, we update

$$\mathcal{B} = \{e_{a_1} + e_{a_3}, e_{a_2} + e_{a_3}, e_{a_3} + e_{a_5}, e_{a_4} + e_{a_5}\}.$$

In Step 5, we compute \mathcal{A} -degrees of $\mathcal{A}' = \mathcal{A}$ to obtain the weight vector

$$L = \left(\frac{2}{3}, 1, 1, 1, 1, 1\right).$$

Using SINGULAR [GPS05], we compute an L -Gröbner basis

$$\begin{aligned} y_2y_3 - y_1y_4, \quad y_1^3 - y_2^2, \quad y_1^2y_3 - y_2y_4, \quad y_1y_3^2 - y_4^2, \\ y_4^2y_5 - y_1^2y_6, \quad y_3y_4y_5 - y_2y_6, \quad y_3^2y_5 - y_1y_6 \end{aligned}$$

of $I_{\mathcal{A}'}$ and thus a Gröbner basis

$$y_2y_3, \quad y_1^3 - y_2^2, \quad y_1^2y_3, \quad y_1y_3^2, \quad y_4^2y_5, \quad y_3y_4y_5, \quad y_3^2y_5$$

of $\text{gr}^L(I_{\mathcal{A}'})$ that reduces \mathcal{B} in Step 6 to

$$\mathcal{B} = \{e_{a_1} + e_{a_3}, e_{a_3} + e_{a_5}, e_{a_4} + e_{a_5}\}.$$

Following Step 7, we set

$$a'_7 = a_1 + a_3 = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}, \quad a'_8 = a_3 + a_5 = \begin{pmatrix} 2 \\ 1 \\ 1 \end{pmatrix}, \quad a'_9 = a_4 + a_5 = \begin{pmatrix} 3 \\ 1 \\ 1 \end{pmatrix}$$

and update

$$\begin{aligned} \mathcal{A}' &= \mathcal{A} \cup \{a'_7, a'_8, a'_9\}, \\ L &= \left(\frac{2}{3}, 1, 1, 1, 1, 1, \frac{4}{3}, \frac{4}{3}, \frac{5}{3} \right), \\ \mathcal{B} &= \{e_{a_1} + e_{a_3}, e_{a_3} + e_{a_5}, e_{a_3} + e_{a_7}, e_{a_3} + e_{a_8}, e_{a_3} + e_{a_9}, \\ &\quad e_{a_4} + e_{a_5}, e_{a_4} + e_{a_8}, e_{a_4} + e_{a_9}, e_{a_5} + e_{a_7}, e_{a_7} + e_{e_8}, e_{a_7} + e_{a_9}\}. \end{aligned}$$

The next Gröbner basis

$$(3.13) \quad \begin{aligned} &y_1y_3, y_4y_5, y_3y_5, y_2y_4 - y_1y_7, y_2y_3, y_1^3 - y_2^2, y_4y_8, y_3y_8, y_2y_8 - y_1y_9, \\ &y_5y_7, y_3y_7, y_1^2y_4 - y_2y_7, y_4y_9, y_3y_9, y_7y_8, y_1^2y_8 - y_2y_9, y_1y_5y_6 - y_8^2, \\ &y_1y_4^2 - y_7^2, y_7y_9, y_2y_5y_6 - y_8y_9, y_1y_8^2 - y_9^2, y_8^4 - y_5y_6y_9^2 \end{aligned}$$

of $\text{gr}^L(I_{\mathcal{A}'})$ in Step 6 reduces \mathcal{B} to zero and the procedure terminates. The relations (3.13) define a presentation of $\text{gr}^{\mathcal{A}'}(\mathbb{K}[H])$ as quotient of $\mathbb{K}[y_1, \dots, y_9]$.

4. COHEN–MACAULAYNESS OF THE NEWTON GRADED TORIC RING

In this section we investigate Cohen–Macaulayness of the Newton graded ring $\text{gr}^{\mathcal{A}'}(\mathbb{K}[H])$. For $1 \leq k \leq r$, let

$$A_k = A_k(H) = \bigoplus_{\tau \in \Phi_{\mathcal{A}'}^{1, k-1}} \mathbb{K}[H_{\tau}];$$

in particular,

$$A_r = \bigoplus_{\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}} \mathbb{K}[H_{\sigma}].$$

By Lemmas 3.4 and 2.1, for $\tau \in \Phi_{\mathcal{A}'}^1$, $\mathbb{K}[H \setminus C_{\tau}]$ is an ideal in $\text{gr}^{\mathcal{A}'}(\mathbb{K}[H])$ and therefore

$$\text{gr}^{\mathcal{A}'}(\mathbb{K}[H]) \twoheadrightarrow \text{gr}^{\mathcal{A}'}(\mathbb{K}[H])/\mathbb{K}[H \setminus C_{\tau}] = \mathbb{K}[H_{\tau}] \subseteq \text{gr}^{\mathcal{A}'}(\mathbb{K}[H])$$

are maps of \mathbb{K} -algebras. In particular, for $\sigma \in \Phi_{\mathcal{A}'}^{1, r-1}$,

$$(4.1) \quad \mathfrak{m} \cdot \mathbb{K}[H_{\sigma}] = \mathfrak{m}_{\sigma}$$

and hence, for $1 \leq k \leq r$,

$$H_{\mathfrak{m}}^i(A_k) = \bigoplus_{\tau \in \Phi_{\mathcal{A}'}^{1, k-1}} H_{\mathfrak{m}_{\tau}}^i \mathbb{K}[H_{\tau}]$$

for all i . By [Kou76, Prop. 2.6], there is an exact sequence

$$(4.2) \quad 0 \rightarrow \text{gr}^{\mathcal{A}'}(\mathbb{K}[H]) \rightarrow A_r \rightarrow A_{r-1} \rightarrow \dots \rightarrow A_1 \rightarrow 0$$

in the category of $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ -modules. This sequence expresses $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ as the product of the rings $\mathbb{K}[H_{\sigma}]$ over facets $\sigma \in \dot{\Phi}_{\mathcal{A}}^{1,r-1}$ modulo the identification of border rings.

Applying the local cohomology functor $R\Gamma_{\mathfrak{m}}$ to (4.2), one obtains a convergent spectral sequence

$$(4.3) \quad E_1^{p,q} = H_{\mathfrak{m}}^q(A_{r-p}) \implies H_{\mathfrak{m}}^{r-p+q}(\text{gr}^{\mathcal{A}}(\mathbb{K}[H])).$$

It follows that if each $\mathbb{K}[H_{\tau}]$, $\tau \in \dot{\Phi}_{\mathcal{A}}^1$, is Cohen–Macaulay then the spectral sequence collapses on the E_1 -page and $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ must also be Cohen–Macaulay as was already shown in [Kou76] with a slightly different argument.

The local cohomology approach facilitates the search for examples where $\mathbb{K}[H]$ is Cohen–Macaulay but where $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ is not Cohen–Macaulay and where hence not all of the $\mathbb{K}[H_{\tau}]$ can be Cohen–Macaulay themselves. First, observe that the multi-degrees of A_k and hence those of $R\Gamma_{\mathfrak{m}}(A_k)$ are contained in

$$Q_k = \bigcup_{\tau \in \dot{\Phi}_{\mathcal{A}}^{1,k-1}} \mathbb{Q}\tau.$$

Next, suppose that $\mathbb{K}[H]$ is of dimension $r = 2$. Then the spectral sequence degenerates into a long exact sequence of $\mathbb{Z}H$ -graded modules which, since each $\mathbb{K}[H_{\tau}]$ is a domain, starts with

$$0 \rightarrow H_{\mathfrak{m}}^1(\text{gr}^{\mathcal{A}}(\mathbb{K}[H])) \rightarrow H_{\mathfrak{m}}^1(A_2) \rightarrow H_{\mathfrak{m}}^1(A_1) \rightarrow \dots$$

Thus, if one can find a Cohen–Macaulay ring $\mathbb{K}[H]$ where the multi-degrees of $H_{\mathfrak{m}}^1(A_2)$ are not contained in Q_1 then $H_{\mathfrak{m}}^1(\text{gr}^{\mathcal{A}}(\mathbb{K}[H]))$ cannot be zero and therefore $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ not Cohen–Macaulay.


In order to arrive at such example one can start with a two-dimensional cone C_H of a semigroup H where $\mathbb{K}[H]$ is not Cohen–Macaulay. Then one strategically adds new elements outside C_H to H to make $H = \bar{H}$, cf. Corollary 2.3. We illustrate the idea by an example.

Example 4.1. Let H be the affine semigroup generated by

$$\mathcal{A} = \{a_1, \dots, a_4\} = \left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 4 \end{pmatrix} \right\}$$

so that $\mathbb{K}[H]$ is the coordinate ring of the pinched rational quartic. As is well-known, $H_{\mathfrak{m}}^1(\mathbb{K}[H])$ is the one-dimensional vector space spanned by the class of the 1-cocycle $(y_2^2/y_1, y_3^2/y_4)$ in the Čech complex to y_1 and y_4 , cf. Example 2.5. The multi-degree of this generator equals $(1, 2)$ and $\{(1, 2)\} = \bar{H} \setminus H$.

FIGURE 3. The $(\mathcal{A}, \mathbf{1})$ -polyhedron for Example 4.1



We are interested in a new element $a' \in \mathbb{Z}^2 \setminus C_H$ such that the toric ring $\mathbb{K}[H']$ of $H' = H + \mathbb{N}a'$ becomes Cohen–Macaulay. Note that it will be a domain regardless of the choice of a' . By Proposition 2.3.(4), a' has to be chosen such that $(1, 2) + \mathbb{N}a'$ does not meet H' . If, for instance, $a' = (0, k)$ with $k \geq 3$ then pairs of elements in $C_H \cap \mathbb{Z}^2$ that differ by a multiple of a' are either both inside or both outside of H . With such a' we have therefore $(1, 2) \notin H' - \mathbb{N}a'$ and hence $(1, 2) \notin \bar{H}'$. One verifies that $\bar{H}' = H'$, or equivalently

$$H_{\mathfrak{m}_{H'}}^1(\mathbb{K}[H']) = (\mathbb{K}[H' - \mathbb{N}a_1] \cap \mathbb{K}[H' - \mathbb{N}a']) / \mathbb{K}[H'] = 0,$$

and hence $\mathbb{K}[H']$ is Cohen–Macaulay. However the summand $\mathbb{K}[H]$ of $A_2(H')$ is not Cohen–Macaulay.

It is natural to wonder whether the spectral sequence (4.3) always degenerates at the E_1 -page. This would be tantamount to the equivalence

$$[\forall \tau \in \check{\Phi}_{\mathcal{A}}^1 : \mathbb{K}[H_\tau] \text{ is Cohen–Macaulay}] \Leftrightarrow [\text{gr}^{\mathcal{A}}(\mathbb{K}[H]) \text{ is Cohen–Macaulay}].$$

The purpose of the following example is to show that this is not the case.

Example 4.2. Let H be the affine semigroup generated by


$$\mathcal{A} = \{a_1, \dots, a_6\} = \left\{ \begin{pmatrix} 0 \\ 2 \end{pmatrix}, \begin{pmatrix} 0 \\ 3 \end{pmatrix}, \begin{pmatrix} -1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 2 \end{pmatrix}, \begin{pmatrix} -2 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right\}.$$

There are two facet cones $C_\sigma = \mathbb{Q}_+a_1 + \mathbb{Q}_+a_3$ and $C_{\sigma'} = \mathbb{Q}_+a_1 + \mathbb{Q}_+a_6$ where

$$\mathbb{K}[H_{\sigma'}] \cong \mathbb{K}[y_1, y_2, y_6] / \langle y_1^3 - y_2^2 \rangle$$

is a complete intersection. On the other hand, the coordinate transformation

FIGURE 4. The $(\mathcal{A}, \mathbf{1})$ -polyhedron for Example 4.2



$(k_1, k_2) \mapsto (k_1, k_1 + k_2)$ in \mathbb{Z}^2 reveals that

$$\mathbb{K}[H_\sigma] \cong \mathbb{K}[y_1, y_2, y_3, y_4] / \langle y_3^2 y_1 - y_4^2, y_3^3 y_2 - y_4^3, y_1^3 - y_2^2, y_2 y_3 - y_4 y_1 \rangle$$

which is not Cohen–Macaulay since $H_{\mathfrak{m}_\sigma}^1(\mathbb{K}[H_\sigma])$ is the one-dimensional \mathbb{K} -vector space generated by the Čech cocycle

$$(y_2/y_1) \oplus (y_4/y_3) \in \mathbb{K}[H_\sigma - \mathbb{N}a_1] \oplus \mathbb{K}[H_\sigma - \mathbb{N}a_3].$$

Let $\tau = \sigma \cap \sigma'$, then the natural map $H_{\mathfrak{m}_\sigma}^1(\mathbb{K}[H_\sigma]) \rightarrow H_{\mathfrak{m}_\tau}^1(\mathbb{K}[H_\tau])$ induced by the map $\mathbb{K}[H_\sigma] \rightarrow \mathbb{K}[H_\tau]$ of $\mathbb{K}[H_\sigma]$ -modules is injective since

$$H_{\mathfrak{m}_\tau}^1(\mathbb{K}[H_\tau]) = \mathbb{K}[H_\tau - \mathbb{N}a_1]/\mathbb{K}[H_\tau] \cong \mathbb{K} \cdot (y_2/y_1) \oplus \bigoplus_{k \geq 1} \mathbb{K} \cdot (y_2/y_1^2)^k$$

as one immediately verifies. It follows that in the spectral sequence (4.3) the only potentially non zero differential $d_1^{0,1}: E_1^{0,1} \rightarrow E_1^{1,1}$ is in fact injective. Thus the E_2 -page has only terms on the diagonal $p+q=2$ and it follows that $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ is Cohen–Macaulay while the summand $\mathbb{K}[H_\sigma]$ of $A_2(H)$ is not.

Remark 4.3. Using the commands `dim` and `depth` in SINGULAR [GPS05], one can verify that $\mathbb{K}[H]$ in Example 3.15 is Cohen–Macaulay while $\text{gr}^{\mathcal{A}}(\mathbb{K}[H])$ is not. By Proposition 2.3, $\mathbb{K}[H_{\sigma_2}]$ is not Cohen–Macaulay because $a_2 - a_1 = (1, 0, 0) \in \bar{H}_{\sigma_2} \setminus H_{\sigma_2}$. Here we choose $\mathcal{B} = \{a_1, a_4\}$ in Definition 2.2 and observe that $a_2 - a_1 + a_4 = a_7$. However, one can verify as in Example 2.5 that $a_2 - a_1$ is not an actual obstruction to the Cohen–Macaulayness of $\mathbb{K}[H]$.

From [Kou76] the following implications are known:

$$\begin{aligned} \mathbb{K}[H_\tau] \text{ is Cohen–Macaulay } \forall \tau \in \dot{\Phi}_{\mathcal{A}}^1 \\ \implies \text{gr}^{\mathcal{A}}(\mathbb{K}[H]) \text{ is Cohen–Macaulay} \\ \implies \mathbb{K}[H] \text{ is Cohen–Macaulay.} \end{aligned}$$

The examples in this section show that neither implication can be reversed in general. In particular, the proofs for [Oku06b, Prop. 3.1] and [Oku06a, Prop. 3.4] are faulty. However, the statements of these propositions themselves are correct by [MMW05, Cor. 9.2] which is based on vanishing theorems for Euler–Koszul homology on toric modules.

In the sequel we give a Newton filtration proof for [Oku06b, Prop. 3.1] and [Oku06a, Prop. 3.4]. This generalizes the approach of [Ado99, Thm. 1.2] in the case of normal H .

By positivity of H there is a linear form $\ell: \mathbb{Z}H \rightarrow \mathbb{Z}$ such that $\ell(\mathcal{A}) \subseteq \mathbb{N} \setminus \{0\}$.

Lemma 4.4. *For a suitable $q \in \mathbb{N} \setminus \{0\}$, the affine hyperplane $P_q = \ell^{-1}(q)$ intersects the one-dimensional faces of C_H in elements of H .*

Proof. Let $q = \text{lcm}(\ell(\mathcal{A})) > 0$. Then $\ell(qa/\ell(a)) = q$ and $qa/\ell(a) \in \mathbb{N}a$. Thus $P_q \cap \mathbb{N}a \neq \emptyset$ for all $a \in \mathcal{A}$ and the claim follows. \square

Note that natural multiples of the q from Lemma 4.4 also satisfy the conclusion of the lemma. As H is Noetherian, there is $\ell: \mathbb{Q}H \rightarrow \mathbb{Q}$ with $\ell(H) \geq 0$, satisfying Lemma 4.4 with $q = 1$, for which

$$\mathcal{A} = \{h \in H \mid \ell(h) \leq 1\}$$

is a finite set of generators of H . Thus, $\Delta_{\mathcal{A}}^1 = C_H \cap \ell^{-1}([0, 1])$ and $\Phi_{\mathcal{A}}^{1, r-1} = \{C_H \cap \ell^{-1}(1)\}$ is a singleton. In this case, Algorithm 3.9 terminates trivially with $\mathcal{A}' = \mathcal{A}$ returning the Newton weight vector

$$L = (\deg_{\mathcal{A}}(a))_{a \in \mathcal{A}} = (\ell(a))_{a \in \mathcal{A}}.$$

Then, by Lemma 3.4,

$$\text{gr}^L(\mathbb{K}[F_{\mathcal{A}}]/I_{\mathcal{A}}) = \text{gr}^{\mathcal{A}}(\mathbb{K}[H]) \cong \mathbb{K}[H].$$

For any other set \mathcal{A}' of generators for $\mathbb{K}[H]$ with corresponding presentation $\mathbb{K}[F_{\mathcal{A}'}] \rightarrow \mathbb{K}[H]$ one can define a positive weight vector L' on \mathcal{A}' by $L'_{a'} = \deg_{\mathcal{A}'}(a')$. Then $\Phi_{\mathcal{A}'}^{L', r-1}$ is the same singleton $\{C_H \cap \ell^{-1}(1)\}$ as $\Phi_{\mathcal{A}}^{1, r-1}$, $\text{gr}^{L'}(\mathbb{K}[F_{\mathcal{A}'}]/I_{\mathcal{A}'}) \cong \mathbb{K}[H]$.

The preceding arguments can be summarized as follows.

Proposition 4.5. *On any presentation (3.6) of a positive semigroup H , there is a strictly positive weight vector L such that $\text{gr}^L(\mathbb{K}[F_{\mathcal{A}}]/I_{\mathcal{A}}) \cong \mathbb{K}[H]$. \square*

We return to the A -hypergeometric system $M_A(\beta)$ defined in the introduction where A is an integer $d \times n$ matrix and β a complex d -vector. The leading actor in the homological study of the A -hypergeometric system is the *Euler–Koszul complex* $K_{\bullet}(S_A, \beta)$ whose 0-th homology module is $M_A(\beta)$ [MMW05]. For a \mathbb{Z}^d -graded R_A -module N , $K_{\bullet}(N, \beta)$ is by definition the Koszul complex of the d commuting left D_A -endomorphisms $E - \beta$ on $D_A \otimes_{R_A} N$ defined by $E_i \circ y = (E_i - \deg_i(y))y$ for all \mathbb{Z}^d -homogeneous $y \in M$ and by \mathbb{C} -linear extension.

Corollary 4.6. *Assume that the columns of $A \in \mathbb{C}^{d \times n}$ span a positive affine semigroup $\mathbb{N}A$ such that $\mathbb{Z}A = \mathbb{Z}^d$ and that $S_A = \mathbb{C}[\mathbb{N}A]$ is a Cohen–Macaulay ring. Then the Euler–Koszul complex $K_{\bullet}(S_A, \beta)$ is acyclic for all $\beta \in \mathbb{C}^d$.*

Proof. By Proposition 4.5, there is a strictly positive weight vector L on ∂ such that $\text{gr}^L(R_A/I_A) \cong S_A$. Choose a rational $c > 0$ and set $\deg_L(x_j) = c - \deg_L(\partial_j)$ for all j . One obtains an induced filtration on D_A with $L_k(R_A) = L_k(D_A) \cap R_A$ satisfying $\text{gr}^L(D_A) = \mathbb{C}[x, \partial]$, and so

$$(4.4) \quad \text{gr}^L(D_A \otimes_{R_A} S_A) = \mathbb{C}[x] \otimes_{\mathbb{C}} S_A.$$

The L -symbols E of the Euler operators $E - \beta$ form a system of parameters on (4.4), cf. [GZK89]. As S_A is Cohen–Macaulay by hypothesis, the Koszul complex of E on (4.4) is a resolution. The latter being the L -graded Euler–Koszul complex $\text{gr}^L(K_{\bullet}(S_A, \beta))$, acyclicity of the ungraded complex $K_{\bullet}(S_A, \beta)$ follows. \square

Remark 4.7. Note the “limit case” $c \rightarrow \infty$, which corresponds to the weight filtration on D_A given by $(\mathbf{1}, \mathbf{0})$. This filtration has been curiously overlooked as an elementary tool to show exactness of the Euler–Koszul complex for Cohen–Macaulay S_A .

REFERENCES

- [Ado94] Alan Adolphson, *Hypergeometric functions and rings generated by monomials*, Duke Math. J. **73** (1994), no. 2, 269–290. MR **96c**:33020
- [Ado99] ———, *Higher solutions of hypergeometric systems and Dwork cohomology*, Rend. Sem. Mat. Univ. Padova **101** (1999), 179–190. MR **MR1705287 (2001b**:14032)
- [BS98] M. P. Brodmann and R. Y. Sharp, *Local cohomology: an algebraic introduction with geometric applications*, Cambridge Studies in Advanced Mathematics, vol. 60, Cambridge University Press, Cambridge, 1998. MR **MR1613627 (99h**:13020)
- [GGZ87] I. M. Gelfand, M. I. Graev, and A. V. Zelevinskiĭ, *Holonomic systems of equations and series of hypergeometric type*, Dokl. Akad. Nauk SSSR **295** (1987), no. 1, 14–19. MR **MR902936 (88j**:58118)
- [GPS05] G.-M. Greuel, G. Pfister, and H. Schönemann, *SINGULAR 3.0*, A Computer Algebra System for Polynomial Computations, Centre for Computer Algebra, University of Kaiserslautern, 2005, <http://www.singular.uni-kl.de>.
- [GZK89] I. M. Gelfand, A. V. Zelevinskiĭ, and M. M. Kapranov, *Hypergeometric functions and toric varieties*, Funktsional. Anal. i Prilozhen. **23** (1989), no. 2, 12–26. MR **90m**:22025

- [Hoc72] M. Hochster, *Rings of invariants of tori, Cohen-Macaulay rings generated by monomials, and polytopes*, Ann. of Math. (2) **96** (1972), 318–337. MR MR0304376 (46 #3511)
- [HT88] Lê Tuấn Hoa and Ngô Việt Trung, *Corrigendum to: “Affine semigroups and Cohen-Macaulay rings generated by monomials”* [Trans. Amer. Math. Soc. **298** (1986), no. 1, 145–167; MR0857437 (87j:13032)], Trans. Amer. Math. Soc. **305** (1988), no. 2, 857. MR MR924781 (88m:13008)
- [Kou76] A. G. Kouchnirenko, *Polyèdres de Newton et nombres de Milnor*, Invent. Math. **32** (1976), no. 1, 1–31. MR MR0419433 (54 #7454)
- [M2] Daniel R. Grayson and Michael E. Stillman, *Macaulay 2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [MMW05] Laura Felicia Matusevich, Ezra Miller, and Uli Walther, *Homological methods for hypergeometric families*, J. Amer. Math. Soc. **18** (2005), no. 4, 919–941 (electronic). MR MR2163866
- [Oku06a] Go Okuyama, *A-hypergeometric ranks for toric threefolds*, Int. Math. Res. Not. (2006), Art. ID 70814, 38. MR MR2250006
- [Oku06b] ———, *Local cohomology modules of A-hypergeometric systems of Cohen-Macaulay type*, Tohoku Math. J. (2) **58** (2006), no. 2, 259–275. MR MR2248433
- [SW06] M. Schulze and U. Walther, *Irregularity of hypergeometric systems via slopes along coordinate subspaces*, arXiv.org **math.AG** (2006), no. 0608668, To appear in Duke Math. J.
- [TH86] Ngô Việt Trung and Lê Tuấn Hoa, *Affine semigroups and Cohen-Macaulay rings generated by monomials*, Trans. Amer. Math. Soc. **298** (1986), no. 1, 145–167. MR MR857437 (87j:13032)

M. SCHULZE, OKLAHOMA STATE UNIVERSITY, DEPT. OF MATHEMATICS, 401 MSCS, STILLWATER, OK 74078, USA

E-mail address: mschulze@math.okstate.edu

U. WALTHER, PURDUE UNIVERSITY, DEPT. OF MATHEMATICS, 150 N. UNIVERSITY ST., WEST LAFAYETTE, IN 47907, USA

E-mail address: walther@math.purdue.edu