


- Least Square > Projection

$$A\vec{x} = \vec{b} \quad A \in \mathbb{R}^{m \times n} \quad m \begin{matrix} n \\ \boxed{A} \end{matrix} \quad m > n$$



$$A\vec{x} = \vec{b}$$

$$\Rightarrow A^T A \hat{x} = A^T \vec{b}$$

$$\Rightarrow \hat{x} = (A^T A)^{-1} A^T \vec{b}$$

$$\Rightarrow A\hat{x} = A(A^T A)^{-1} A^T \vec{b}$$

- ① In order for $(A^T A)^{-1}$ to exist, A need to have independent column vectors.

If we want to project a vector \vec{b} onto a subspace or a span, find a basis for that subspace. Use basis column vectors to form A.

- ② If $A = [\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3]$ has orthogonal columns,

$$\text{then } A^T A = \begin{matrix} \text{---} \\ \text{---} \\ \text{---} \end{matrix} \begin{matrix} | \\ | \\ | \end{matrix} = \begin{bmatrix} \|\vec{v}_1\|^2 & 0 & 0 \\ 0 & \|\vec{v}_2\|^2 & 0 \\ 0 & 0 & \|\vec{v}_3\|^2 \end{bmatrix}$$

\Rightarrow The projection of \vec{b} onto $\text{Col}(A)$ is also

$$\left[A(A^T A)^{-1} A^T \vec{b} = \right] \frac{\langle \vec{b}, \vec{v}_1 \rangle}{\|\vec{v}_1\|^2} \vec{v}_1 + \frac{\langle \vec{b}, \vec{v}_2 \rangle}{\|\vec{v}_2\|^2} \vec{v}_2 + \frac{\langle \vec{b}, \vec{v}_3 \rangle}{\|\vec{v}_3\|^2} \vec{v}_3$$

Example: Projection of \vec{b} onto $\text{Span}\{\vec{v}\}$ is $\frac{\langle \vec{b}, \vec{v} \rangle}{\|\vec{v}\|^2} \vec{v}$

③ If $A = [\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3]$ has orthonormal columns,

$$\text{then } A^T A = \begin{array}{|c|c|c|} \hline \text{---} & \text{---} & \text{---} \\ \hline \text{---} & \text{---} & \text{---} \\ \hline \text{---} & \text{---} & \text{---} \\ \hline \end{array} \begin{array}{|c|c|c|} \hline \text{---} & \text{---} & \text{---} \\ \hline \text{---} & \text{---} & \text{---} \\ \hline \text{---} & \text{---} & \text{---} \\ \hline \end{array} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

\Rightarrow The projection of \vec{b} onto $\text{Col}(A)$ is also

$$\left[A(A^T A)^{-1} A^T \vec{b} = \right] \langle \vec{b}, \vec{v}_1 \rangle \vec{v}_1 + \langle \vec{b}, \vec{v}_2 \rangle \vec{v}_2 + \langle \vec{b}, \vec{v}_3 \rangle \vec{v}_3$$

• Gram-Schmit for generating orthogonal vectors

Question: Given $\text{Span} \{ \vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4, \dots, \vec{v}_7 \}$,
how to find an orthogonal basis?

Answer: 1) Find a basis, say, $\{ \vec{v}_1, \vec{v}_2, \vec{v}_3 \}$

2) Apply Gram-Schmit to $\{ \vec{v}_1, \vec{v}_2, \vec{v}_3 \}$

$$\vec{u}_1 = \vec{v}_1$$

$$\vec{u}_2 = \vec{v}_2 - \frac{\langle \vec{v}_2, \vec{u}_1 \rangle}{\| \vec{u}_1 \|^2} \vec{u}_1$$

$$\vec{u}_3 = \vec{v}_3 - \frac{\langle \vec{v}_3, \vec{u}_1 \rangle}{\| \vec{u}_1 \|^2} \vec{u}_1 - \frac{\langle \vec{v}_3, \vec{u}_2 \rangle}{\| \vec{u}_2 \|^2} \vec{u}_2$$

• Solutions to $\frac{d}{dt} \vec{u}(t) = A \vec{u}(t)$

$$\frac{d}{dt} \begin{bmatrix} u_1(t) \\ u_2(t) \\ u_3(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} u_1(t) \\ u_2(t) \\ u_3(t) \end{bmatrix}$$

① Find eigenvalues λ_i with eigenvectors \vec{v}_i

② $\vec{u}(t) = e^{\lambda_1 t} \vec{v}_1$ is one solution

③ If $\{\vec{v}_1, \vec{v}_2, \vec{v}_3\}$ are independent eigenvectors
then $e^{\lambda_1 t} \vec{v}_1, e^{\lambda_2 t} \vec{v}_2, e^{\lambda_3 t} \vec{v}_3$ are
independent solutions (abstract vectors)

④ (IVP)
initial value
Problem

$$\begin{cases} \frac{d}{dt} \begin{bmatrix} u_1(t) \\ u_2(t) \\ u_3(t) \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \begin{bmatrix} u_1(t) \\ u_2(t) \\ u_3(t) \end{bmatrix} \\ \begin{bmatrix} u_1(0) \\ u_2(0) \\ u_3(0) \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix} \end{cases}$$

Assume $A = VDV^{-1}$, then

$$A_t = V [D_t] V^{-1}$$

$$= V \begin{bmatrix} e^{\lambda_1 t} & 0 & 0 \\ 0 & e^{\lambda_2 t} & 0 \\ 0 & 0 & e^{\lambda_3 t} \end{bmatrix} V^{-1}$$

The solution to IVP is

$$\vec{u}(t) = e^{At} \vec{u}(0)$$

$$= V \begin{bmatrix} e^{\lambda_1 t} & 0 & 0 \\ 0 & e^{\lambda_2 t} & 0 \\ 0 & 0 & e^{\lambda_3 t} \end{bmatrix} V^{-1} \vec{u}(0)$$