

GREEN - Test Version 01

NAME _____ INSTRUCTOR _____

1. You must use a **#2 pencil** on the mark-sense sheet (answer sheet).
2. On the mark-sense sheet, fill in the **instructor's** name (if you do not know, write down the class meeting time and location) and the **course number** which is **MA265**.
3. Fill in your **NAME** and blacken in the appropriate spaces.
4. Fill in the **SECTION Number** boxes with section number of your class (see below if you are not sure) and blacken in the appropriate spaces.

172	2:30pm	MWF	Brown, Johnny		173	10:30am	MWF	Chen, Ying
174	10:30am	TR	Ho, Meng-Che		175	12:00pm	TR	Ho, Meng-Che
176	10:30am	TR	Liu, Baiying		177	4:30pm	TR	Liu, Baiying
178	1:30pm	MWF	Liu, Tong		179	10:30am	MWF	Liu, Tong
180	1:30pm	TR	Luo, Tao		181	12:00pm	TR	Luo, Tao
182	4:30pm	TR	Madsen, Caroline		183	3:00pm	TR	Madsen, Caroline
184	12:30pm	MWF	Moon, Yong Suk		185	11:30am	MWF	Moon, Yong Suk
186	3:30pm	MWF	Patz, Peter		187	4:30pm	MWF	Patz, Peter
188	10:30am	MWF	Wang, Xu		189	11:30am	MWF	Wang, Xu
190	9:30am	MWF	Wang, Yating		191	8:30am	MWF	Wang, Yating
192	1:30pm	MWF	Wei, Ning		193	3:30pm	MWF	Wei, Ning
194	9:30am	MWF	Xu, Ping		195	10:30am	MWF	Xu, Ping
196	1:30pm	TR	Yang, Zhiguo		197	3:00pm	TR	Yang, Zhiguo

5. Fill in the correct TEST/QUIZ NUMBER (**GREEN** is 01).
6. Fill in the 10-DIGIT PURDUE ID and blacken in the appropriate spaces.
7. Sign the mark-sense sheet.
8. Fill in your name and your instructor's name on the question sheets (above).
9. There are 25 questions, each worth 8 points. Blacken in your choice of the correct answer in the spaces provided for questions 1–25 in the answer sheet. Do all your work on the question sheets, in addition, also **CIRCLE** your answer choice for each problem on the question sheets in case your scantron is lost. **Turn in both the mark-sense sheets and the question sheets to your instructor when you are finished.**
10. Show your work on the question sheets. Although no partial credit will be given, any disputes about grades or grading will be settled by examining your written work on the question sheets.
11. **NO CALCULATORS, BOOKS, NOTES, PHONES OR CAMERAS ARE ALLOWED** on this exam. Turn off or put away all electronic devices. Use the back of the test pages for scrap paper.

1. The matrix below represents the augmented matrix of a system of linear equations. Assume that the variables in this system are x_1, x_2, x_3, x_4, x_5 , and x_6 , and let A be the coefficient matrix:

ii) '2' is not the only non-zero entry of 5th col \Rightarrow not RREF.

$$\left(\begin{array}{cccccc|c} \boxed{1} & 0 & 1 & 2 & 0 & 0 & 1 \\ 0 & \boxed{1} & 0 & 4 & \boxed{1} & 0 & 2 \\ 0 & 0 & 0 & 0 & \boxed{1} & 0 & c \\ 0 & 0 & 0 & 0 & 0 & \boxed{1} & d \end{array} \right)$$

x_1, x_2 are free b/c (no leading entry)

*1 i. there is one row $[0 \ 0 \ \dots \ 0 \ | \ b]$
 $b \neq 0 \Rightarrow$ system has no solution.*

Which of the following statements are true?

- (i) For any given c and d , the system above is consistent.
- (ii) The coefficient matrix A is in reduced echelon form.
- (iii) The right hand side vector is in the column space of matrix A .
- (iv) The system has no solution.
- (v) The system has infinitely many solutions.

- A. (i), (ii) only
- B. (i), (ii), (iii) only
- C. (ii), (iii), (iv) only
- D. (i), (iii), (v) only
- E. all of the above

*iii RHS vector is in the span of all cols of A
 \Leftrightarrow there exist c_1, c_2, \dots, c_6 all scalars, such that
 $c_1 \vec{x}_1 + c_2 \vec{x}_2 + \dots + c_6 \vec{x}_6 = \vec{b}$
 $\Leftrightarrow (\vec{x}_1 \ \dots \ \vec{x}_6 \ | \ \vec{b})$ has a solution.*

2. Suppose the set $\left\{ \begin{bmatrix} -1 \\ -3 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ -2 \\ a \end{bmatrix}, \begin{bmatrix} -1 \\ -2 \\ 1 \end{bmatrix} \right\}$ is linearly dependent. Find a .

- A. $a = -5$.
- B. $a = -2$.
- C. $a = 2$.
- D. $a = 1$.
- E. $a = -3$.

linear dep: $Ax = 0$ has non-trivial solutions

$\Leftrightarrow A$ has free variables

\Leftrightarrow there exist some cols which have no leading entry.

$$\left(\begin{array}{ccc} -1 & -2 & -1 \\ -3 & -2 & -2 \\ 1 & a & 1 \end{array} \right) \xrightarrow[R_3 \rightarrow 0]{R_2 = R_2 - 3R_1} \left(\begin{array}{ccc} -1 & -2 & -1 \\ 0 & 4 & 1 \\ \boxed{1} & a & 1 \end{array} \right) \xrightarrow[R_1 \rightarrow 0]{R_3 = R_3 + R_1} \left(\begin{array}{ccc} -1 & -2 & -1 \\ 0 & 4 & 1 \\ 0 & a+2 & 0 \end{array} \right)$$

If $a+2 = 0 \Rightarrow$ the 3rd col has no leading entry $\Rightarrow x_3$ is free

\Rightarrow non-trivial solution \Rightarrow linear dep.

$$3^A \vec{v}_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \vec{v}_2 = \begin{pmatrix} 2 \\ 0 \\ 0 \end{pmatrix} \vec{v}_3 = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

$$\Leftrightarrow [\vec{v}_1 \ \vec{v}_2 \ \vec{v}_3 \ | \ \vec{b}]$$

3. Which of the following statements is **false**?

- A. ~~X~~ If $\{v_1, v_2, v_3\}$ is linearly dependent, then v_3 is a linear combination of v_1 and v_2 .
- B. Suppose that the columns of A are \vec{v}_1, \vec{v}_2 , and \vec{v}_3 . Then the matrix equation $A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = b$ is equivalent to the vector equation $x_1\vec{v}_1 + x_2\vec{v}_2 + x_3\vec{v}_3 = b$. *set of the solutions to the $Ax=b$*
- C. Suppose that X_0 is a solution to the linear system $AX = b$. Then $\{X | AX = b\} = X_0 + \{X | AX = 0\}$. (sec 1.6)
- D. The columns of A are linearly independent if and only if A has a pivot position in every column. $\Leftrightarrow A$ has only trivial solution. \Leftrightarrow no free variable
- E. A homogeneous linear system has a non-trivial solution if and only if it has at least one free variable.

4. Let $L : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a linear transformation such that $L(\underbrace{\begin{bmatrix} 1 \\ 2 \end{bmatrix}}_{\vec{u}}) = \underbrace{\begin{bmatrix} 0 \\ 1 \\ 4 \end{bmatrix}}_{\vec{v}}$ and $L(\underbrace{\begin{bmatrix} 3 \\ 2 \end{bmatrix}}_{\vec{w}}) =$

$$\underbrace{\begin{bmatrix} -2 \\ 0 \\ 1 \end{bmatrix}}_{\vec{x}}. \text{ Find } L(\underbrace{\begin{bmatrix} -1 \\ 2 \end{bmatrix}}_{\vec{y}}).$$

A. $\begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix}$

B. $\begin{bmatrix} 2 \\ 2 \\ 7 \end{bmatrix}$

C. $\begin{bmatrix} -2 \\ 4 \\ 5 \end{bmatrix}$

D. $\begin{bmatrix} 0 \\ 6 \\ -3 \end{bmatrix}$

E. $\begin{bmatrix} 5 \\ 3 \\ 7 \end{bmatrix}$

If you can find $c_1, c_2 \in \mathbb{R}$

such that $\vec{x} = c_1\vec{u} + c_2\vec{v}$

$L(\vec{x}) = L(c_1\vec{u} + c_2\vec{v}) \stackrel{\text{property of the linear transformation}}{=} c_1L(\vec{u}) + c_2L(\vec{v})$

Now, $c_1\vec{u} + c_2\vec{v} = \vec{x}$ *vector form*

$\Leftrightarrow [\vec{u}_1, \vec{v} \ | \ \vec{x}]$ *augmented form*

$$\left[\begin{array}{cc|c} 1 & 3 & -1 \\ 2 & 2 & 2 \end{array} \right] \xrightarrow{R_2 = R_2 - 2R_1} \left[\begin{array}{cc|c} 1 & 3 & -1 \\ 0 & -4 & 4 \end{array} \right]$$

Solve from bottom to top.

2nd $-4c_2 = 4 \Rightarrow c_2 = -1$

1st $c_1 + 3c_2 = -1 \Rightarrow c_1 = 2$

$$L(\vec{x}) = 2 \cdot \begin{pmatrix} 0 \\ 1 \\ 4 \end{pmatrix} - \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 2 \\ 2 \\ 7 \end{pmatrix}$$

2

Q3. $A \in \mathbb{R}^{n \times n}$

$Ax = 0$ has only trivial solution

\Leftrightarrow cols of A are linearly indep.

\Leftrightarrow no free variable.

\Leftrightarrow each col has a pivot position (leading entry)

\Leftrightarrow each col is ⁱⁿ the basis of $\text{col}(A)$ (see 2.8 thm)

$\Leftrightarrow \text{dim}(\text{col}(A)) = n$ (# of vectors in the basis. # of cols of A)

$\Leftrightarrow \text{rank}(A) = n \rightarrow \text{rank}(A) = \text{dim}(\text{col}(A))$

i $\text{rank}(A) = m = \# \text{ rows of } A$ (thm 2.9 (rank) \rightarrow # cols)

$\Leftrightarrow \text{dim}(\text{null}(A)) = n - m \rightarrow \text{rank} = \text{dim}(\text{col}(A))$

ii $Ax = 0$ has ^{only} trivial solution.

$\text{null}(A) = \{x \mid Ax = 0\} \Leftrightarrow \text{null}(A) = \{0\} \Leftrightarrow \text{dim}(\text{null}(A)) = 0$

i + ii $\Rightarrow n - m = 0 \Leftrightarrow n = m \Leftrightarrow A$ is a square matrix $\Leftrightarrow \det(A) \neq 0 \Leftrightarrow A$ is invertible $\Leftrightarrow \dots$

R1 $L: \mathbb{R}^m \rightarrow \mathbb{R}^n$ linear. $A = \{U(\vec{e}_1), U(\vec{e}_2), \dots, U(\vec{e}_m)\}$ trivial solution.
 there exists $A_{n \times m}$ such that

$L(x) = A \cdot x_{m \times 1}$

R2 Linear L is 1-1

iff $L(x) = 0$ has only $Ax = 0$

R3 Linear L is 1-1 iff cols of A are linearly indep.

5. Let $L: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation whose standard matrix is $\begin{bmatrix} t-1 & 2t-2 \\ 1 & t \end{bmatrix}$ where t is a real number. Find ALL values of t such that L is one-to-one.

R4 T maps \mathbb{R}^n onto \mathbb{R}^m iff cols of the std A span \mathbb{R}^m .

- A. $t \neq 1$
- B. $t \neq 0, 1$
- C. $t \neq 1, 2$
- D. $t = 1$
- E. $t = 2$

$\begin{pmatrix} t-1 & 2t-2 \\ 1 & t \end{pmatrix} \xrightarrow{R_2 \leftrightarrow R_1} \begin{pmatrix} 1 & t \\ t-1 & 2t-2 \end{pmatrix}$ variables are basic.
 $\Rightarrow t+2$ & $t+1$
 $R_2 = R_2 - R_1 \cdot (t-1)$
 $t-1 \rightarrow 0$
 $\begin{pmatrix} 1 & t \\ 0 & 2(t-1) - (t-1)t \end{pmatrix}$ RREF

iff $2(t-1) - (t-1)t \neq 0, \Rightarrow$ all $(2-t)(t-1) \neq 0$

6. Let $A = \begin{bmatrix} 2 & 0 & 2 \\ 1 & 0 & 2 \\ -3 & 1 & -3 \end{bmatrix}$ and let its inverse $A^{-1} = [b_{ij}]$. Find the trace of the matrix A^{-1} . In other words, compute the sum $b_{11} + b_{22} + b_{33}$.

- A. -1
- B. 0
- C. $\frac{1}{2}$
- D. 1
- E. 2

7. Find the third column of the matrix D , given that $C = \begin{bmatrix} 1 & -1 \\ 3 & 2 \end{bmatrix}$ and $CD = \begin{bmatrix} 2 & 0 & -2 & 1 & 0 \\ 0 & 3 & 1 & 7 & 0 \end{bmatrix}$.

A. $\begin{bmatrix} -\frac{12}{5} \\ \frac{7}{5} \end{bmatrix}$

B. $\begin{bmatrix} -\frac{3}{5} \\ \frac{7}{5} \end{bmatrix}$

C. $\begin{bmatrix} -\frac{2}{5} \\ \frac{1}{5} \end{bmatrix}$

D. $\begin{bmatrix} -5 \\ -7 \end{bmatrix}$

E. $\begin{bmatrix} -7 \\ -5 \end{bmatrix}$

$AB = \{ \underbrace{A \vec{b}_1}_{m \times 1}, A \vec{b}_2, \dots, A \vec{b}_p \}$

each col of AB is a linear combination of cols of A using the weights from the corresponding col of B .

$A \vec{b}_1 = \vec{a}_1 \cdot b_{11} + \vec{a}_2 \cdot b_{21} + \dots + \vec{a}_n \cdot b_{n1}$

$\vec{b}_1 = \begin{pmatrix} b_{11} \\ b_{21} \\ \vdots \\ b_{n1} \end{pmatrix}$

7. sol: the third col of $CD =$ linear combination of cols of C using the weights from the third col of D .

$\begin{pmatrix} -2 \\ 1 \end{pmatrix} = x_1 \begin{pmatrix} 1 \\ 3 \end{pmatrix} + x_2 \begin{pmatrix} -1 \\ 2 \end{pmatrix}$, where 3rd col of $D = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$
 3rd col of CD 3rd col of C 2nd col of C .

Q8. $A = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 3 \end{pmatrix}_{2 \times 5}$, nullity(A) = 3, basis of col(A) = $\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \}$

$A^T = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}_{5 \times 2}$, nullity(A^T) = 0, basis of col(A^T) = $\{ \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \\ 3 \end{pmatrix} \}$

no free variable.

8. Let A be an m x n matrix. Which of the following statement is necessarily true?

- A. The nullity of A is the same as the nullity of A^T. (X)
 - B. The rank of A is the same as the rank of A^T.
 - C. The column space of A is the same as the null space of A^T.
 - D. The columns of A form a basis of the column space of A.
 - E. The columns of A^T form a basis of the null space of A.
- ① linear indep
② span
- nullity = dim(null(A))
dim(col(A)) = 2
dim(null(A^T)) = 0

9. Let

$A = \begin{bmatrix} 1 & 2 & 2 & 2 & 2 \\ 3 & 2 & 3 & 1 & 3 \\ 3 & 1 & 2 & 2 & 2 \end{bmatrix}_{3 \times 5} \cdot \begin{pmatrix} | \\ | \\ | \\ | \\ | \end{pmatrix}_{5 \times 1} = 0$

Which of the following is a basis of the null space of A?

A. $\begin{bmatrix} -4 \\ -8 \\ 9 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ -1 \\ 0 \\ 1 \end{bmatrix}$

B. $\begin{bmatrix} 1 \\ 3 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 2 \end{bmatrix}$

C. $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$

D. $\begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$

E. $\begin{bmatrix} 4 \\ 8 \\ -11 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} -8 \\ -16 \\ 19 \\ 2 \\ -1 \end{bmatrix}, \begin{bmatrix} -4 \\ -8 \\ 0 \\ 1 \\ 9 \end{bmatrix}$

$\begin{pmatrix} 1 & 2 & 2 & 2 & 2 \\ 3 & 2 & 3 & 1 & 3 \\ 3 & 1 & 2 & 2 & 2 \end{pmatrix} \xrightarrow{R_2 - R_1, R_3 - R_1} \begin{pmatrix} 1 & 2 & 2 & 2 & 2 \\ 0 & -4 & -3 & -3 & -3 \\ 0 & -1 & -2 & -2 & -2 \end{pmatrix}$

$\xrightarrow{R_2 - R_3} \begin{pmatrix} 1 & 2 & 2 & 2 & 2 \\ 0 & -4 & -3 & -3 & -3 \\ 0 & -1 & -2 & -2 & -2 \end{pmatrix} \xrightarrow{R_2 - R_3} \begin{pmatrix} 1 & 2 & 2 & 2 & 2 \\ 0 & -3 & -1 & -1 & -1 \\ 0 & -1 & -2 & -2 & -2 \end{pmatrix}$

$\xrightarrow{R_2 - R_3} \begin{pmatrix} 1 & 2 & 2 & 2 & 2 \\ 0 & -2 & 1 & 1 & 1 \\ 0 & -1 & -2 & -2 & -2 \end{pmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{pmatrix} 1 & 2 & 2 & 2 & 2 \\ 0 & -1 & -2 & -2 & -2 \\ 0 & -2 & 1 & 1 & 1 \end{pmatrix}$

3rd row $\Rightarrow x_3 - 9s + t = 0 \Rightarrow x_3 = 9s - t$
 2nd row $\Rightarrow x_2 + (9s - t) - s + t = 0 \Rightarrow x_2 = -8s$
 1st row $\Rightarrow x_1 - 16s + 18s - 2t + 2s + 2t = 0 \Rightarrow x_1 = -4s$

$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} -4s + 0t \\ -8s + 0t \\ 9s - t \\ s + 0t \\ 0s + t \end{pmatrix} = \begin{pmatrix} -4s \\ -8s \\ 9s \\ s \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ -t \\ -t \\ 0 \\ t \end{pmatrix}$
 $= s \begin{pmatrix} -4 \\ -8 \\ 9 \\ 1 \\ 0 \end{pmatrix} + t \begin{pmatrix} 0 \\ -1 \\ -1 \\ 0 \\ 1 \end{pmatrix}$
 basis of null(A)

verify A * (0, 0, 0, 1, 0) = (0, 0, 0, 1, 0)

$A \xrightarrow{\text{transpose}} \begin{pmatrix} a & d & g \\ b & e & h \\ c & f & i \end{pmatrix} \xrightarrow{R_2 = R_2 \cdot 2} \begin{pmatrix} a & d & g \\ 2b & 2e & 2h \\ c & f & i \end{pmatrix} \xrightarrow{\text{transpose}} \begin{pmatrix} a & 2b & c \\ d & 2e & f \\ g & 2h & i \end{pmatrix} \xrightarrow{R_2 \leftrightarrow R_3} \begin{pmatrix} a & 2b & c \\ g & 2h & i \\ d & 2e & f \end{pmatrix}$

$\det(A) = \det(A_1)$
 $\det(A_2) = 2 \det(A) = 8$
 $\det(A_3) = \det(A_2) = 8$
 $\det(A_4) = -\det(A_3) = -8$

10. Suppose a 3×3 matrix $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix}$ has determinant 4. What is the determinant of

$B = \begin{bmatrix} a & 2b & c \\ g & 2h & i \\ d + 3a & 2e + 6b & f + 3c \end{bmatrix}?$

$R_3 = R_3 + 3R_1$
 $\det(B) = \det(A_4) = -8$

- A. 4.
- B. 8.
- C. -8.**
- D. 24.
- E. -24.

11. Compute the value of the following determinant:

$$\begin{vmatrix} 4 & -9 & 2 & 3 \\ 0 & 3 & 0 & -4 \\ -5 & 0 & 0 & 3 \\ 0 & 5 & 0 & -7 \end{vmatrix}$$

- A. 10.
- B. -10.
- C. 410.
- D. -410.
- E. 90.

12. Suppose $A = PDP^{-1}$, where P is a 3×3 invertible matrix and $D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.

Let $B = 2I + 3A + A^2$, which of the following is **true**?

- A. B is not diagonalizable.
- B. B is diagonalizable, and $B = PCP^{-1}$, where $C = \begin{bmatrix} 6 & 0 & 0 \\ 0 & 12 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.
- C. B is diagonalizable, and $B = PCP^{-1}$, where $C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & -3 \end{bmatrix}$.
- D. B is diagonalizable, and $B = PCP^{-1}$ for some C , but there is not enough information to determine C .
- E. There is not enough information to determine whether B is diagonalizable.

13. Which of the following statements are **true**?

(i) If λ is an eigenvalue for A , then $-\lambda$ is an eigenvalue for $-A$.

(ii) If zero is an eigenvalue of A , then A is not invertible.

(iii) If an $n \times n$ matrix A is diagonalizable, then A has n distinct eigenvalues.

(iv) Let $A = \begin{bmatrix} 2 & 1 \\ 0 & 2 \end{bmatrix}$, then A is both invertible and diagonalizable.

A. (i) and (ii) only

B. (i) and (iii) only

C. (i), (ii) and (iii) only

D. (i), (ii) and (iv) only

E. (i), (ii), (iii) and (iv)

14. Define $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $T(x) = Ax$, where $A = \begin{bmatrix} 2 & -6 \\ -1 & 3 \end{bmatrix}$. Which of the following is a basis \mathcal{B} for \mathbb{R}^2 with the property that the \mathcal{B} -matrix for T is a diagonal matrix?

A. $\left\{ \begin{bmatrix} 3 \\ 1 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right\}$

B. $\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 3 \\ 4 \end{bmatrix} \right\}$

C. $\left\{ \begin{bmatrix} 4 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 4 \end{bmatrix} \right\}$

D. $\left\{ \begin{bmatrix} 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\}$

E. $\left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$.

15. Let $A = \begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}$, where $i = \sqrt{-1}$. Then A^{32} equals

- A. $\begin{bmatrix} 1 & i \\ 1 & i \end{bmatrix}$
- B. $\begin{bmatrix} i & 0 \\ 0 & i \end{bmatrix}$
- C. $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
- D. $\begin{bmatrix} 0 & i \\ i & 0 \end{bmatrix}$
- E. $\begin{bmatrix} -i & 0 \\ 0 & -i \end{bmatrix}$.

16. Consider the dynamical system $x' = Ax$, where $A = \begin{bmatrix} 1 & 1 \\ 2 & 0 \end{bmatrix}$. Then the origin is

- A. an attractor
- B. a repeller
- C. a saddle point
- D. a spiral point
- E. none of the above

17. Which one of the following is the solution to the differential equation

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 2 & 2 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

with initial condition $\begin{bmatrix} x(0) \\ y(0) \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \end{bmatrix}$?

- A. $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 3e^{4t} \\ 3e^{4t} \end{bmatrix} - \begin{bmatrix} 2e^{-t} \\ 3e^{-t} \end{bmatrix}$
- B. $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 3e^{4t} \\ 3e^{4t} \end{bmatrix} + \begin{bmatrix} 2e^{-t} \\ -3e^{-t} \end{bmatrix}$
- C. $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 2e^{4t} \\ 2e^{4t} \end{bmatrix} + \begin{bmatrix} 2e^{-t} \\ -3e^{-t} \end{bmatrix}$
- D. $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 2e^{4t} \\ 2e^{4t} \end{bmatrix} - \begin{bmatrix} 3e^{-t} \\ -2e^{-t} \end{bmatrix}$
- E. $\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} 2e^{4t} \\ 2e^{4t} \end{bmatrix} + \begin{bmatrix} 3e^{-t} \\ -2e^{-t} \end{bmatrix}$

(i) $\vec{u} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ $\vec{v} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$

\vec{u} & \vec{v} satisfy $2xyz = 0$.

$\vec{u} + \vec{v} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$ & set

(ii) (iii) then (2-y), the nullspace of an m-n matrix is a subspace of \mathbb{R}^n .

18. Which of the following subsets of the vector space \mathbb{R}^3 are subspaces of \mathbb{R}^3 ?

(i) The set of all vectors $v = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ with the property $2xyz = 0$.

(ii) The set of all the solutions of the equation $x - 5y + 2z = 0$.

(iii) The set of all solutions for the system $\begin{bmatrix} 2 & 3 & 2 \\ 5 & 2 & 8 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$.

(iv) The set of all the solutions of the equation $x + 3y = 2z + 1$.

- A. (ii) and (iii) only
- B. (ii) and (iv) only
- C. (iii) and (iv) only
- D. (ii), (iii) and (iv) only
- E. (i), (ii), (iii) and (iv)

(2-8)
 ① 0
 ② the set is closed under +, for any u, v in the set $u+v$ is also in it.
 ③ the set is closed under \cdot , for any u in the set & any $\alpha \in \mathbb{R}$ αu is also in the set.

$\begin{bmatrix} 1 \\ -5 \\ 2 \end{bmatrix}$
 $\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \in \mathbb{R}^3$, $0 \cdot 1 + 3 \cdot 0 \neq 2 \cdot 0 + 1$

19. Determine a basis for the set spanned by the vectors

$$v_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, v_2 = \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}, v_3 = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}, v_4 = \begin{bmatrix} 5 \\ 11 \\ 17 \end{bmatrix}, v_5 = \begin{bmatrix} 2 \\ 7 \\ 12 \end{bmatrix}, v_6 = \begin{bmatrix} 2 \\ 0 \\ 0 \end{bmatrix}$$

- A. $\{v_1, v_3, v_4\}$
- B. $\{v_1, v_3, v_5\}$
- C. $\{v_2, v_3, v_4\}$
- D. $\{v_3, v_4, v_5\}$
- E. $\{v_1, v_3, v_6\}$

20. Performing the Gram-Schmidt process on the vectors $\left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix} \right\}$ yields an orthonormal basis $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ of \mathbb{R}^3 . What is \mathbf{u}_3 ?

A. $\frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$

B. $\frac{1}{\sqrt{26}} \begin{bmatrix} -1 \\ -4 \\ 3 \end{bmatrix}$

C. $\frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$

D. $\frac{1}{\sqrt{14}} \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$

E. $\frac{1}{\sqrt{17}} \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}$

21. Find the least squares solution to

$$\begin{bmatrix} 1 & 3 \\ 2 & 2 \\ 1 & 5 \end{bmatrix} \mathbf{x} = \begin{bmatrix} 0 \\ 5 \\ 8 \end{bmatrix}.$$

A. $(0, 1)$

B. $(1, 1)$

C. $(1, 2)$

D. $(0, 2)$

E. $(2, 1)$

22. Find the distance from the vector \mathbf{y} to the subspace $W = \text{Span}\{\mathbf{u}, \mathbf{v}\}$, where

$$\mathbf{y} = \begin{bmatrix} -1 \\ -5 \\ 10 \end{bmatrix}, \quad \mathbf{u} = \begin{bmatrix} -2 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}.$$

- A. 12.
- B. $2\sqrt{2}$.
- C. $3\sqrt{3}$.
- D. 8.
- E. $3\sqrt{5}$.

23. Let A be an $n \times n$ matrix. Which of the following statements is/are **NOT** equivalent to that A is invertible?

- (i) Columns of A are linearly independent.
- (ii) A is diagonalizable.
- (iii) Columns of A is an orthonormal set.
- (iv) The dimension of the null space of A is 0.
- (v) The linear system $AX = b$ always has solution for any $b \in \mathbb{R}^n$.

- A. (i), (ii) and (iii) only.
- B. (i) and (ii) only.
- C. (ii) and (iii) only.
- D. (i) and (iv) only.
- E. (ii), (iv), (v) only.

24. Let $C[-1, 1]$ be the space of all continuous functions over $[-1, 1]$ with the inner product

$$\langle f(t), g(t) \rangle = \int_{-1}^1 f(t)g(t)dt \quad \text{for any } f(t), g(t) \in C[-1, 1].$$

Which of the following set is an orthogonal basis of $\text{Span}\{1, t - 1, t^2 + t\}$?

- A. $1, t, t^2$
- B. $1, t - 1, t^2 + t$
- C. $1, t, t^2 - 1$
- D. $1, t - 1, t^2$
- E. $1, t, t^2 - \frac{1}{3}$.

25. Suppose that $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} = QDQ^T$ where $D = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}$ and Q is an orthogonal matrix. In the following select a pair of Q and D with required properties.

A. $Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{6}} & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{3}} \end{bmatrix}, D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{bmatrix}.$

B. $Q = \begin{bmatrix} 1 & 1 & 1 \\ -1 & 0 & 1 \\ 0 & -1 & 1 \end{bmatrix}, D = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$

C. $Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{3}} \\ 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{3}} \end{bmatrix}, D = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$

D. $Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{bmatrix}, D = \begin{bmatrix} 2 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}.$

E. $Q = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} \\ 0 & -\frac{2}{\sqrt{6}} & \frac{1}{\sqrt{3}} \end{bmatrix}, D = \begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$